

ON SLICING INVARIANTS OF KNOTS

BRENDAN OWENS

ABSTRACT. The slicing number of a knot, $u_s(K)$, is the minimum number of crossing changes required to convert K to a slice knot. This invariant is bounded above by the unknotting number and below by the slice genus $g_s(K)$. We show that for many knots, previous bounds on unknotting number obtained by Ozsváth and Szabó and by the author in fact give bounds on the slicing number. Livingston defined another invariant $U_s(K)$ which takes into account signs of crossings changed to get a slice knot, and which is bounded above by the slicing number and below by the slice genus. We exhibit an infinite family of knots K_n with slice genus n and Livingston invariant greater than n . Our bounds are based on restrictions (using Donaldson’s diagonalisation theorem or Heegaard Floer homology) on the intersection forms of four-manifolds bounded by the double branched cover of a knot.

1. INTRODUCTION

The unknotting number of a knot is the minimum number of crossing changes required to convert it to an unknot. Ozsváth and Szabó used Heegaard Floer theory to provide a powerful obstruction to a knot having unknotting number one [22]. This obstruction was generalised in [17] to higher unknotting numbers. In this paper we show that similar techniques yield information about the number of crossing changes required to convert to a slice knot.

The slice genus $g_s(K)$ of a knot K in the three-sphere is the minimum genus of a connected oriented smoothly properly embedded surface in the four-ball with boundary K . A knot is called *slice* if $g_s(K) = 0$. Given any diagram D for a knot K , a new knot may be obtained by changing one or more crossings of D . The slicing number $u_s(K)$ is the minimum number of crossing changes required to obtain a slice knot, where the minimum is taken over all diagrams for K . A “movie” of a sequence of crossing changes represents an immersed annulus in $S^3 \times [0, 1]$ with a singularity for each crossing change. A neighbourhood of each singular point may be removed and replaced with an annulus; if the last frame of the movie is a slice knot then the corresponding boundary component of the resulting embedded surface may be capped off with a disk, yielding a surface in B^4 with genus $u_s(K)$ and boundary K .

Recall that crossings in a knot diagram may be given a sign as in Figure 1 (independent of the choice of orientation of the knot). Suppose that K may be *sliced* (converted to a slice knot) by changing p positive and n negative crossings (in some diagram). Form the immersed annulus in $S^3 \times [0, 1]$ as before. The sign of each self-intersection of this annulus agrees with the sign of the corresponding crossing in the changed diagram. Take two self-intersections of opposite sign, and in each case remove a disk neighbourhood of the singularity from just one of the intersecting sheets and connect the boundary components by a tube. This leads to an oriented surface in B^4 with genus

Received by the editors July 9, 2008.

2000 *Mathematics Subject Classification.* 57M25.

B. Owens was supported in part by NSF grant DMS-0604876.




FIGURE 1. Signed crossings in a knot diagram.

$\max(p, n)$. Livingston defined the following slicing invariant:

$$U_s(K) = \min(\max(p, n)),$$

where the minimum is taken over all diagrams for K and over all sets of crossing changes in a diagram which give a slice knot.

From the preceding discussion we see that

$$g_s(K) \leq U_s(K) \leq u_s(K).$$

Livingston showed in [12] that the two-bridge knot $S(15, 4)$, also known as 7_4 , has $u_s = 2$ and $g_s = 1$, thus giving a negative answer to a question of Askitas [16]. (Murakami and Yasuhara showed in [14] that 8_{16} has $u_s = 2$ and $g_s = 1$. Their proof is based on a four-manifold bounded by the double-branched cover of the knot. We take a similar approach here.) Livingston also asked whether in fact U_s is always equal to the slice genus, and suggested that 7_4 may be a counterexample.

Let $\sigma(K)$ denote the signature of a knot K . It is shown in [3, Proposition 2.1] (also [24, Theorem 5.1]) that if K' is obtained from K by changing a positive crossing, then

$$\sigma(K') \in \{\sigma(K), \sigma(K) + 2\};$$

similarly if K' is obtained from K by changing a negative crossing then

$$\sigma(K') \in \{\sigma(K), \sigma(K) - 2\}.$$

Now suppose that K may be sliced by changing p positive and n negative crossings (in some diagram). Since a slice knot has zero signature, it follows that a bound for n is given by

$$(1) \quad n \geq \sigma(K)/2.$$

In this paper we give an obstruction to equality in (1).

Let $\Sigma(K)$ denote the double cover of S^3 branched along K , and suppose that crossing changes in some diagram for K result in a slice knot J . It follows from ‘‘Montesinos’ trick’’ ([13], or see [17]) that $\Sigma(K)$ is given by Dehn surgery on some framed link in $\Sigma(J)$ with half-integral framing coefficients.

Definition 1. *An integer-valued symmetric bilinear form Q on a free abelian group of rank $2r$ is said to be of half-integer surgery type if it admits a basis $\{x_1, \dots, x_r, y_1, \dots, y_r\}$ with*

$$\begin{aligned} Q(x_i, x_j) &= 2\delta_{ij}, \\ Q(x_i, y_j) &= \delta_{ij}. \end{aligned}$$

Examples. The positive-definite rank 2 unimodular form is of half-integer surgery type since it may be represented by the matrix $\begin{pmatrix} 2 & 1 \\ 1 & 1 \end{pmatrix}$. The form represented by the matrix $\begin{pmatrix} 4 & 1 \\ 1 & 4 \end{pmatrix}$ is not of half-integer surgery type since it has no vectors of square 2 (the vector x with coefficients (a, b) has square $4a^2 + 2ab + 4b^2 = (a+b)^2 + 3(a^2 + b^2)$; since a and b are integers one cannot solve $Q(x, x) = 2$).

Converting the half-integer surgery description above to integer surgery in the standard way gives a cobordism W from $\Sigma(J)$ to $\Sigma(K)$ whose intersection form Q_W is of half-integer surgery type. Since J is slice, $\Sigma(J)$ bounds a rational homology ball B . Joining B to W along $\Sigma(J)$ gives a smooth closed four-manifold X bounded by $\Sigma(K)$. The second Betti number of X is twice the number of crossing changes used to get from K to J .

Suppose now that K is converted to J by changing p positive and n negative crossings, with $n = \sigma(K)/2$. Then K bounds a disk in $B^4 \#^{p+n} \mathbb{C}\mathbb{P}^2$. As in [3] (see also [17]) we may take the double cover X' of the blown-up four-ball branched along this disk. It follows from a theorem of Cochran and Lickorish [3, Theorem 3.7] that X' is positive-definite with $b_2(X') = 2(p+n)$. The following theorem is based on the idea that in fact X is diffeomorphic to X' .

Theorem 2. *Suppose that a knot K can be converted to a slice knot by changing p positive and n negative crossings, with $n = \sigma(K)/2$. Then the branched double cover $\Sigma(K)$ bounds a positive-definite smooth four-manifold X with $b_2(X) = 2(p+n)$ whose intersection form Q_X is of half-integer surgery type, with exactly n of $Q_X(x_1, x_1), \dots, Q_X(x_{p+n}, x_{p+n})$ even, and $\det Q_X$ divides $\det K$ with quotient a square.*

For knots whose determinant is square-free, it follows that the first two parts of Ozsváth and Szabó's obstruction to unknotting number one [22, Theorem 1.1] (without the symmetry condition) in fact give an obstruction to $u_s(K) = 1$.

Corollary 3. *The knots*

$$7_4, 8_{16}, 9_5, 9_{15}, 9_{17}, 9_{31}, 10_{19}, 10_{20}, 10_{24}, 10_{36}, 10_{68}, 10_{69}, 10_{86}, \\ 10_{97}, 10_{105}, 10_{109}, 10_{116}, 10_{121}, 10_{122}, 10_{144}, 10_{163}, 10_{165}$$

have slice genus 1 and slicing number 2.

(Note that as mentioned above this was shown for 7_4 in [12] and for 8_{16} in [14]. The slice genus information in Corollaries 3 and 4 is taken from [1].)

Furthermore, the obstruction given in [17, Theorem 5] to unknotting a knot by changing p positive and $n = \sigma(K)/2$ negative crossings is in fact an obstruction to slicing, provided again that $\det K$ is square-free.

Corollary 4. *The knots*

$$9_{10}, 9_{13}, 9_{38}, 10_{53}, 10_{101}, 10_{120}$$

have slice genus 2 and slicing number 3.

The 11-crossing two-bridge knot $S(51, 35)$ (Dowker-Thistlethwaite name 11a365) has slice genus 3 and slicing number 4.

It may also be shown that for some of the knots in Corollaries 3 and 4, Livingston's invariant U_s is not equal to the slice genus. The knot 7_4 is such an example (as Livingston suggested in [12]), and in fact we find that it is the first member of an infinite family of such examples.

Corollary 5. *For each positive integer n , there exists a two-bridge knot K_n with signature $2n$ and slice genus n which cannot be sliced by changing n negative crossings and any number of positive crossings; hence $U_s(K_n) > n$.*

Acknowledgements. It is a pleasure to thank András Stipsicz and Tom Mark for helpful conversations.

2. PROOF OF THEOREM 2

In this section we prove our main result.

Recall that a positive-definite integer-valued symmetric bilinear form Q on a free abelian group A gives an integer lattice L in Euclidean space on tensoring with \mathbb{R} . We say a lattice L in \mathbb{R}^n is of half-integer surgery type if the corresponding form Q is of the form given in Definition 1. Also a matrix representative for Q is referred to as a Gram matrix for L . For convenience we will frequently denote $Q(x, y)$ by $x \cdot y$, and $Q(x, x)$ by x^2 .

The proof of Theorem 2 consists of a topological and an algebraic step. Following [17] we show using careful analysis of Montesinos' trick that, under the hypotheses of the theorem, $\Sigma(K)$ bounds a positive-definite manifold X and that the intersection pairing of X is of half-integer surgery type when restricted to some finite index sublattice. We then show that if a lattice M has an odd index sublattice L of half-integer surgery type then in fact M is of half-integer surgery type.

We begin with a couple of lemmas.

Lemma 2.1. *Let Q be a form of half-integer surgery type, with $m_i = Q(y_i, y_i)$. Then*

$$\det Q \equiv \prod_{i=1}^r (2m_i - 1) \pmod{4}.$$

Proof. This follows from the discussion after Lemma 2.2 in [17]. □

Lemma 2.2. *Let Q be a block matrix of $r \times r$ blocks of the form $\begin{pmatrix} 2I & * \\ * & * \end{pmatrix}$ which is congruent modulo 2 to $\begin{pmatrix} 2I & I \\ I & X \end{pmatrix}$. Then there exists $P = \begin{pmatrix} I & * \\ 0 & R \end{pmatrix} \in GL(2r, \mathbb{Z})$ with*

$$P^T Q P = \begin{pmatrix} 2I & I \\ I & X' \end{pmatrix},$$

and $X' \equiv X \pmod{2}$.

Proof. Let Q be the Gram matrix of a lattice with basis $x_1, \dots, x_r, z_1, \dots, z_r$. By successively adding multiples of x_i to each of z_1, \dots, z_r we get a new basis $x_1, \dots, x_r, y_1, \dots, y_r$ with $x_i \cdot y_j = \delta_{ij}$; since x_i has even square, this preserves parities on the diagonal. □

The following was originally proved by Ozsváth and Szabó [22] in the case $p + n = 1$ and J is the unknot.

Proposition 2.3. *Suppose that a knot K can be converted to a slice knot J by changing p positive and n negative crossings, with $n = \sigma(K)/2$. Then the branched double cover $\Sigma(K)$ bounds a positive-definite four-manifold X with $b_2(X) = 2(p + n)$. The lattice $(H_2(X; \mathbb{Z}), Q_X)$ contains a finite index sublattice of half-integer type, which has a basis as in Definition 1 with exactly p elements of odd square.*

Proof. We adapt the proof of [17, Lemma 3.2]. By Montesinos' lemma ([13], or see [17, Lemma 3.1]), $\Sigma(J)$ is the result of surgery on some link L in S^3 with half-integer framing coefficients. Convert to integer surgery (see [6] or [17, Lemma 2.2]): the framing $(2m - 1)/2$ on any component is replaced with m , and a small 2-framed meridian is added to each component. Let Q_J be the resulting linking matrix of half-integer type. We may assume (after possibly adding a $-1/2$ framed unknot to L) that $\det Q_J$ is positive. This integer framed link is a Kirby diagram for a four-manifold X_J obtained by

attaching a two-handle to the four-ball along each framed link component, i.e. a four-dimensional two-handlebody. The two-handlebody X_J has boundary Σ_J and intersection form represented by Q_J .

Note that since J is slice it has signature zero and determinant $\det J = \det Q_J = k^2$ for some odd integer k . Suppose that K_- is a knot of signature 2 which may be converted to J by changing a single negative crossing c . Then $\Sigma(K)$ is the result of surgery on $L \cup C$ for some knot C in S^3 , with framing $(2m - 1)/2$ on C . Let K_0 be the result of taking the oriented resolution of the crossing c ; then as in [17, Lemma 3.2] we have that $\Sigma(K_0)$ is surgery on $L \cup C$ with framing m on C . Converting to integer surgery we find that $\Sigma(K_-), \Sigma(K_0)$ are given by integer surgeries with linking matrices

$$Q_- = \begin{pmatrix} 2 & 1 & 0 \\ 1 & m & * \\ 0 & * & Q_J \end{pmatrix}, \quad Q_0 = \begin{pmatrix} m & * \\ * & Q_J \end{pmatrix}.$$

Let $\Delta_L(t)$ denote the Conway-normalised Alexander polynomial of a link L (see for example [9, Chapter 8]). For a knot K , the sign of the Alexander polynomial at -1 is given by

$$\Delta_K(-1) = (-1)^{\sigma(K)/2} \det K.$$

(This may be seen by diagonalising the symmetrised Seifert form as in [15, Proof of Theorem 5.6].) This shows that

$$\Delta_J(-1) = k^2, \quad \Delta_{K_-}(-1) = -|\det Q_-| = -|2 \det Q_0 - k^2|.$$

The skein relation for the Alexander polynomial then yields

$$k^2 + |2 \det Q_0 - k^2| = 2|\det Q_0|,$$

from which we conclude that $\det Q_- = 2 \det Q_0 - k^2$ is positive. Now using Lemma 2.1 we have

$$2m - 1 \equiv \det Q_- = \det K_- \equiv 3 \pmod{4},$$

and thus m is even. (The last congruence is due to Murasugi [15]: $\det K \equiv \sigma(K) + 1 \pmod{4}$.)

Now suppose K_+ is a knot of signature 0 which may be converted to J by changing a positive crossing. Again $\Sigma(K)$ is obtained by half-integer surgery on $L \cup C$ for some knot C in S^3 , with framing $(2m - 1)/2$ on C . Let Q_+ denote the linking matrix after converting to integer surgery. A similar argument as above (see [17] for the case where J is the unknot) shows that $\det Q_+ > 0$ and m is odd.

Let c_1, \dots, c_{p+n} be the set of crossings (p positive, n negative) in some chosen diagram of K that we change to convert to J . Then $\Sigma(K)$ is Dehn surgery on the link $L \cup C_1 \cup \dots \cup C_{p+n}$, with half-integer framing coefficients. Each C_i corresponds to a crossing c_i . Dehn surgery on L union a sublink of $C_1 \cup \dots \cup C_{p+n}$ gives the double branched cover of a knot which is obtained from K by changing a subset of the crossings c_1, \dots, c_{p+n} . In particular surgery on the knot $L \cup C_i$ yields the double branched cover of the knot K'_i which is obtained from K by changing all of the crossings except c_i . By the condition $n = \sigma(K)/2$ the knot signature changes every time a negative crossing is changed and remains constant when a positive crossing changes. It follows from the discussion above applied to K'_i that the framing on each C_i is of the form $(2m_i - 1)/2$ and exactly those m_i which correspond to changing negative crossings of K are even.

Denote by X_K the two-handlebody with boundary $\Sigma(K)$ that results from converting to integer surgery (i.e. surgery on the link $L \cup C_1 \cup \dots \cup C_{p+n}$, followed by surgery on a 2-framed meridian of each component). Then X_K has intersection form of half-integer surgery type; moreover we can view X_K as the union of X_J and a surgery cobordism W along the common boundary $\Sigma(J)$. We will show by an induction argument that this cobordism is positive-definite.

Let K_j denote the knot obtained from K by changing crossings c_{j+1}, \dots, c_{p+n} , and let Q_j be the linking matrix of half-integer type obtained by converting the corresponding Dehn surgery diagram of $\Sigma(K_j)$ to integer type.

Suppose that $\det Q_{j-1}$ is positive and hence equal to $\det K_{j-1}$. We have

$$Q_j = \begin{pmatrix} 2 & 1 & 0 \\ 1 & m_j & * \\ 0 & * & Q_{j-1} \end{pmatrix},$$

and by Lemma 2.1

$$\det Q_j \equiv (2m_j - 1) \det Q_{j-1} \pmod{4}.$$

If c_j is a positive crossing then m_j is odd and so

$$(2) \quad \det Q_j \equiv \det Q_{j-1} \pmod{4}.$$

On the other hand the signature of K_{j-1} is equal to that of K_j and so

$$(3) \quad \det K_j \equiv \det K_{j-1} \pmod{4}.$$

Comparing (2) and (3) we see that $\det Q_j$ is congruent modulo 4 to its absolute value. Since it is an odd number it must be positive.

On the other hand if c_j is negative we find both congruences (2) and (3) do *not* hold, and again it follows that $\det Q_j$ is positive.

By induction we see that $Q_J = Q_0, Q_1, \dots, Q_{p+n}$ all have positive determinants. Thus the surgery cobordism W is built by attaching $2(p+n)$ two-handles to the two-handlebody X_J , and before and after each attachment we have a four-manifold whose intersection pairing has positive determinant. It follows that W is positive-definite.

We claim $(H_2(W; \mathbb{Z}), Q_W)$ contains a finite index sublattice with a basis as in Definition 1 with exactly p elements of odd square. Suppose that L has r components, so that $b_2(X_K) = 2(p+n+r)$. Let $\{x_i, y_i\}$ be a basis for $H_2(X_K; \mathbb{Z})$ as in Definition 1, chosen so that $\{x_i, y_i\}_{i>p+n}$ is a basis for the sublattice $H_2(X_J; \mathbb{Z})$. For $i \leq p+n$, take y_i to be the class corresponding to the two-handle attached along C_i , and x_i that corresponding to the two-handle attached along the meridian of C_i . Note that each x_i is contained in $H_2(W; \mathbb{Z})$. The rough idea is to form a sublattice by projecting the span of $\{x_i, y_i\}_{i \leq p+n}$ orthogonally to $H_2(W; \mathbb{Z})$. Of course we cannot quite do this; however since $H_1(\Sigma(J); \mathbb{Z})$ has order k^2 we may write

$$k^2 y_i = z_i + w_i, \quad i = 1, \dots, p+n$$

with $z_i \in H_2(W; \mathbb{Z})$ and $w_i \in H_2(X_J; \mathbb{Z})$. We claim that the self-intersection of z_i has the same parity as that of y_i . To see this note that for each $j > p+n$, y_i is orthogonal to x_j and hence so is w_i . It follows that w_i is in the span of $\{x_i\}_{i>p+n}$ and so has even self-intersection. Thus we have a full rank sublattice of $(H_2(W; \mathbb{Z}), Q_W)$ with basis $\{x_i, z_i\}_{i \leq p+n}$; by Lemma 2.2, this sublattice has a basis as in Definition 1 with exactly p elements of odd square.

Form the manifold X with one boundary component by capping off the $\Sigma(J)$ end of W with the rational ball B given as the double branched cover of B^4 along a slice disk bounded by J [2]. Then $(H_2(W; \mathbb{Z}), Q_W)$ is a sublattice of $(H_2(X; \mathbb{Z}), Q_X)$, and therefore so is $\text{Span}(\{x_i, z_i\}_{i \leq p+n})$. \square

Proposition 2.4. *Suppose M is a positive-definite integer lattice of rank $2r$ and L is a sublattice of M of odd index l . If L is of half-integer type then so is M . Moreover, the number of elements of a basis of M as in Definition 1 with odd square is the same as that for L .*

This will follow from the next two lemmas, the first of which is standard.

Lemma 2.5. *The natural map from $GL(n, \mathbb{Z})$ to $GL(n, \mathbb{Z}/2)$ is onto for any n .*

Proof. Use induction on n . Suppose that $R \in M(n, \mathbb{Z})$ has odd determinant. The cofactor expansion across the first row yields

$$\det R = r_{11}R_{11} + r_{12}R_{12} + \cdots + r_{1n}R_{1n}.$$

Since the determinant is odd, so is at least one $r_{1j}R_{1j}$. By induction we may choose $\tilde{R} \equiv R \pmod{2}$ with $\tilde{R}_{1j} = 1$, then adjust the value of r_{1j} to get $\det \tilde{R} = 1$. \square

Lemma 2.6. *Suppose M is a positive-definite integer lattice of rank $2r$, and that L is a lattice of half-integer type which is a sublattice of M of odd index l . Let $x_1, \dots, x_r, y_1, \dots, y_r$ be a basis for L as in Definition 1, and let Q_L be the Gram matrix of L in this basis. Then x_1, \dots, x_r may be extended to a basis $x_1, \dots, x_r, z_1, \dots, z_r$ for M with*

$$Q_M \equiv Q_L \pmod{2}.$$

Proof. Let $m_i = y_i \cdot y_i$. In the given basis Q_L is in block form $\begin{pmatrix} 2I & I \\ I & X \end{pmatrix}$, with $\text{Diag}(X) = (m_1, \dots, m_r)$. By Theorem 6 in Chapter 1, Section 3 of [7] there exists a basis $a_1, \dots, a_r, z_1, \dots, z_r$ of M with $x_i \in \text{Span}_{\mathbb{Z}}\{a_1, \dots, a_i\}$. A simple induction argument using the fact that $x_i \cdot y_j = \delta_{ij}$ shows that in fact (after possibly multiplying by -1) we have $a_i = x_i$ for $i = 1, \dots, r$.

Let $P \in M(2r, \mathbb{Z})$ be the matrix whose i th column is the coefficient vector of the i th basis vector of L in the basis $x_1, \dots, x_r, z_1, \dots, z_r$. Then P is in block form $\begin{pmatrix} I & * \\ 0 & R \end{pmatrix}$, and

$$Q_L = P^T Q_M P.$$

Since $\det Q_L = l^2 \det Q_M$ we have $\det R = l$ is odd. By Lemma 2.5 we may choose $\tilde{R} \in GL(r, \mathbb{Z})$ with $\tilde{R} \equiv R \pmod{2}$. Applying the transition matrix $\tilde{P} = \begin{pmatrix} I & * \\ 0 & \tilde{R} \end{pmatrix}$ to M yields the required basis. \square

Proof of Proposition 2.4. Let $x_1, \dots, x_r, z_1, \dots, z_r$ be the basis of M given by Lemma 2.6, in which

$$Q_M = \begin{pmatrix} 2I & * \\ * & * \end{pmatrix} \equiv Q_L = \begin{pmatrix} 2I & I \\ I & X \end{pmatrix} \pmod{2}.$$

The proposition now follows from Lemma 2.2.

Proof of Theorem 2. Let X be the four-manifold bounded by $\Sigma(K)$ given by Proposition 2.3. A standard argument using the long exact homology sequence of the pair $(X, \Sigma(K))$ shows that $\det Q_X$ divides $\det K$ with quotient a square (see for example [18, Lemma 2.1]). The determinant of a knot is always odd; it follows that Q_X has odd order and the sublattice given by Proposition 2.3 has odd index in $(H_2(X; \mathbb{Z}), Q_X)$. Theorem 2 now follows immediately from Proposition 2.4.

3. EXAMPLES

Theorem 2 tells us that to show that a knot K cannot be converted to a slice knot by changing p positive and $n = \sigma(K)/2$ negative crossings, we must show that its double branched cover cannot bound a four-manifold with an intersection form with certain properties.

We will make use of two very effective gauge-theoretic obstructions to a rational homology three-sphere Y bounding a positive-definite form Q . On the one hand, Ozsváth and Szabó define a function

$$d : \text{Spin}^c(Y) \rightarrow \mathbb{Q}$$

coming from the absolute grading in Heegaard Floer homology, and they show that for each spin^c structure \mathfrak{s} on a positive-definite four manifold bounded by Y the following must hold:

$$(4) \quad c_1(\mathfrak{s})^2 - b_2(X) \geq 4d(\mathfrak{s}|_Y),$$

$$(5) \quad \text{and } c_1(\mathfrak{s})^2 - b_2(X) \equiv 4d(\mathfrak{s}|_Y) \pmod{2}.$$

The left hand side depends on the intersection form of X .

To determine if a given knot K may be sliced by changing p positive and $n = \sigma(K)/2$ negative crossings we carry out the following steps:

1. Compute $d : \Sigma(K) \rightarrow \mathbb{Q}$;
2. Find a complete finite set of representatives Q_1, \dots, Q_m of forms of rank $2(p+n)$ satisfying the conclusion of Theorem 2;
3. Check using (4,5) whether $\Sigma(K)$ is obstructed from bounding each of Q_1, \dots, Q_m .

Details on Heegaard Floer theory and the d invariant may be found in [20, 21, 22]; for a summary of how this theory may be used in our context see [17].

Another approach to understanding the set of positive-definite forms that a three-manifold Y may bound is to make use of Donaldson's diagonalisation theorem [4]. This approach works well for Seifert fibred rational homology spheres, and especially for lens spaces (see e.g. [10], [11], [19]). Knowledge of a particular negative-definite four-manifold X_1 bounded by Y constrains the intersection form of a positive-definite X_2 with the same boundary, since $X = X_2 \cup -X_1$ is a closed positive-definite manifold with $(H_2(X; \mathbb{Z}), Q_X) \cong \mathbb{Z}^m$ for some m . We will illustrate this technique in Subsection 3.3.

The slicing number of a knot is the same as that for its reflection. We assume in what follows that all knots have nonnegative signature. (This distinguishes between the knot and its reflection unless the signature is zero.)

3.1. Knots with slice genus one. For a knot with $\sigma(K) = 2$ and $u_s(K) = 1$, it follows from inequality (1) and Theorem 2 that $\Sigma(K)$ bounds a four-manifold whose intersection form is represented by the matrix

$$\begin{pmatrix} m & 1 \\ 1 & 2 \end{pmatrix},$$

with $\det K = (2m-1)t^2$ for some integer t .

For a knot with $\sigma(K) = 0$ and $u_s(K) = 1$ we find that either $\Sigma(K)$ or $-\Sigma(K)$ must bound such a positive-definite four-manifold.

The knots listed in Corollary 3 have square-free determinant. For each of them, Ozsváth and Szabó have shown in [22], using (4,5), that $\pm\Sigma(K)$ cannot bound

$$\begin{pmatrix} \frac{\det K + 1}{2} & 1 \\ 1 & 2 \end{pmatrix};$$

it is also known in each case that the knot can be unknotted with two crossing changes. We conclude that $u_s(K) = 2$.

Remark 3.1. *Each of the knots 10_{29} , 10_{40} , 10_{65} , 10_{67} , 10_{89} , 10_{106} , and 10_{108} has signature 2 and $\det K = st^2$ for some $t > 1$. In each case Ozsváth and Szabó have shown $\Sigma(K)$ cannot bound $\begin{pmatrix} \frac{\det K + 1}{2} & 1 \\ 1 & 2 \end{pmatrix}$ and hence K has unknotting number two.*

However we find in each case $\Sigma(K)$ is not obstructed from bounding $\begin{pmatrix} \frac{s+1}{2} & 1 \\ 1 & 2 \end{pmatrix}$.

3.2. Knots with slice genus two or three. We now consider the knots in Corollary 4. Each of $9_{10}, 9_{13}, 9_{38}, 10_{53}, 10_{101}, 10_{120}$ has signature 4 and slice genus 2. In [17] we have shown, using (4, 5), that in each case $\Sigma(K)$ cannot bound a positive-definite form Q as in Definition 1 with rank 4, $\det Q = \det K$ and $Q(x_i, x_i)$ even. Since the knots have square-free determinant, it follows from Theorem 2 that they cannot be sliced with two crossing changes. Each can be unknotted with three crossing changes, and so in each case $u_s(K) = 3$.

Similarly $K = 11a365$ is shown in [17] to have unknotting number 4, and since its determinant is square-free the same argument shows it has $u_s(K) = 4$.

3.3. Knots with large slice genus. In this subsection we prove Corollary 5. We define K_n to be the 4-plat closure of the four-string braid $(\sigma_1^4 \sigma_2^4)^n$, as illustrated in Figure 2 for $n = 2$. For $n = 1$ this is the knot $7_4 = S(15, 11)$ shown by Lickorish to have unknotting number 2 and by Livingston to have slicing number 2 [8, 12].




FIGURE 2. **The knot K_2 .** The two pairs of dashed arcs indicate where to attach ribbons to go from K_n to K_{n-1} .

As illustrated in the diagram, two oriented ribbon moves convert K_n to K_{n-1} . Since K_0 is the unknot this shows that the slice genus of K_n is at most n . The signature of K_n may be shown (see below) to be $2n$. We conclude that

$$g_s(K_n) = n.$$

Let $P(a_1, a_2, \dots, a_m)$ denote the plumbing of disk bundles over two-spheres corresponding to the linear graph with m vertices, where the i th vertex has weight a_i . The double cover of S^3 branched along K_n is the boundary of $P(4, 4, \dots, 4)$. Let Q_n denote the rank $2n$ intersection form of this plumbing, and let L_n denote the associated lattice in \mathbb{R}^{2n} . The following sequence of blow-ups and blow-downs exhibits $\Sigma(K_n)$ as the boundary of a negative-definite plumbing:

$$\begin{aligned} \Sigma(K_n) &\cong \partial P(4, 4, \dots, 4) \\ &\cong \partial P(-1, 2, -1, 2, \dots, -1, 2, -1) \\ &\cong \partial P(-2, -1, 1, -2, -1, 1, \dots, -2, -1, 1, -1) \\ &\cong \partial P(-2, -2, -3, -2, -3, -2, \dots, -3, -2, -3, -2, -2). \end{aligned}$$

(We note the above shows that K_n may also be represented by the alternating diagram which is the closure of the braid

$$(\sigma_1^{-1} \sigma_2^2)^{2n} \sigma_1^{-1};$$

using the formula of Gordon and Litherland [5] it is easy to compute the signature from this diagram.)

Let X'_n denote the positive-definite plumbing $P(2, 2, 3, 2, 3, 2, \dots, 3, 2, 2)$ with boundary $-\Sigma(K_n)$. Let Q'_n denote its intersection form and let L'_n denote the associated lattice. Note that L'_n has dimension $2n + 3$: there are n vertices with weight 3 and $n + 3$ with weight 2.

Lemma 3.2. *Suppose $\Sigma(K_n)$ is given as the boundary of a smooth four-manifold X with positive-definite intersection form Q_X . Then $(H_2(X; \mathbb{Z}), Q_X)$ embeds as a finite index sublattice of $L_n \oplus \mathbb{Z}^k$ for some $k \geq 0$.*

Proof. Gluing X to X'_n along their boundary gives a closed positive-definite manifold. It follows from Donaldson's theorem that the lattice L'_n embeds as a sublattice of \mathbb{Z}^m with the lattice $(H_2(X; \mathbb{Z}), Q_X)$ contained in its orthogonal complement.

Let e_1, \dots, e_m be an orthonormal basis of \mathbb{Z}^m . Up to automorphism of \mathbb{Z}^m there is a unique way to embed L'_n : the first vertex vector must map to $e_1 + e_2$, the second to $e_2 + e_3$, the third to $e_3 + e_4 + e_5$, the fourth to $e_5 + e_6$ and so on. Thus the image of L'_n is contained in a \mathbb{Z}^{3n+4} sublattice of \mathbb{Z}^m . An easy calculation shows that the orthogonal complement of L'_n in \mathbb{Z}^{3n+4} is spanned by the vectors $e_1 - e_2 + e_3 - e_4, e_4 - e_5 + e_6 - e_7, \dots, e_{3n+1} - e_{3n+2} + e_{3n+3} - e_{3n+4}$. These span a copy of L_n , from which the conclusion follows. \square

Lemma 3.3. *For any $n \geq 1, k \geq 0$, the lattice $L_n \oplus \mathbb{Z}^k$ does not admit any finite index sublattices of half-integer surgery type.*

Proof. For $k = 0$ this is immediate since L_n has no nonzero vectors of square less than 4. If $k > 0$ let e_1, \dots, e_k be an orthonormal basis of \mathbb{Z}^k , and suppose we have a sublattice of $L_n \oplus \mathbb{Z}^k$ of half-integer surgery type with basis $\{x_i, y_i\}$ as in Definition 1. Up to an automorphism of $L_n \oplus \mathbb{Z}^k$ we have $x_1 = e_1 + e_2$. Then x_2 is orthogonal to x_1 . We cannot have $x_2 = e_1 - e_2$ since y_1 pairs evenly with x_2 and oddly with x_1 . Thus up to automorphism, $x_2 = e_3 + e_4$. It follows that any sublattice of $L_n \oplus \mathbb{Z}^k$ of half-integer surgery type has rank at most k . \square

Corollary 5 now follows from Lemmas 3.2 and 3.3 and Theorem 2.

Remark 3.4. *Livingston conjectured in [12] that the difference $U_s - g_s$ can be arbitrarily large. It is possible to unknot K_n by changing $2n$ positive crossings in the diagram as in Figure 2. In the absence of any further evidence it is tempting to conjecture that for these knots $U_s - g_s = n$; in any case this would seem to be a good candidate with which to attempt to verify Livingston's conjecture.*

Remark 3.5. *The trace of a homotopy from a knot K to a slice knot J is an immersed annulus in $S^3 \times I$; capping this off with a slice disk yields an immersed disk D in B^4 bounded by K . If J is obtained from K by changing p positive and n negative crossings then the resulting disk D has p negative self-intersections and n positive self-intersections. (This is why changing a positive crossing is often referred to as a "negative crossing change".) Instead of considering crossing changes one may ask whether a knot K bounds an immersed disk in B^4 with a prescribed number of self-intersections, or with prescribed numbers of self-intersections of each sign.*

Rudolph has shown in [23] that the minimal number of self-intersections in a ribbon immersed disk bounded by K is equal to the minimal number of crossing changes to get from K to a ribbon knot. (Here a ribbon surface in B^4 is one on which the radial distance function has no maxima, and a ribbon knot is a knot which bounds an embedded ribbon disk.) Knowing whether a result analogous to Rudolph's for non-ribbon disks and slice knots holds would be very interesting. It is to be expected that the conclusion of Theorem 2 holds under the weaker hypothesis that K bounds an immersed disk with p negative, $n = \sigma(K)/2$ positive self-intersections. The expected proof would generalise that of [14] and also make use of [3, Theorem 3.7].

REFERENCES

- [1] J. C. Cha & C. Livingston, *Table of knot invariants*, <http://www.indiana.edu/~knotinfo>.
- [2] A. J. Casson & C. McA. Gordon, *On slice knots in dimension three*, Algebraic and geometric topology, Part 2, Proc. Sympos. Pure Math. **XXXII**, Amer. Math. Soc., 1978, 39–53.
- [3] T. D. Cochran & W. B. R. Lickorish, *Unknotting information from 4-manifolds*, Trans. Amer. Math. Soc. **297** 1986, 125–142.
- [4] S. K. Donaldson, *An application of gauge theory to four-dimensional topology*, J. Diff. Geom. **18** 1983, 279–315.
- [5] C. McA. Gordon & R. A. Litherland, *On the signature of a link*, Invent. Math. **47** 1978, 53–69.
- [6] R. E. Gompf & A. I. Stipsicz, *4-manifolds and Kirby calculus*, Graduate Studies in Math. **20**, Amer. Math. Soc., 1999.
- [7] P. M. Gruber & C. G. Lekkerkerker, *Geometry of numbers, 2nd ed.*, North-Holland Mathematical Library **37**, North-Holland, 1987.
- [8] W. B. R. Lickorish, *The unknotting number of a classical knot*, Contemp. Math. **44** 1985, 117–121.
- [9] W. B. R. Lickorish, *An introduction to knot theory*, Graduate Texts in Math. **175**, Springer, 1997.
- [10] P. Lisca, *Symplectic fillings and positive scalar curvature*, Geometry and Topology **2** 1998, 103–116.
- [11] P. Lisca, *Lens spaces, rational balls and the ribbon conjecture*, Geometry and Topology **11** 2007, 429–472.
- [12] C. Livingston, *The slicing number of a knot*, Algebraic and Geometric Topology **2** 2002, 1051–1060.
- [13] J. M. Montesinos, *Varietades de Seifert que son cubiertas ciclicas ramificados de dos hojas*, Bol. Soc. Mat. Mexicana **18** (1973), 1–32.
- [14] H. Murakami & A. Yasuhara, *Four-genus and four-dimensional clasp number of a knot*, Proc. Amer. Math. Soc. **128** 2000, 3693–3699.
- [15] K. Murasugi, *On a certain numerical invariant of link types*, Trans. Amer. Math. Soc. **117** 1965, 387–422.
- [16] T. Ohtsuki, *Problems on invariants of knots and 3-manifolds*, Invariants of knots and 3-manifolds (Kyoto 2001), Geometry and Topology Monographs **4** 2002, 377–572.
- [17] B. Owens, *Unknotting information from Heegaard Floer homology*, Advances in Mathematics, **217** (2008), pages 2353–2376.
- [18] B. Owens & S. Strle, *Rational homology spheres and the four-ball genus of knots*, Advances in Mathematics, **200** (2006), pages 196–216.
- [19] B. Owens & S. Strle, *A characterisation of the $n\langle 1 \rangle \oplus \langle 3 \rangle$ form and applications to rational homology spheres*, Math. Res. Lett., **13** (2006), pages 259–271.
- [20] P. Ozsváth & Z. Szabó, *Absolutely graded Floer homologies and intersection forms for four-manifolds with boundary*, Advances in Mathematics **173** 2003, 179–261.
- [21] P. Ozsváth & Z. Szabó, *On the Floer homology of plumbed three-manifolds*, Geometry and Topology **7** 2003, 225–254.
- [22] P. Ozsváth & Z. Szabó, *Knots with unknotting number one and Heegaard Floer homology*, Topology **44** 2005, 705–745.
- [23] L. Rudolph, *Braided surfaces and Seifert ribbons for closed braids*, Comment. Math. Helv. **58** 1983, 1–37.
- [24] A. Stoimenow, *Polynomial values, the linking form and unknotting numbers*, Math. Res. Lett. **11** 2004, 755–769.

DEPARTMENT OF MATHEMATICS, LOUISIANA STATE UNIVERSITY, BATON ROUGE, LA 70803

Current address: Department of Mathematics, University of Glasgow, Glasgow G12 8QW, UK