

Notes on Hopf algebras over fields

Notes for SMSTC supplementary course 2024–5

Andrew Baker

University of Glasgow

© Andrew Baker Last updated 29/03/2025

Introduction

These notes are intended to cover some basic ideas and results on Hopf algebras over fields, especially finite dimensional ones.

Throughout, \mathbb{k} will be a field, although much of the theory works over a general commutative ring provided due care is taken over flatness and finiteness conditions.

We will make use of basic category theory terminology some of which will be explained as it arises. The books by Mac Lane and Riehl [ML98, Rie16] are good sources for this.

The References contain several books and expository articles that cover aspects of the theory that will be covered in these notes. Radford's book [Rad12] is probably the most complete source for the general theory of Hopf algebras, while Montgomery [Mon93] is more terse but extremely useful. The recent book by Cartier & Patras [CP21] covers examples from areas such as combinatorics and is a good introduction to the 'classical' theory. The lecture notes by Brown & Goodearl [BG02] are wide ranging although their main focus is quantum groups. Lorentz [Lor18] is an amazing book which contains a lot on Hopf algebras. Waterhouse [Wat79] is a very accessible introduction to group schemes and the functorial viewpoint in Algebraic Geometry, while Milne [Mil17] is more exhaustive and recent. The classic Milnor & Moore [MM65] was one of the earliest accounts of the theory of \mathbb{Z} -graded Hopf algebras and is still an important source for topologists and there is an updated presentation of some aspects by May & Ponto [MP12].

Contents

Introduction	2
Chapter 1. Some background material	5
1.1. Some category theory odds and ends	5
1.2. Vector spaces over a field as a symmetric monoidal category	11
1.3. Graded vector spaces	15
Chapter 2. Algebras, coalgebras, bialgebras and Hopf algebras	17
2.1. Algebras	17
2.2. Coalgebras	20
2.3. Dualising between coalgebras and algebras	23
2.4. Bialgebras	24
2.5. Interlude: Convolution monoids	25
2.6. Hopf algebras	28
2.7. SubHopf algebras, adjoint actions and normal subalgebras	32
Chapter 3. A cornucopia of examples	37
3.1. Endomorphism algebras of vector spaces	37
3.2. Polynomial algebras and their duals	37
3.3. The free vector space	38
3.4. Group algebras and dual group algebras	39
3.5. Poset coalgebras and algebras	40
3.6. Free algebras, bialgebras and Hopf algebras	40
3.7. Free bialgebras and free Hopf algebras	41
3.8. Enveloping algebras of Lie algebras	42
3.9. Restricted Lie algebras	43
3.10. Affine group schemes	44
3.11. Combinatorial Hopf algebras	46
3.12. Taft algebras	46
3.13. Frobenius algebras	47
3.14. Quivers and their path algebras	48
3.15. Graded (co)algebra	49
Chapter 4. Modules and comodules	51
4.1. Modules over an algebra	51
4.2. Comodules over a coalgebra	55
4.3. Tensor and cotensor products	59
4.4. Modules over a Hopf algebra	60
4.5. Hopf module algebras and coalgebras	65
Chapter 5. Finite dimensional Hopf algebras	67
5.1. The Nichols-Zoeller Theorem	67

5.2. Antipodes and finite dimensionality	67
5.3. Hopf modules	69
5.4. Applications to finite dimensional Hopf algebras	71
5.5. Semisimplicity for finite dimensional Hopf algebras	74
Chapter 6. A brief introduction to quantum groups	77
6.1. q -combinatorics	77
6.2. The Quantum Plane	79
6.3. Quasitriangular Hopf algebras	79
6.4. Braidings on module categories of quasitriangular Hopf algebras	81
6.5. Some examples of quasitriangular structures	82
Bibliography	85

CHAPTER 1

Some background material

In this chapter we give a rapid tour of some important notions that will be used.

We will assume basic familiarity with categories, functors, etc. The category theory met will be useful to most mathematicians so it is worth becoming familiar with it. The books by Saunders Mac Lane [ML98] and Emily Riehl [Rie16] are both recommended as references and for learning the subject. We will make a lot of use of commutative diagrams and basic notions from homological algebra such as exactness.

It is also assumed that readers are familiar with basic linear algebra and we focus on aspects that are more likely to be covered in advanced books on abstract algebra such as Lang [Lan02] or other encyclopaedic works.

1.1. Some category theory odds and ends

Given a category \mathbf{C} , we write $\mathbf{C}(c, d)$ for the set of morphisms $c \rightarrow d$ and $c \xrightarrow{1_c} c$ for the identity morphism of c . It is often convenient to identify an object c with its identity morphism 1_c and so we can think of the category as ‘just’ its morphisms with the objects being the subcategory of identity morphisms. We will write $\text{dom } f$ (domain/source) and $\text{codom } f$ (codomain/target) although it is very common in the literature to encounter sf and tf used for these.

For morphism $c \xrightarrow{f} d$, a morphism $d \xrightarrow{g} c$ is a *left inverse* of f if $gf = 1_c$, and $d \xrightarrow{h} c$ is a *right inverse* of f if $hf = 1_d$. If f has both a left and a right inverse they are equal and we set $f^{-1} = g = h$; then f is called an *isomorphism* and the objects c and d are *isomorphic*. Notice that for each c the set of isomorphisms $c \rightarrow c$ forms a group.

DEFINITION 1.1. Let \mathbf{C} be a category.

An object $i \in \mathbf{C}$ is *initial* if for each $c \in \mathbf{C}$, $\mathbf{C}(i, c)$ has exactly one element.

An object $t \in \mathbf{C}$ is *terminal* if for each $c \in \mathbf{C}$, $\mathbf{C}(c, t)$ has exactly one element.

If \mathbf{C} has both a terminal object and an initial object which are isomorphic then any such object is a *null object*. Such a category is sometimes called *pointed*.

It is easy to see that any two terminal objects are isomorphic, and similarly for initial objects.

Initial and terminal objects are typically denoted $\mathbf{0}$ and $\mathbf{1}$. When there is a null object, it is isomorphic to every terminal and initial object.

EXAMPLE 1.2. Here are some basic examples.

- The category of sets **Set** has \emptyset as its unique initial object, and any singleton set (i.e., a set with one element) as a terminal object.
- For the category of based (or pointed) sets **Set**_{*}, the sets with one element are null objects. Morphisms here are functions that map base points to basepoints. We can

think of a based set as a morphism $\{0\} \rightarrow X$ and morphism as a commutative diagram.

$$\begin{array}{ccc} & \{0\} & \\ \swarrow & & \searrow \\ X & \xrightarrow{f} & Y \end{array}$$

So \mathbf{Set}_* is the *slice* category of sets under $\{0\}$.

- Let \mathbf{C} be any category and c an object in it. The *slice category of objects under c* is the category c/\mathbf{C} whose objects are morphisms $c \xrightarrow{u} \text{codom } u$ and where a morphism $u \xrightarrow{f} v$ is a morphism $f \in \mathbf{C}(\text{codom } u, \text{codom } v)$ for which

$$\begin{array}{ccc} & c & \\ \swarrow u & & \searrow v \\ \text{codom } u & \xrightarrow{f} & \text{codom } v \end{array}$$

is a commutative diagram. Composition is the same as in \mathbf{C} . Notice that c/\mathbf{C} has a unique initial object $c \xrightarrow{1_c} c$, the identity morphism of c .

Similarly we can define the *slice category of objects over c* , \mathbf{C}/c .

- The categories of groups \mathbf{Gp} and abelian groups \mathbf{AbGp} have the trivial groups as null objects.
- The category of (unital) rings \mathbf{Ring} and with unital homomorphisms has \mathbb{Z} as an initial object. If we allow a trivial ring $\{0\}$ then it is a terminal object; however, for some purposes it is useful to require that $0 \neq 1$ in a unital ring. The category \mathbf{Rng} of not necessarily unital rings (i.e., rings without identity) has any trivial ring as a null object.
- In any *abelian category* there is a null object, also called a zero object. For example, this applies the categories of abelian groups, modules over a ring or sheaves of modules over a sheaf of rings on a space.


DEFINITION 1.3. Let \mathbf{C} be a category.

Suppose that we have a set of morphisms $\{c \xrightarrow{p_i} c_i : i \in I\}$ in \mathbf{C} where I is some indexing set. Then c is a *product* of the c_i (or for the p_i) if given any set of morphisms $\{d \xrightarrow{f_i} c_i : i \in I\}$ there is a unique morphism $f: d \rightarrow c$ such that for all $i \in I$, $f_i = p_i f$. If $I = \emptyset$ the *empty product* is a terminal object.

Suppose that we have a set of morphisms $\{c_i \xrightarrow{j_i} c : i \in I\}$ in \mathbf{C} where I is some indexing set. Then c is a *coproduct* of the c_i (or for the j_i) if given any set of morphisms $\{c_i \xrightarrow{g_i} d : i \in I\}$ there is a unique morphism $g: c \rightarrow d$ such that for all $i \in I$, $g_i = g j_i$. If $I = \emptyset$ the *empty coproduct* is an initial object.


There are also related notions of *pullbacks* and *pushouts* which are worth becoming familiar with. These can be subsumed into the above by working *slice categories* of objects above/below a fixed one. Another generalisation is *limits* and *colimits* but we won't discuss these.

When the indexing set is $I = \{1, 2\}$ we can express things diagrammatically: Given the diagram of solid arrows




there is a unique dotted arrow f making the whole diagram commute. A similar diagram with all arrows reversed applies to define the coproduct.

Given two products $\{c \xrightarrow{p_i} c_i : i \in I\}$ and $\{c' \xrightarrow{p'_i} c_i : i \in I\}$ it turns out that there is an isomorphism $c \xrightarrow{\cong} c'$ such that for every $i \in I$,



commutes, so it is usual to refer to *the* product and denote it by $\prod_I c_i$ or $\prod_{i \in I} c_i$; when $I = \{1, 2\}$ this is also written $c_1 \amalg c_2$. Similarly the coproduct is unique up to isomorphism and denoted $\coprod_I c_i$ and $c_1 \amalg c_2$.

It is also important that products and coproducts are *functorial* in their variables: Given products $\{c \xrightarrow{p_i} c_i : i \in I\}$ and $\{d \xrightarrow{q_i} d_i : i \in I\}$ and morphisms $f_i : c_i \rightarrow d_i$, there is a unique morphism $h : c \rightarrow d$ such that for every $j \in I$,



commutes; it is standard to denote h by $\prod_I f_i : \prod_I c_i \rightarrow \prod_I d_i$. A similar result and notation applies for coproducts.

Suppose that $\sigma : I \rightarrow J$ is a bijection. If for each $j \in J$, $d_j = c_{\sigma^{-1}(j)}$ then there is a *switch* isomorphism $T_\sigma : \prod_I c_i \xrightarrow{\cong} \prod_J d_j$. For examples, when $I = J = \{1, 2\}$, the transposition $(1\ 2)$ gives rise to an isomorphism

$$T = T_{(1\ 2)} : c_1 \amalg c_2 \xrightarrow{\cong} c_2 \amalg c_1$$

which switches the factors. Notice that the composition of switch isomorphisms

$$c_1 \amalg c_2 \xrightarrow{\cong} c_2 \amalg c_1 \xrightarrow{\cong} c_1 \amalg c_2$$

is the identity function. Similar considerations apply to coproducts.

EXAMPLE 1.4. Here are some examples of products and coproducts.

- In the category **Set**, the categorical product is the Cartesian product, the coproduct is disjoint union.
- In **Gp** the categorical product is the Cartesian product, the coproduct is *free product*.

- In an abelian category, finite products and coproducts agree and are denoted by \oplus and \bigoplus . When discussing vector spaces or modules, \times and \oplus are often used interchangeably.
- In **Ring** the product is the Cartesian product, the coproduct is much more complicated.
- In **Top** the product topology allows a product of topological spaces to be defined. Similarly there is an obvious topology on the disjoint union of a collection of spaces that makes it the coproduct.

PROPOSITION 1.5. *Suppose that in the category \mathbf{C} all products of two objects exist and there is a terminal object $\mathbf{1}$. Then all finite products exist, and for each object c ,*

$$\mathbf{1} \amalg c \cong c \cong c \amalg \mathbf{1}.$$

Similarly if all coproducts of two objects exist and there is an initial object $\mathbf{0}$, then all finite coproducts exist, and for each object c ,

$$\mathbf{0} \amalg c \cong c \cong c \amalg \mathbf{0}.$$

Monoids and comonoids. Let's recall the notion of a *monoid* in algebra. A set M together with a product/multiplication $\mu: M \times M \rightarrow M$ and a map $\iota: \mathbf{1} \rightarrow M$ (where $\mathbf{1}$ is a one element set) defines a *monoid* (M, μ, ι) if the following diagrams commute in the category **Set**.

$$\begin{array}{ccc}
 & M \times M \times M & \\
 \mu \times \text{Id} \swarrow & & \searrow \text{Id} \times \mu \\
 M \times M & & M \times M \\
 \mu \searrow & & \swarrow \mu \\
 & M &
 \end{array}
 \qquad
 \begin{array}{ccccc}
 \mathbf{1} \times M & \xleftarrow{\cong} & M & \xrightarrow{\cong} & M \times \mathbf{1} \\
 \downarrow \iota \times \text{Id} & & \downarrow \text{Id} & & \downarrow \text{Id} \times \iota \\
 M \times M & & M & & M \times M \\
 \mu \searrow & & \downarrow \mu & & \swarrow \mu \\
 & M &
 \end{array}$$

If the following diagram commutes then M is *commutative*.

$$\begin{array}{ccc}
 M \times M & \xleftarrow[\cong]{\text{T}} & M \times M \\
 \mu \searrow & & \swarrow \mu \\
 & M &
 \end{array}$$

Of course a *group* is a monoid together with a self-map $\chi: M \rightarrow M$ that satisfies some additional commutative diagrams defining left and right inverses.

Now let's generalise to a category \mathbf{C} with finite products and terminal objects. A *monoid* in \mathbf{C} is a triple $(M, \mu: M \amalg M \rightarrow M, \iota: \mathbf{1} \rightarrow M)$ where the following diagrams in \mathbf{C} commute.

$$(1.1) \qquad
 \begin{array}{ccc}
 & M \amalg M \amalg M & \\
 \mu \amalg \text{Id} \swarrow & & \searrow \text{Id} \amalg \mu \\
 M \amalg M & & M \amalg M \\
 \mu \searrow & & \swarrow \mu \\
 & M &
 \end{array}
 \qquad
 \begin{array}{ccccc}
 \mathbf{1} \amalg M & \xleftarrow{\cong} & M & \xrightarrow{\cong} & M \amalg \mathbf{1} \\
 \downarrow \iota \amalg \text{Id} & & \downarrow \text{Id} & & \downarrow \text{Id} \amalg \iota \\
 M \amalg M & & M & & M \amalg M \\
 \mu \searrow & & \downarrow \mu & & \swarrow \mu \\
 & M &
 \end{array}$$

If the following diagram commutes then M is *commutative*.

$$(1.2) \qquad
 \begin{array}{ccc}
 M \amalg M & \xleftarrow[\cong]{\text{T}} & M \amalg M \\
 \mu \searrow & & \swarrow \mu \\
 & M &
 \end{array}$$

Now one of the magical tricks of Category Theory is that any definition involving commutative diagrams can be dualised by reversing arrows, replacing products by coproducts and terminal objects by initial objects. So if \mathbf{C} has finite coproducts and initial objects then a *comonoid* in \mathbf{C} is a triple $(C, \gamma: C \rightarrow C \amalg C, \varepsilon: C \rightarrow \mathbf{0})$ making the following diagrams commute.

$$(1.3) \qquad
 \begin{array}{ccc}
 & C \amalg C \amalg C & \\
 \gamma \amalg \text{Id} \swarrow & & \searrow \text{Id} \amalg \gamma \\
 C \amalg C & & C \amalg C \\
 \gamma \searrow & & \swarrow \gamma \\
 & C &
 \end{array}
 \qquad
 \begin{array}{ccccc}
 \mathbf{0} \amalg C & \xrightarrow{\cong} & C & \xleftarrow{\cong} & C \amalg \mathbf{0} \\
 \uparrow \varepsilon \amalg \text{Id} & & \uparrow \text{Id} & & \uparrow \text{Id} \amalg \varepsilon \\
 C \amalg C & & C & & C \amalg C \\
 \gamma \searrow & & \downarrow \gamma & & \swarrow \gamma \\
 & C &
 \end{array}$$

If the following diagram commutes then C is *cocommutative*.

$$(1.4) \quad \begin{array}{ccc} C \amalg C & \xrightleftharpoons[\cong]{T} & C \amalg C \\ & \searrow \gamma \quad \nearrow \gamma & \\ & C & \end{array}$$

Note that comonoids don't exist in **Set**, but do exist in other settings such as the homotopy category of based spaces where they are called co- H -spaces. A monoid in a category **C** gives rise to a comonoid in the opposite category **C**^{op} and vice versa.

We can also introduce notions of monoids and comonoids in a *monoidal category* $(\mathbf{C}, \otimes, \mathbf{1})$; for the definition of a (symmetric) monoidal category see [ML98, Rie16]. A monoid M then has morphisms $M \otimes M \rightarrow M$ and $\mathbf{1} \rightarrow M$ fitting into commutative diagrams like (1.1) while a comonoid C has morphisms $C \rightarrow C \otimes C$ and $C \rightarrow \mathbf{1}$ with diagrams like (1.3). We will discuss the important case of vector spaces under tensor product in Section 1.2.

Here is a really important and illuminating example.

EXAMPLE 1.6. Let **AbGp** be the abelian category of abelian groups made symmetric monoidal using the tensor product \otimes . The unit object here is \mathbb{Z} since for any abelian group M ,

$$\mathbb{Z} \otimes M \cong M \cong M \otimes \mathbb{Z}.$$

The symmetry condition stems from the switch isomorphism

$$T: M \otimes N \xrightarrow{\cong} N \otimes M.$$

It is important that the composition

$$M \otimes N \xrightarrow{\cong} N \otimes M \xrightarrow{\cong} M \otimes N$$

is the identity function, which is required for (\mathbf{AbGp}, \otimes) to be *symmetric* monoidal rather than *braided* monoidal.

A monoid in $(\mathbf{AbGp}, \otimes, \mathbb{Z})$ is an abelian group R equipped with a group homomorphism $\varphi: R \otimes R \rightarrow R$ which gives a map

$$R \times R \rightarrow R \otimes R \rightarrow R; \quad (x, y) \mapsto xy = \varphi(x \otimes y)$$

and this is associative. The unit homomorphism $\eta: \mathbb{Z} \rightarrow R$ satisfies

$$\eta(1)x = x = x\eta(1),$$

so $1_R = \eta(1)$ behaves as the unity in a ring should.

The distributive laws are hidden in the fact that φ is a homomorphism of abelian groups and the tensor product is constructed to be bilinear so that

$$(x_1 + x_2) \otimes y = x_1 \otimes y + x_2 \otimes y, \quad x \otimes (y_1 + y_2) = x \otimes y_1 + x \otimes y_2.$$

Therefore a monoid in $(\mathbf{AbGp}, \otimes, \mathbb{Z})$ is a (unital) ring and a commutative monoid is just a commutative (unital) ring.

This example can be generalised by replacing \mathbb{Z} with any commutative ring, **AbGp** the abelian category of modules over it and \otimes with the tensor product of modules; a monoid is then an algebra over the ring.

Adjoint functors. Adjunctions play an important rôle in studying algebraic structures, in particular free functors and (co)monads are ubiquitous. For full definitions and properties see [ML98, Rie16].

Let \mathbf{C} and \mathbf{D} be two categories.

DEFINITION 1.7. The pair of functors $L: \mathbf{C} \rightarrow \mathbf{D}$ and $R: \mathbf{D} \rightarrow \mathbf{C}$ between categories \mathbf{C} and \mathbf{D} is an *adjoint pair* or are *adjoint functors* if there are natural (in the two variables) isomorphisms

$$\mathbf{D}(L(-), -) \cong \mathbf{C}(-, R(-)).$$

The functor L is a *left adjoint* of R and R is the *right adjoint* of L . This is indicated by writing $L \dashv R$ with the *Kan turnstyle* \dashv or \perp .

$$\begin{array}{ccc} & L & \\ \mathbf{C} & \xrightarrow{\quad} & \mathbf{D} \\ & \perp & \\ & R & \end{array}$$

It can be shown that given two left adjoints for R are naturally isomorphic and similarly for two right adjoints of L ; see [Rie16, proposition 4.4.1]. The existence of a left (or right) adjoint requires strong conditions on a functor. For example, left adjoints must preserve coproducts and right adjoints must preserve products. Results on this are discussed in [Rie16, section 4.6].

Notice that for any $X \in \mathbf{C}$, under the isomorphism $\mathbf{D}(L(X), L(X)) \cong \mathbf{C}(X, R(L(X)))$, the identity morphism $I_{L(X)}$ corresponds to a morphism $i_X: X \rightarrow R(L(X))$ usually called a *universal morphism*. Similarly for any $Y \in \mathbf{D}$ there is a morphism $i^Y: L(R(Y)) \rightarrow Y$ corresponding to $I_{R(Y)}$ under $\mathbf{C}(R(Y), R(Y)) \cong \mathbf{D}(L(R(Y)), Y)$. We can think of i_X as part of a natural transformation $i_{(\cdot)}: I_{\mathbf{C}} \Rightarrow RL$ and i^Y as part of a natural transformation $i^{(\cdot)}: LR \Rightarrow I_{\mathbf{D}}$.

EXAMPLE 1.8. The forgetful functor $R: \mathbf{Gp} \rightarrow \mathbf{Set}$ which sends a group (G, μ, e) to its underlying set G has a left adjoint $L: \mathbf{Set} \rightarrow \mathbf{Gp}$ which sends a set X to the *free group* generated by it, namely $\langle X \rangle$. The function $i_X: X \rightarrow \langle X \rangle$ just sends an element $x \in X$ to the word x , while $i^G: \langle G \rangle \rightarrow G$ sends a word in elements of G to the element of G obtained by evaluating the product of the letters.

Given an adjoint pair L, R , we can define endofunctors $RL: \mathbf{C} \rightarrow \mathbf{C}$ and $LR: \mathbf{D} \rightarrow \mathbf{D}$. Then there are natural transformations

$$(RL)(RL) = R(LR)L \xrightarrow{Ri^{(\cdot)}L} RI_{\mathbf{D}}L = RL$$

$$LR = LI_{\mathbf{C}}R \xrightarrow{Li_{(\cdot)}R} L(RL)R = (LR)(LR)$$

so that $(RL, i_{(\cdot)}, Ri^{(\cdot)}L)$ is a *monad* in \mathbf{C} and $(LR, i^{(\cdot)}, Li_{(\cdot)}R)$ is a *comonad* in \mathbf{D} . So each is a ‘monoid or comonoid in one of the categories of endofunctors of \mathbf{C} and \mathbf{D} ’; see [Rie16, page 154].

1.2. Vector spaces over a field as a symmetric monoidal category

In this section we will discuss an important example that underlies the study of Hopf algebras and many other things. This material requires knowledge of tensor products of vector spaces and we will review some of the ideas. Throughout, \mathbb{k} will be a field.

The abelian category of (left) \mathbb{k} -vector spaces $\mathbf{Vect}_{\mathbb{k}}$ is very simple in terms of its *additive* structure. For example, every short exact sequence splits

$$0 \rightarrow U \rightarrow V \rightarrow W \rightarrow 0$$

so that $V \cong U \oplus W$. As a result there is no homological algebra like there is for modules over a general ring. Of course this is a consequence of the existence of bases, which also implies that every vector space is a direct sum of 1-dimensional ones.

However, additional structure is available: $\mathbf{Vect}_{\mathbb{k}}$ is also a *closed symmetric monoidal category* under tensor product $\otimes = \otimes_{\mathbb{k}}$ and with the internal function object given by

$$\mathrm{hom}(-, -) = \mathrm{Hom}_{\mathbb{k}}(-, -) = \mathbf{Vect}_{\mathbb{k}}(-, -).$$

The vector space \mathbb{k} is a unit object for these since there are functorial isomorphisms

$$\mathbb{k} \otimes V \cong V \cong V \otimes \mathbb{k}, \quad \mathrm{hom}(\mathbb{k}, V) \cong V.$$

It is symmetric because of the functorial switch isomorphism

$$T: U \otimes V \xrightarrow{\cong} V \otimes U; \quad T(x \otimes y) = y \otimes x.$$

The tensor product is functorial in the two variables: given linear mappings $f: U \rightarrow U'$ and $g: V \rightarrow V'$ there is a linear mapping $f \otimes g: U \otimes V \rightarrow U' \otimes V'$ fitting into a commutative diagram of linear mappings.

$$\begin{array}{ccc} U \otimes V & \xrightarrow{f \otimes \mathrm{Id}_V} & U' \otimes V \\ \mathrm{Id}_{U'} \otimes g \downarrow & \searrow f \otimes g & \downarrow \mathrm{Id}_U \otimes g \\ U \otimes V' & \xrightarrow{f \otimes \mathrm{Id}_{V'}} & U' \otimes V' \end{array}$$

The tensor product is associative in the sense that for three vector spaces U, V, W , there is a canonical isomorphism

$$(U \otimes V) \otimes W \xrightarrow{\cong} U \otimes (V \otimes W)$$

ultimately induced from the canonical bijection of sets

$$(U \times V) \times W \longleftrightarrow U \times (V \times W)$$

using the universal property of \otimes repeatedly (this is left as an exercise for those who have not seen it before). Then for linear mappings $f: U \rightarrow U'$, $g: V \rightarrow V'$ and $h: W \rightarrow W'$ there is a commutative diagram.

$$\begin{array}{ccc} (U \otimes V) \otimes W & \xleftarrow{\cong} & U \otimes (V \otimes W) \\ (f \otimes g) \otimes h \downarrow & & \downarrow f \otimes (g \otimes h) \\ (U' \otimes V') \otimes W' & \xleftarrow{\cong} & U' \otimes (V' \otimes W') \end{array}$$

Because of this we usually just write $U \otimes V \otimes W$ for $(U \otimes V) \otimes W$ and $f \otimes g \otimes h: U \otimes V \otimes W \rightarrow U' \otimes V' \otimes W'$ for $(f \otimes g) \otimes h: (U \otimes V) \otimes W \rightarrow U' \otimes V' \otimes W'$, and identify it with $f \otimes (g \otimes h): U \otimes (V \otimes W) \rightarrow U' \otimes (V' \otimes W')$ using the isomorphisms and diagram above.

REMARK 1.9. Of course all of these properties of tensor products are consequences of the basic universal property they satisfy. We recall that given three \mathbb{k} -vector spaces U, V, W , a function $f: U \times V \rightarrow W$ is *bilinear* if for all $s_1, s_2, t_1, t_2 \in \mathbb{k}$, $u_1, u_2 \in U$ and $v_1, v_2 \in V$,

$$f(s_1 u_1 + s_2 u_2, t_1 v_1 + t_2 v_2) = \sum_{i,j=1,2} s_i t_j f(u_i, v_j).$$

Now a bilinear function $\tau: U \times V \rightarrow T$ into a vector space T is a tensor product for U, V if for every bilinear map $f: U \times V \rightarrow W$ there is a unique linear mapping $f': T \rightarrow W$ such that $f' \circ \tau = f$.

$$\begin{array}{ccc} U \times V & \xrightarrow{\tau} & T \\ & \searrow f & \swarrow \exists! f' \\ & W & \end{array}$$

This universal property is used to build maps out of the tensor product into a vector space by specifying bilinear maps. As usual the vector space T is unique up to isomorphism and there is a standard construction that can be found in many books. We denote it by $U \otimes V = U \otimes_{\mathbb{k}} V$ and then the map τ is given by $\tau(u, v) = u \otimes v$ where such *basic tensors* span $U \otimes V$.

Some adjunctions for vector spaces. Our next result gives an example of an adjunction that has many variations for modules over rings. It can be proved using the universal property of \otimes . It also justifies the claim that $(\mathbf{Vect}_{\mathbb{k}}, \otimes, \text{hom})$ is a *closed* monoidal category as mentioned earlier.

Note that $\mathbf{Vect}_{\mathbb{k}}(U, V)$ is not just a set but a vector space. The adjunction isomorphisms in the next result are actually vector space isomorphisms (i.e., invertible linear mappings).

THEOREM 1.10. *For a \mathbb{k} -vector space V , the functors*

$$(-) \otimes V: \mathbf{Vect}_{\mathbb{k}} \rightarrow \mathbf{Vect}_{\mathbb{k}}, \quad \text{hom}(V, -): \mathbf{Vect}_{\mathbb{k}} \rightarrow \mathbf{Vect}_{\mathbb{k}}$$

are adjoint, $(-) \otimes V \dashv \text{hom}(V, -)$. Hence for any three vector spaces U, V, W there is an adjunction isomorphism

$$\mathbf{Vect}_{\mathbb{k}}(U \otimes V, W) \xrightarrow{\cong} \mathbf{Vect}_{\mathbb{k}}(U, \text{hom}(V, W))$$

which is functorial in the variables.

This means that for linear mappings $f: U \rightarrow U'$, $g: V \rightarrow V'$ and $h: W \rightarrow W'$ there are commutative diagrams involving these isomorphisms.

$$\begin{array}{ccc} \mathbf{Vect}_{\mathbb{k}}(U' \otimes V, W) & \xleftarrow{\cong} & \mathbf{Vect}_{\mathbb{k}}(U', \text{hom}(V, W)) \\ (f \otimes \text{Id})^* \downarrow & & \downarrow f^* \\ \mathbf{Vect}_{\mathbb{k}}(U \otimes V, W) & \xleftarrow{\cong} & \mathbf{Vect}_{\mathbb{k}}(U, \text{hom}(V, W)) \end{array}$$

$$\begin{array}{ccc} \mathbf{Vect}_{\mathbb{k}}(U \otimes V', W) & \xleftarrow{\cong} & \mathbf{Vect}_{\mathbb{k}}(U, \text{hom}(V', W)) \\ (\text{Id} \otimes g)^* \downarrow & & \downarrow (g^*)^* \\ \mathbf{Vect}_{\mathbb{k}}(U \otimes V, W) & \xleftarrow{\cong} & \mathbf{Vect}_{\mathbb{k}}(U, \text{hom}(V, W)) \end{array}$$

$$\begin{array}{ccc} \mathbf{Vect}_{\mathbb{k}}(U \otimes V, W) & \xleftarrow{\cong} & \mathbf{Vect}_{\mathbb{k}}(U, \text{hom}(V, W)) \\ h_* \downarrow & & \downarrow (h_*)^* \\ \mathbf{Vect}_{\mathbb{k}}(U \otimes V, W') & \xleftarrow{\cong} & \mathbf{Vect}_{\mathbb{k}}(U, \text{hom}(V, W')) \end{array}$$

The *dual (space)* of V is $V^* = \text{hom}(V, \mathbb{k})$; if V is finite dimensional, a choice of basis leads to an isomorphism $V \xrightarrow{\cong} V^*$ dependent on the basis used.

The (*strongly*) *dualisable* objects V are characterised by the condition that for all W ,

$$\text{hom}(V, W) \cong W \otimes V^*,$$

and

$$V^{**} = \text{hom}(V^*, \mathbb{k}) \cong V,$$

where the latter isomorphism can be chosen to be independent of choice of basis and functorial in V ; these turn out to be precisely the finite dimensional vector spaces. Notice also that $\mathbb{k}^* \cong \mathbb{k}$ and $\text{End}_{\mathbb{k}}(V) \cong V \otimes V^*$. There are also functorial isomorphisms

$$\text{hom}(U \otimes V, W) = \mathbf{Vect}_{\mathbb{k}}(U \otimes V, W) \cong \mathbf{Vect}_{\mathbb{k}}(U, W \otimes V^*) = \text{hom}(U, W \otimes V^*).$$

When U and V are finite dimensional, we will make the canonical identification

$$(1.5) \quad (U \otimes V)^* \cong V^* \otimes U^*$$

not with $U^* \otimes V^*$, although these are isomorphic via the switch isomorphism; the literature has varying conventions on this and some minor differences occur as a result. Warning: when U or V is infinite dimensional there is a canonical injective linear mapping

$$(1.6) \quad V^* \otimes U^* \rightarrow (U \otimes V)^*$$

which is not an isomorphism.

For later use we mention an important construction. There is a *forgetful functor* $\mathbf{Vect}_{\mathbb{k}} \rightarrow \mathbf{Set}$ which ‘forgets’ the algebraic structure and just remembers the underlying set; it sends each \mathbb{k} -linear mapping to itself just viewed as a function between sets. Thus $\mathbf{Vect}_{\mathbb{k}}$ is an example of a *concrete category*; many of the familiar examples of mathematical structures form concrete categories.

PROPOSITION 1.11. *There is a functor $\mathbb{F}: \mathbf{Set} \rightarrow \mathbf{Vect}_{\mathbb{k}}$ which is left adjoint to the forgetful functor, i.e., for every set X and vector space V there is a bijection*

$$\mathbf{Vect}_{\mathbb{k}}(\mathbb{F}(X), V) \cong \mathbf{Set}(X, V),$$

and this gives a natural isomorphism of bifunctors

$$\mathbf{Vect}_{\mathbb{k}}(\mathbb{F}(-), -) \cong \mathbf{Set}(-, -).$$

Furthermore, there is a natural isomorphism $\mathbb{F} \circ (- \times -) \cong \mathbb{F}(-) \otimes \mathbb{F}(-)$ between the bifunctors

$$\begin{aligned} \mathbb{F} \circ (- \times -): \mathbf{Set} \times \mathbf{Set} &\rightarrow \mathbf{Vect}_{\mathbb{k}}; & (X, Y) &\mapsto \mathbb{F}(X \times Y), \\ \mathbb{F}(-) \otimes \mathbb{F}(-): \mathbf{Set} \times \mathbf{Set} &\rightarrow \mathbf{Vect}_{\mathbb{k}}; & (X, Y) &\mapsto \mathbb{F}(X) \otimes \mathbb{F}(Y). \end{aligned}$$

We usually think of the vector space $\mathbb{F}X = \mathbb{F}(X)$ as having X as a basis and it is called the (*free*) *vector space on X* . One construction is

$$\mathbb{F}(X) = \{(\alpha: X \rightarrow \mathbb{k}) : \alpha \text{ is finitely supported}\},$$

where a function is *finitely supported* if it is zero except on finitely many elements. Of course if we take $X = \{1, 2, \dots, n\}$, $\mathbb{F}X \cong \mathbb{k}^n$; in particular, if $n = 1$, $\mathbb{F}X \cong \mathbb{k}$.

Of course every vector space V has a basis B , and there is a unique linear mapping $\mathbb{F}(B) \rightarrow V$ corresponding to the inclusion function $B \hookrightarrow V$; this is easily seen to be an isomorphism, so $\mathbb{F}(B) \cong V$. Of course this just says that every vector space is a free vector space but in many ways corresponding to different choices of bases.

We will see later that free functors and adjunctions provide powerful tools for creating algebraic objects with useful properties.

Some homological algebra. As mentioned before, the abelian category of vector spaces $\mathbf{Vect}_{\mathbb{k}}$ is *semi-simple* since every short exact sequence in $\mathbf{Vect}_{\mathbb{k}}$ splits, so it has no homological algebra in the sense of derived functors. More specifically, for any vector space W , the additive functors

$$W \otimes (-), (-) \otimes W, \text{hom}(W, -), \text{hom}(-, W): \mathbf{Vect}_{\mathbb{k}} \rightarrow \mathbf{Vect}_{\mathbb{k}}$$

are all *exact*, i.e., they send every short exact sequence to a short exact sequence. So if

$$0 \rightarrow V' \rightarrow V \rightarrow V'' \rightarrow 0$$

is short exact then so is each of the induced sequences

$$\begin{aligned} 0 \rightarrow W \otimes V' \rightarrow W \otimes V \rightarrow W \otimes V'' \rightarrow 0, \\ 0 \rightarrow V' \otimes W \rightarrow V \otimes W \rightarrow V'' \otimes W \rightarrow 0, \\ 0 \rightarrow \text{hom}(W, V') \rightarrow \text{hom}(W, V) \rightarrow \text{hom}(W, V'') \rightarrow 0, \\ 0 \rightarrow \text{hom}(V'', W) \rightarrow \text{hom}(V, W) \rightarrow \text{hom}(V', W) \rightarrow 0. \end{aligned}$$

1.3. Graded vector spaces

In many contexts, vectors spaces, modules, etc, are equipped with *gradings*. In principal we can do this with an arbitrary abelian group but in these notes we will only discuss the very common case of the integers.

A \mathbb{Z} -graded \mathbb{k} -vector space means a collection of vector spaces $V_* = \{V_n : n \in \mathbb{Z}\}$ or $V^* = \{V^n : n \in \mathbb{Z}\}$. Here the upper (homology) and lower (cohomology) indexing is often used to distinguish ‘homological’ and ‘cohomological’ degrees (for example in Algebraic Topology).

Given graded vector spaces U_* and V^* , together with an integer d , we can form new graded vector spaces $U[d]_*$ or $V[d]^*$ where

$$U[d]_n = U_{n-d}, \quad V[d]_n = V^{n-d}.$$

From now on we will focus on lower gradings but the upper gradings have similar properties.

Given two such objects U_* and V_* , a *homomorphism of graded vector spaces* $f_*: U_* \rightarrow V_*$ is a collection of homomorphisms $f_* = \{f_n: U_n \rightarrow V_n : n \in \mathbb{Z}\}$. We can also talk about a *homomorphism of degree d* which is a homomorphism $U[d]_* \rightarrow V_*$. Notice that this is the same as a homomorphism $U[d+k]_* \rightarrow V[k]_*$ for any k . We can form a graded version of $\text{Hom} = \text{Hom}_{\mathbb{k}}$ by taking $\text{Hom}^d(U_*, V_*)$ to be the set of all homomorphisms of degree d from U_* to V_* . However, this can be large since

$$\text{Hom}^d(U_*, V_*) = \prod_{n \in \mathbb{Z}} \text{Hom}(U_{d+n}, V_n).$$

We can also tensor graded vector spaces. The tensor product of U_* and V_* is the graded vector space $U_* \otimes_{\mathbb{k}} V_* = (U \otimes_{\mathbb{k}} V)_*$ with

$$(U \otimes_{\mathbb{k}} V)_n = \bigoplus_{k \in \mathbb{Z}} U_k \otimes_{\mathbb{k}} V_{n-k}.$$

It is often useful to work with graded objects where the only non-zero terms are in non-negative degrees. A graded vector space V_* is *d -connected* if $V_k = 0$ whenever $k \leq d$ it is called *bounded below* if it is d -connected for some d . We can also talk about *coconnected* and *bounded above* objects.

A graded vector space V_* is of *finite type* if for every $n \in \mathbb{Z}$, $\dim_{\mathbb{k}} V_n < \infty$. It is (*totally*) *finite* if it of finite type and only finitely many of the $\dim_{\mathbb{k}} V_n$ are non-zero.

If V_* is of finite type we can define its *dual* to be V^* where

$$V^n = \text{Hom}_{\mathbb{k}}(V_n, \mathbb{k}),$$

so V^* is also of finite type.

It is possible to make graded vector spaces into a symmetric monoidal category and this will be useful for Section 3.15.

The Koszul sign convention. In graded contexts there is often a sign built in depending on degrees. The origin of this lies in the switch isomorphism

$$U \otimes V \xrightarrow[\cong]{T} V \otimes U.$$

When we are working with graded vector spaces U_* and V_* this is replaced with the isomorphism

$$U_m \otimes V_n \xrightarrow[\cong]{(-1)^{mn}T} V_n \otimes U_m.$$

Of course the sign factor here is only -1 if both m and n are odd. So if we are working with graded vector spaces which are only non-zero in even degrees this sign does not differ from that in the ungraded world. However, this new sign convention affects the notion of graded commutativity as we will see.

Notice that when $\text{char } \mathbb{k} = 2$ the Koszul signs are all 1 anyway.

CHAPTER 2

Algebras, coalgebras, bialgebras and Hopf algebras

In this chapter we will introduce two types of algebraic structures, namely *algebras* and *coalgebras*, then combine them into a new structure called a *bialgebra*. Finally we will focus on bialgebras which possess an *antipode*; these are our central objects of study, namely *Hopf algebras*.

Throughout, \mathbb{k} will be a field. In fact, much of the theory works for an arbitrary commutative ring, although additional assumptions such as flatness are sometimes required when discussing coalgebras. We will often set $\otimes = \otimes_{\mathbb{k}}$ and $\text{hom} = \text{Hom}_{\mathbb{k}}$.

2.1. Algebras

A \mathbb{k} -*algebra* (A, φ, η) is a *monoid* in the symmetric monoidal category $(\mathbf{Vect}_{\mathbb{k}}, \otimes, \mathbb{k})$, i.e., a \mathbb{k} -vector space A equipped with a \mathbb{k} -linear *product* $\varphi: A \otimes A \rightarrow A$ and *unit* $\eta: \mathbb{k} \rightarrow A$, which make the following diagrams in $\mathbf{Vect}_{\mathbb{k}}$ commute where the isomorphisms are the canonical ones.

$$(2.1) \quad \begin{array}{ccc} (A \otimes A) \otimes A & \xleftarrow{\cong} & A \otimes (A \otimes A) \\ \varphi \otimes \text{Id} \downarrow & & \downarrow \text{Id} \otimes \varphi \\ A \otimes A & & A \otimes A \\ \varphi \searrow & & \swarrow \varphi \\ & A & \end{array} \quad \begin{array}{ccc} \mathbb{k} \otimes A & \xleftarrow{\cong} & A \xrightarrow{\cong} A \otimes \mathbb{k} \\ \eta \otimes \text{Id} \downarrow & & \downarrow \text{Id} \otimes \eta \\ A \otimes A & \xrightarrow{\text{Id}} & A \otimes A \\ \varphi \searrow & & \swarrow \varphi \\ & A & \end{array}$$

If in addition the following diagram commutes then A is *commutative*.

$$(2.2) \quad \begin{array}{ccc} A \otimes A & \xleftarrow[\cong]{\tau} & A \otimes A \\ \varphi \searrow & & \swarrow \varphi \\ & A & \end{array}$$

An algebra is really a special kind of ring and when working with elements we set $xy = \varphi(x \otimes y)$ and $1 = 1_A = \eta(1)$ if this is unlikely to lead to confusion. Of course commutativity means that for all $x, y \in A$, $xy = yx$.

Unpacking the definition we find that an algebra is a ring with the additional structure of a specified (injective) ring homomorphism $\eta: \mathbb{k} \rightarrow A$ whose image is contained in the centre of A and makes A a \mathbb{k} -vector space. Commutativity means that A is a commutative ring.

A *homomorphism* $\theta: (A, \varphi, \eta) \rightarrow (A', \varphi', \eta')$ between two \mathbb{k} -algebras is a \mathbb{k} -linear mapping $\theta: A \rightarrow A'$ making the following diagrams commute.

$$\begin{array}{ccc} A \otimes A & \xrightarrow{\theta \otimes \theta} & A' \otimes A' \\ \varphi \downarrow & & \downarrow \varphi' \\ A & \xrightarrow{\theta} & A' \end{array} \quad \begin{array}{ccc} & \mathbb{k} & \\ \eta \swarrow & & \searrow \eta' \\ A & \xrightarrow{\theta} & A' \end{array}$$

So a homomorphism is a ring homomorphism which is also a \mathbb{k} -linear mapping. Of course the kernel of a homomorphism θ is an ideal, $\ker \theta \triangleleft A$, and the image of θ is a subalgebra of A' isomorphic to the quotient algebra $A/\ker \theta$. The conditions for a subspace $I \subseteq A$ to be a two-sided ideal amount to saying that there is commutative diagram of the following form.

$$\begin{array}{ccccc} A \otimes I & \xrightarrow{\text{id} \otimes \text{inc}} & A \otimes A & \xleftarrow{\text{inc} \otimes \text{id}} & I \otimes A \\ \vdots \downarrow & & \downarrow \varphi & & \downarrow \vdots \\ I & \xrightarrow{\text{inc}} & A & \xleftarrow{\text{inc}} & I \end{array}$$

To show that the kernel of the algebra homomorphism θ is an ideal we consider the diagram of solid arrows

$$(2.3) \quad \begin{array}{ccccccc} 0 & \longrightarrow & A \otimes \ker \theta & \xrightarrow{\text{Id} \otimes \text{inc}} & A \otimes A & \xrightarrow{\text{Id} \otimes \theta} & A \otimes A' \\ & & \vdots \downarrow & & \downarrow \varphi & & \downarrow \varphi' \circ (\text{Id} \otimes \theta) \\ 0 & \longrightarrow & \ker \theta & \xrightarrow{\text{inc}} & A & \xrightarrow{\theta} & A' \\ & & \uparrow \vdots & & \uparrow \varphi & & \uparrow \varphi' \circ (\theta \otimes \text{Id}) \\ 0 & \longrightarrow & \ker \theta \otimes A & \xrightarrow{\text{inc} \otimes \text{Id}} & A \otimes A & \xrightarrow{\text{Id} \otimes \theta} & A' \otimes A \end{array}$$

in which the rows are exact. The existence of the dotted arrows making the resulting diagram commute is now an easy exercise.

The *trivial \mathbb{k} -algebra* is \mathbb{k} with the product given by the canonical isomorphism $\mathbb{k} \otimes \mathbb{k} \xrightarrow{\cong} \mathbb{k}$ which is given on basic tensors by

$$r \otimes s \mapsto rs.$$

Since \mathbb{k} is a field, for any algebra the unit $\eta: \mathbb{k} \rightarrow A$ is an injective homomorphism of \mathbb{k} -algebras and it is usual to identify its image with \mathbb{k} , thus making \mathbb{k} a subring of A . The exact sequence of vector spaces

$$0 \rightarrow \mathbb{k} \rightarrow A \rightarrow \text{coker } \eta \rightarrow 0$$

splits, giving a linear isomorphism $A \cong \mathbb{k} \oplus \text{coker } \eta$. However this isomorphism depends on choosing a basis of A which extends a basis of \mathbb{k} . If additional structure is present then it can sometimes be made canonical.

A \mathbb{k} -algebra A is *augmented* if there is a given homomorphism of \mathbb{k} -algebras $\varepsilon: A \rightarrow \mathbb{k}$, so $\ker \varepsilon \triangleleft A$. Notice that the commutative diagram of vector spaces and linear mappings

$$\begin{array}{ccccccc} & & 0 & & & & \\ & & \downarrow & & & & \\ & & \mathbb{k} & & & & \\ & & \downarrow \eta & & & & \\ 0 & \longrightarrow & \ker \varepsilon & \longrightarrow & A & \xrightarrow{\varepsilon} & \mathbb{k} \longrightarrow 0 \\ & & \searrow \cong & & \downarrow & & \\ & & & & \text{coker } \eta & & \\ & & & & \downarrow & & \\ & & & & 0 & & \end{array}$$

has exact row and column, and the composition

$$\ker \varepsilon \xrightarrow{\quad} A \xrightarrow{\quad} \operatorname{coker} \eta$$

is an isomorphism. Therefore there is a canonical decomposition of vector spaces

$$A \cong \mathbb{k} \oplus \ker \varepsilon \cong \mathbb{k} \oplus \operatorname{coker} \eta.$$

Given an algebra A , its *opposite algebra* A^{op} has the same underlying vector space but product

$$\varphi^{\operatorname{op}} = \varphi \circ T.$$

So if we denote $a \in A$ viewed as an element of A^{op} by a^{op} ,

$$a^{\operatorname{op}} b^{\operatorname{op}} = \varphi^{\operatorname{op}}(a^{\operatorname{op}} \otimes b^{\operatorname{op}}) = (ba)^{\operatorname{op}}.$$

The identity function $A \rightarrow A^{\operatorname{op}}$ is an algebra homomorphism if and only if A is commutative. It is not always possible to find an isomorphism $A \rightarrow A^{\operatorname{op}}$ but it does sometimes occur.

EXAMPLE 2.1. The ring $M_n(\mathbb{k})$ of $n \times n$ matrices with entries in \mathbb{k} is an algebra which has opposite algebra $M_n(\mathbb{k})^{\operatorname{op}}$. Transposition defines an isomorphism

$$(-)^T: M_n(\mathbb{k}) \rightarrow M_n(\mathbb{k})^{\operatorname{op}}; \quad X \mapsto (X^T)^{\operatorname{op}}.$$

More generally, if A is any algebra with a specified isomorphism $\varphi: A \rightarrow A^{\operatorname{op}}$ then $M_n(A)$, $M_n(A)^{\operatorname{op}}$, $M_n(A^{\operatorname{op}})$ and $M_n(A^{\operatorname{op}})^{\operatorname{op}}$ are all algebras and there is an isomorphisms

$$M_n(A) \rightarrow M_n(A^{\operatorname{op}})^{\operatorname{op}}; \quad X \mapsto (X^\dagger)^{\operatorname{op}}$$

where given the matrix X with (i, j) entry x_{ij} , X^\dagger has (i, j) entry $\varphi(x_{ji})$.

A possibly familiar example of this occurs when $\mathbb{k} = \mathbb{R}$ and $A = \mathbb{H}$ (the quaternions).

EXAMPLE 2.2. The ring of polynomials $\mathbb{k}[X_1, \dots, X_n]$ is a commutative algebra. If we take an ideal $I \triangleleft \mathbb{k}[X_1, \dots, X_n]$ then the quotient ring $\mathbb{k}[X_1, \dots, X_n]/I$ is also a commutative algebra and the quotient homomorphism $\mathbb{k}[X_1, \dots, X_n] \rightarrow \mathbb{k}[X_1, \dots, X_n]/I$ is an algebra homomorphism.

The ring of non-commutative polynomials $\mathbb{k}\langle X_1, \dots, X_n \rangle$ is an algebra which has all the (ordered) monomials $X_{r_1} \cdots X_{r_\ell}$ as a basis. The product is of course obtained from juxtaposing monomials. Again the quotient by an ideal is also an algebra.

If we choose a non-zero $h \in \mathbb{k}$ then the quotient of $\mathbb{k}\langle X, Y \rangle$ by the principal ideal generated by $XY - YX - h$, $\mathbb{k}\langle X, Y \rangle / (XY - YX - h)$, is called the *Weyl algebra*. We can think of Y as the differential operator $h \frac{d}{dX}$ and then $\mathbb{k}\langle X, Y \rangle / (XY - YX - h)$ is the ring of formal differential operators.

Given two algebras (A_1, φ_1, η_1) and (A_2, φ_2, η_2) , their tensor product $A_1 \otimes A_2$ becomes an algebra with product and unit given by the compositions

$$(A_1 \otimes A_2) \otimes (A_1 \otimes A_2) \xrightarrow[\cong]{\operatorname{Id}_{A_1} \otimes T \otimes \operatorname{Id}_{A_2}} (A_1 \otimes A_1) \otimes (A_2 \otimes A_2) \xrightarrow{\varphi_1 \otimes \varphi_2} A_1 \otimes A_2$$

and

$$\mathbb{k} \xrightarrow[\cong]{} \mathbb{k} \otimes \mathbb{k} \xrightarrow{\eta_1 \otimes \eta_2} A_1 \otimes A_2.$$

So given basic tensors $a_1 \otimes a_2, b_1 \otimes b_2 \in A_1 \otimes A_2$, their product is

$$(a_1 \otimes a_2)(b_1 \otimes b_2) = a_1 b_1 \otimes a_2 b_2.$$

EXAMPLE 2.3. Given an algebra A and its opposite algebra A^{op} , the algebra $A^e = A \otimes A^{\text{op}}$ is called the *enveloping algebra* of A . It is encountered when studying bimodules over A and Hochschild (co)homology.

We can summarise the main properties of \mathbb{k} -algebras and their homomorphisms in the following.

THEOREM 2.4. *Algebras and commutative algebras form the symmetric monoidal categories $\mathbf{Alg}_{\mathbb{k}}$ and $\mathbf{Alg}_{\mathbb{k}}^{\text{co}}$ under \otimes . In $\mathbf{Alg}_{\mathbb{k}}^{\text{co}}$, \otimes is the categorical coproduct and \mathbb{k} is an initial object. In $\mathbf{Alg}_{\mathbb{k}}$ and $\mathbf{Alg}_{\mathbb{k}}^{\text{co}}$ the Cartesian product $\times = \oplus$ is the categorical product.*

2.2. Coalgebras

The dual notion to an algebra is that of a \mathbb{k} -coalgebra, which is a triple (C, ψ, ε) , with C a \mathbb{k} -vector space, a *coproduct* $\psi: C \rightarrow C \otimes C$, and a *counit* $\varepsilon: C \rightarrow \mathbb{k}$ fitting into the commutative diagrams shown.

$$(2.4) \quad \begin{array}{ccc} (C \otimes C) \otimes C & \xleftarrow{\cong} & C \otimes (C \otimes C) \\ \psi \otimes \text{Id} \uparrow & & \uparrow \text{Id} \otimes \psi \\ C \otimes C & & C \otimes C \\ \psi \swarrow & & \searrow \psi \\ & C & \end{array} \quad \begin{array}{ccccc} \mathbb{k} \otimes C & \xrightarrow{\cong} & C & \xleftarrow{\cong} & C \otimes \mathbb{k} \\ \varepsilon \otimes \text{Id} \uparrow & & \uparrow \text{Id} & & \uparrow \text{Id} \otimes \varepsilon \\ C \otimes C & & C & & C \otimes C \\ \psi \swarrow & & \uparrow & & \searrow \psi \\ & C & \end{array}$$

This says that (C, ψ, ε) is a *comonoid* in $\mathbf{Vect}_{\mathbb{k}}$.

If the following diagram commutes then C is *cocommutative*.

$$(2.5) \quad \begin{array}{ccc} C \otimes C & \xleftarrow[\cong]{\tau} & C \otimes C \\ \psi \swarrow & & \searrow \psi \\ & C & \end{array}$$

A *homomorphism* $\theta: (C, \psi, \varepsilon) \rightarrow (C', \psi', \varepsilon')$ between two \mathbb{k} -coalgebras is a \mathbb{k} -linear mapping $\theta: C \rightarrow C'$ making the following diagrams commute.

$$\begin{array}{ccc} C \otimes C & \xrightarrow{\theta \otimes \theta} & C' \otimes C' \\ \psi \uparrow & & \uparrow \psi' \\ C & \xrightarrow{\theta} & C' \end{array} \quad \begin{array}{ccc} & \mathbb{k} & \\ \varepsilon \nearrow & & \nwarrow \varepsilon' \\ C & \xrightarrow{\theta} & C' \end{array}$$

The kernel of θ is a coideal, where a subspace $J \subseteq C$ is a *coideal* if the coproduct ψ restricts to give a map $J \rightarrow C \otimes J + J \otimes C \subseteq C \otimes C$ and $\varepsilon J = 0$; then there is a commutative diagram

$$\begin{array}{ccc} J & \xrightarrow{\quad} & C \otimes J + J \otimes C \\ \downarrow & & \downarrow \\ C & \xrightarrow{\psi} & C \otimes C \\ \downarrow & & \downarrow \\ C/J & \xrightarrow{\bar{\psi}} & C/J \otimes C/J \end{array}$$

To prove that the kernel of a coalgebra homomorphism is a coideal involves an analogue of the diagram (2.3). For this we note first that

$$\ker(\theta \otimes \theta) = C \otimes \ker \theta + \ker \theta \otimes C$$

and so there is a commutative diagram of solid arrows

$$(2.6) \quad \begin{array}{ccccccc} 0 & \longrightarrow & \ker \theta & \xrightarrow{\text{inc}} & C & \xrightarrow{\theta} & C' \\ & & \downarrow \text{dotted} & & \downarrow \psi & & \downarrow \psi' \\ 0 & \longrightarrow & \ker(\theta \otimes \theta) & \xrightarrow{\text{inc}} & C \otimes C & \xrightarrow{\theta \otimes \theta} & C' \otimes C' \\ & & \parallel & & & & \\ & & C \otimes \ker \theta + \ker \theta \otimes C & & & & \end{array}$$

and there is a unique linear map giving the dotted arrow which makes the resulting diagram commute.

The image of θ is a *subcoalgebra* of C' isomorphic to the quotient C/J equipped with the induced coproduct $\bar{\psi}: C/J \rightarrow C/J \otimes C/J$.

The *trivial* \mathbb{k} -coalgebra is \mathbb{k} with the coproduct given by the isomorphism

$$\mathbb{k} \xrightarrow{\cong} \mathbb{k} \otimes_{\mathbb{k}} \mathbb{k} = \mathbb{k} \otimes \mathbb{k}$$

which on basic tensors is just

$$t \mapsto t \otimes 1 = 1 \otimes t.$$

For any coalgebra the counit $\varepsilon: C \rightarrow \mathbb{k}$ is a homomorphism of \mathbb{k} -coalgebras.

A \mathbb{k} -coalgebra C is *(co)augmented* if there is a homomorphism of \mathbb{k} -coalgebras $\eta: \mathbb{k} \rightarrow C$. The commutative diagram of vector spaces and linear mappings has exact row and column

$$\begin{array}{ccccccc} & & 0 & & & & \\ & & \downarrow & & & & \\ & & \ker \varepsilon & & & & \\ & & \downarrow & \searrow \cong & & & \\ 0 & \longrightarrow & \mathbb{k} & \xrightarrow{\eta} & C & \longrightarrow & \text{coker } \eta \longrightarrow 0 \\ & & \parallel & & \downarrow \varepsilon & & \\ & & & & \mathbb{k} & & \\ & & & & \downarrow & & \\ & & & & 0 & & \end{array}$$

and there is a canonical decomposition of vector spaces

$$C \cong \mathbb{k} \oplus \ker \varepsilon \cong \mathbb{k} \oplus \text{coker } \eta.$$

A coalgebra C has an *opposite coalgebra* C^{op} with coproduct

$$\psi^{\text{op}} = T \circ \psi.$$

So on writing $\psi(c) = \sum_i c'_i \otimes c''_i$, we have

$$\psi^{\text{op}}(c^{\text{op}}) = \sum_i (c''_i)^{\text{op}} \otimes (c'_i)^{\text{op}}.$$

The identity function $C \rightarrow C^{\text{op}}$ is a coalgebra homomorphism if and only if C is cocommutative.

REMARK 2.5 (Sweedler notation). Coalgebraists often use the notations

$$\psi(c) = \sum c_{(1)} \otimes c_{(2)} = \sum c_1 \otimes c_2$$

and even drop the summation sign (this is like the *Einstein summation convention* used with tensors). This notation is quite convenient in calculations especially as an alternative to working with humongous commutative diagrams. For example, the coassociativity condition is equivalent to the identity

$$(2.7) \quad (\psi \otimes \text{Id}) \circ \psi(c) = \sum (c_{(1)})_{(1)} \otimes (c_{(1)})_{(2)} \otimes c_{(2)} = \sum c_{(1)} \otimes (c_{(2)})_{(1)} \otimes (c_{(2)})_{(2)} = (\text{Id} \otimes \psi) \circ \psi(c),$$

while the counit conditions become

$$(2.8) \quad \sum \varepsilon(c_{(1)})c_{(2)} = c = \sum \varepsilon(c_{(2)})c_{(1)}.$$

For coalgebras C_1, C_2 , their tensor product $C_1 \otimes C_2$ becomes a coalgebra whose coproduct and counit are the compositions

$$\begin{aligned} C_1 \otimes C_2 &\xrightarrow{\psi_1 \otimes \psi_2} (C_1 \otimes C_1) \otimes (C_2 \otimes C_2) \xrightarrow[\cong]{\text{Id}_{C_1} \otimes \text{T} \otimes \text{Id}_{C_2}} (C_1 \otimes C_2) \otimes (C_1 \otimes C_2), \\ C_1 \otimes C_2 &\xrightarrow[\cong]{\varepsilon_1 \otimes \varepsilon_2} \mathbb{k} \otimes \mathbb{k} \xrightarrow[\cong]{} \mathbb{k}. \end{aligned}$$

So if $c' \in C_1$ and $c'' \in C_2$ with coproducts

$$\psi_{C_1}(c') = \sum c'_{(1)} \otimes c'_{(2)}, \quad \psi_{C_2}(c'') = \sum c''_{(1)} \otimes c''_{(2)},$$

the coproduct on $c' \otimes c'' \in C_1 \otimes C_2$ is

$$\psi_{C_1 \otimes C_2}(c' \otimes c'') = \sum (c'_{(1)} \otimes c''_{(1)}) \otimes (c'_{(2)} \otimes c''_{(2)}) \in (C_1 \otimes C_2) \otimes (C_1 \otimes C_2).$$

Dually to Theorem 2.4 we have

THEOREM 2.6. *Coalgebras and cocommutative coalgebras form symmetric monoidal categories $\mathbf{Coalg}_{\mathbb{k}}$ and $\mathbf{Coalg}_{\mathbb{k}}^{\text{co}}$ under \otimes . In $\mathbf{Coalg}_{\mathbb{k}}^{\text{co}}$, \otimes is the product and \mathbb{k} is a terminal object. In $\mathbf{Coalg}_{\mathbb{k}}$ and $\mathbf{Coalg}_{\mathbb{k}}^{\text{co}}$ the Cartesian product $\times = \oplus$ is the coproduct.*

Here are some basic examples of coalgebras, many more appear in Chapter 3.

EXAMPLE 2.7. Consider the vector space C with basis $\gamma_0, \gamma_1, \gamma_2, \dots$. This can be made into a cocommutative coalgebra with coproduct given by

$$\psi(\gamma_n) = \sum_{0 \leq i \leq n} \gamma_i \otimes \gamma_{n-i}$$

and counit

$$\varepsilon(\gamma_n) = \begin{cases} 1 & \text{if } n = 0, \\ 0 & \text{if } n > 0. \end{cases}$$

This is often called the *Divided Power coalgebra* because we can think of γ_n as representing $X^n/n!$ and the coproduct as coming from

$$X^n \mapsto (X \otimes 1 + 1 \otimes X)^n.$$

EXAMPLE 2.8. Suppose that $(M, \mu, 1)$ is a finite monoid. Then the vector space $\text{Map}(M, \mathbb{k})$ of functions $M \rightarrow \mathbb{k}$ becomes a coalgebra under the coproduct

$$\psi: \text{Map}(M, \mathbb{k}) \rightarrow \text{Map}(M \times M, \mathbb{k}) \cong \text{Map}(M, \mathbb{k}) \otimes \text{Map}(M, \mathbb{k})$$

with

$$\psi(f) = f \circ \mu \longleftrightarrow \sum_{(x,y) \in M^2} f(xy) \delta_x \otimes \delta_y,$$

where $\delta_x \in \text{Map}(M, \mathbb{k})$ is given by

$$\delta_x(y) = \begin{cases} 1 & \text{if } y = x, \\ 0 & \text{if } y \neq x. \end{cases}$$

2.3. Dualising between coalgebras and algebras

The diagrams satisfied by the structure morphisms of algebras and coalgebras are dual in the sense that they are related by ‘reversing all the arrows’. We can exploit this categorical symmetry to dualise coalgebras to algebras and sometimes algebras to coalgebras.

Given a coalgebra (C, ψ, ε) the dual space $C^* = \text{hom}(C, \mathbb{k})$ becomes an algebra by defining the product $C^* \otimes C^* \rightarrow C^*$ to be the following composition.

$$\begin{array}{ccccc} C^* \otimes C^* & \xrightarrow{\quad} & (C \otimes C)^* & \xrightarrow{\psi^*} & C^* \\ & & \parallel & & \parallel \\ & & \text{hom}(C \otimes C, \mathbb{k}) & & \text{hom}(C, \mathbb{k}) \end{array}$$

On elements, for $\alpha, \beta \in C^*$ and $c \in C$,

$$(\alpha\beta)(c) = \sum \alpha(c_{(2)})\beta(c_{(1)})$$

where we use Sweedler notation for the coproduct on c ; notice the switch in order of the indices which is a consequence of our definition of the dual of a tensor product (1.6). The unit is the dual of the counit ε^* ,

$$\varepsilon^*: \mathbb{k} \rightarrow C^*; \quad \varepsilon^*(t) = t\varepsilon.$$

To see that this product on C^* is associative using Sweedler notation, for $\alpha, \beta, \gamma \in C^*$ and $c \in C$, by (2.7)

$$\begin{aligned} ((\alpha\beta)\gamma)(c) &= \sum (\alpha\beta)(c_{(2)})\gamma(c_{(1)}) \\ &= \sum \alpha((c_{(2)})_{(2)})\beta((c_{(2)})_{(1)})\gamma(c_{(1)}) \\ &= \sum \alpha(c_{(2)})\beta((c_{(1)})_{(2)})\gamma((c_{(1)})_{(1)}) \\ &= \sum \alpha(c_{(2)})(\beta\gamma)(c_{(1)}) \\ &= (\alpha(\beta\gamma))(c), \end{aligned}$$

showing that $(\alpha\beta)\gamma = \alpha(\beta\gamma)$; alternatively we could do this with a humongous commutative diagram. A similar calculation using (2.8) shows that ε^* is a unit. Also, if C is cocommutative then C^* is commutative. An important fact is that the algebra C^* acts on C to make it a left C^* -module, as we will see later.

We summarise this discussion in a result.

PROPOSITION 2.9. *Given a coalgebra $(C, \psi_C, \varepsilon_C)$, there is a dual algebra $(C^*, \psi_C^*, \varepsilon_C^*)$. Moreover, C^* is commutative if and only if C is cocommutative.*

A homomorphism of coalgebras $\theta: (C, \psi_C, \varepsilon_C) \rightarrow (C', \psi_{C'}, \varepsilon_{C'})$ induces a homomorphism of algebras $\theta^: ((C')^*, \psi_{C'}^*, \varepsilon_{C'}^*) \rightarrow (C^*, \psi_C^*, \varepsilon_C^*)$.*

PROOF. The last part is left as an exercise. □

If (A, φ, η) is an algebra which is finite dimensional then we can also dualise to get a coalgebra (A^*, φ^*, η^*) . However, if A is infinite dimensional we need to modify the notion of dual appropriately to make this work. There are two ways to do this: the more drastic one involves introducing linearly topologised vector spaces and a notion of completeness, the other

leads to a more ‘algebraic’ outcome by suitably restricting the elements in the dual space. We will take the latter approach, see Remark 2.11 for more on the alternative.

Suppose that V is a vector space. Then a subspace $U \subseteq V$ is *cofinite* or has *finite codimension* if $\dim_{\mathbb{k}} V/U < \infty$.

For an algebra A , we define its *finite* or *restricted dual* by

$$A^\circ = \{\alpha \in A^* : \text{there is a cofinite } I \triangleleft A \text{ such that } I \subseteq \ker \alpha\} \subseteq A^*.$$

So $\alpha \in A^*$ is in A° if it factors through a finite dimensional quotient algebra A/I . Of course when A is finite dimensional, $A^\circ = A^*$, but otherwise $A^\circ \subsetneq A^*$. As before we can ‘dualise’ the algebra structure on A to obtain a coproduct $\varphi^*: A^* \rightarrow (A \otimes A)^*$ but in order to land in a tensor product we need to restrict it to A° using the fact that there is a commutative diagram

$$\begin{array}{ccc} A^\circ & \xrightarrow{\varphi^\circ} & A^\circ \otimes A^\circ \xleftarrow{\cong} (A \otimes A)^\circ \\ \downarrow & & \downarrow \\ A^* & \xrightarrow{\varphi^*} & (A \otimes A)^* \end{array}$$

and the dotted arrow is defined to be the coproduct $\varphi^\circ: A^\circ \rightarrow A^\circ \otimes A^\circ$. There is a counit $\eta^\circ: A^\circ \rightarrow \mathbb{k}$ obtained by precomposing with η . Then $(A^\circ, \varphi^\circ, \eta^\circ)$ is a coalgebra.

PROPOSITION 2.10. *Given an algebra (A, φ_A, η_A) , there is a dual coalgebra $(A^\circ, \varphi_A^\circ, \eta_A^\circ)$. Moreover, A° is cocommutative if and only if A is commutative.*

A homomorphism of algebras $\theta: (A, \varphi_A, \eta_A) \rightarrow (A', \varphi_{A'}, \eta_{A'})$ induces a homomorphism of coalgebras $\theta^\circ: ((A')^\circ, \varphi_{A'}^\circ, \eta_{A'}^\circ) \rightarrow (A^\circ, \varphi_A^\circ, \eta_A^\circ)$.

REMARK 2.11. One way to relate this approach to the other is by noting that given two cofinite ideals $I, J \triangleleft A$, if $I \subseteq J$ there is an induced surjective algebra homomorphism $A/I \rightarrow A/J$; these fit together to form an inverse system whose limit is the *profinite completion* of A ,

$$\widehat{A} = \lim_{I \triangleleft A \text{ cofin.}} A/I.$$

This can be given a certain Hausdorff *profinite topology* so that it is Cauchy complete.

When we take linear duals we get linear maps $(A/J)^* \rightarrow (A/I)^*$ which form an directed system whose colimit is


$$A^\circ = \operatorname{colim}_{I \triangleleft A \text{ cofin.}} (A/I)^*.$$

This is not the linear dual of \widehat{A} but it is the *continuous linear dual* with respect to the *profinite topology* on \widehat{A} and \mathbb{k} (which has the discrete topology). In the profinite topology the open sets are the cosets $a + J$ where $J \triangleleft \widehat{A}$ is a cofinite ideal (these are actually clopen); a continuous linear map $\widehat{A} \rightarrow \mathbb{k}$ is one that factors through a finite dimensional quotient \widehat{A}/J .


2.4. Bialgebras

In order to define a *bialgebra* (also known as a *hyperalgebra* or *bigèbre* in French) we need a vector space equipped with both an algebra and a coalgebra structure, (B, φ, η) and (B, ψ, ε) , which interact appropriately.

Recall that we can give $B \otimes B$ an algebra structure and a coalgebra structure, so it makes sense to ask if $\varphi: B \otimes B \rightarrow B$ and $\eta: \mathbb{k} \rightarrow B$ are coalgebra homomorphisms or if $\psi: B \rightarrow B \otimes B$ and $\varepsilon: B \rightarrow \mathbb{k}$ are algebra homomorphisms. Either of these amounts to requiring that chasing around the following diagrams read from left to right gives the same output.



Here the product for $B \otimes B$ is shown in red, the coproduct in green.



We also have two commutative diagrams for the unit and counit.

$$\begin{array}{ccc}
 \mathbb{k} & \xrightarrow{\cong} & \mathbb{k} \otimes \mathbb{k} \\
 \eta \downarrow & & \downarrow \eta \otimes \eta \\
 B & \xrightarrow{\psi} & B \otimes B
 \end{array}
 \qquad
 \begin{array}{ccc}
 B \otimes B & \xrightarrow{\varphi} & B \\
 \varepsilon \otimes \varepsilon \downarrow & & \downarrow \varepsilon \\
 \mathbb{k} \otimes \mathbb{k} & \xrightarrow{\cong} & \mathbb{k}
 \end{array}$$

If necessary, we denote the structure maps in a bialgebra by writing $(B, \varphi, \eta, \psi, \varepsilon)$.

Of course a homomorphism of bialgebras should be simultaneously an algebra and a coalgebra homomorphism, and there is a category of bialgebras $\mathbf{Bialg}_{\mathbb{k}}$ with full subcategories of commutative and cocommutative bialgebras. If a bialgebra is both commutative and cocommutative then it is called *bicommutative*.

EXAMPLE 2.12 (The Quantum Plane). Let $1 \neq q \in \mathbb{k}$. The *Quantum Plane* is the non-commutative bialgebra

$$\mathcal{O}_q(\mathbb{k}^2) = \mathbb{k}\langle X, Y \rangle / (YX - qXY).$$

We will denote the residue classes of X and Y by x and y , so these satisfy $yx = qxy$; notice that the monomials $x^i y^j$ form a basis of $\mathcal{O}_q(\mathbb{k}^2)$. There is a coproduct ψ and counit ε given by

$$\psi(x) = x \otimes x, \quad \psi(y) = y \otimes 1 + x \otimes y, \quad \varepsilon(x) = 1, \quad \varepsilon(y) = 0.$$

This bialgebra is neither commutative nor cocommutative so it is a *quantum monoid*.

In Example 2.20 this construction will be modified to give a Hopf algebra.

PROPOSITION 2.13. *Suppose that $(B, \varphi, \eta, \psi, \varepsilon)$ is a bialgebra.*

- (a) *If (B, ψ, ε) is a cocommutative coalgebra, then (B, φ, η) is a monoid in $\mathbf{Coalg}_{\mathbb{k}}^{\text{co}}$. In particular, φ and η are coalgebra homomorphisms.*
- (b) *If (B, φ, η) is a commutative algebra then (B, ψ, ε) is a comonoid in $\mathbf{Alg}_{\mathbb{k}}^{\text{co}}$. In particular, ψ and ε are algebra homomorphisms.*

PROOF. (a) In the category $\mathbf{Coalg}_{\mathbb{k}}^{\text{co}}$, \otimes is the categorical product and \mathbb{k} is a terminal object. Now expand the diagrams of (2.1) for (B, φ, η) with $A = B$ and interpret them as being in $\mathbf{Coalg}_{\mathbb{k}}^{\text{co}}$.

(b) In the category $\mathbf{Alg}_{\mathbb{k}}^{\text{co}}$, \otimes is the categorical coproduct and \mathbb{k} is an initial object. Now expand the diagrams of (2.4) for (B, ψ, ε) with $C = B$ and interpret them as being in $\mathbf{Alg}_{\mathbb{k}}^{\text{co}}$. \square

2.5. Interlude: Convolution monoids

In order to define Hopf algebras we will require a construction that can be made using an algebra and a coalgebra as ingredients. Let (A, φ, η) be a \mathbb{k} -algebra and (C, ψ, ε) a \mathbb{k} -coalgebra. The vector space $\text{hom}(C, A)$ can be given a useful product.

DEFINITION 2.14. For $f, g \in \text{hom}(C, A)$, their *convolution product* is

$$f * g = \varphi \circ (f \otimes g) \circ \psi$$

and since this function $C \rightarrow A$ is a composition of linear mappings it is an element of $\text{hom}(C, A)$. In Sweedler notation, we have for $c \in C$,


$$f * g(c) = \sum f(c_{(1)})g(c_{(2)}).$$

There is also the distinguished element $\mathbf{1} = \mathbf{1}_{C,A} = \eta \circ \varepsilon \in \text{hom}(C, A)$.


PROPOSITION 2.15. *With the product $*$, $\text{hom}(C, A)$ is a monoid with identity element $\mathbf{1}$. When A is commutative and C is cocommutative, this monoid is commutative.*

PROOF. To show that $\text{hom}(C, A)$ is a monoid we need to check that $*$ is associative and $\mathbf{1}$ acts as the identity.

Associativity and coassociativity of A and C imply the commutativity of the following diagram which proves $(f * g) * h = f * (g * h)$ for any $f, g, h \in \text{hom}(C, A)$.



By contemplating the following commutative diagram we see that $\mathbf{1} * f = f = f * \mathbf{1}$.



The commutativity result is left as an exercise. □

When we combine $*$ with the vector space structure on $\text{hom}(C, A)$ we get an algebra.

COROLLARY 2.16. *The vector space $\text{hom}(C, A)$ becomes a \mathbb{k} -algebra with product $*$ and unity $\mathbf{1}$.*

Given an algebra homomorphism $\alpha: A \rightarrow A'$ and a coalgebra homomorphism $\gamma: C' \rightarrow C$, there are \mathbb{k} -linear mappings

$$\begin{aligned}\alpha_*: \operatorname{hom}(C, A) &\rightarrow \operatorname{hom}(C, A'); & \alpha_*(f) &= \alpha \circ f, \\ \gamma^*: \operatorname{hom}(C, A) &\rightarrow \operatorname{hom}(C', A); & \gamma^*(f) &= f \circ \gamma.\end{aligned}$$

It is easy to verify that these are monoid homomorphisms and so algebra homomorphisms, i.e., for $f, g \in \operatorname{hom}(C, A)$,

$$\begin{aligned}\alpha_*(f * g) &= \alpha_*(f) * \alpha_*(g), & \gamma^*(f * g) &= \gamma^*(f) * \gamma^*(g), \\ \alpha_*(\mathbf{1}) &= \mathbf{1}, & \gamma^*(\mathbf{1}) &= \mathbf{1}.\end{aligned}$$

In a monoid elements need not have inverses, but sometimes they do. If $f \in \operatorname{hom}(C, A)$ then $\bar{f} \in \operatorname{hom}(C, A)$ is an inverse for f if

$$\bar{f} * f = \mathbf{1} = f * \bar{f},$$

or more explicitly if for every $c \in C$, using Sweedler notation in A we have

$$\sum \bar{f}(c_{(1)})f(c_{(2)}) = \varepsilon(c) = \sum f(c_{(1)})\bar{f}(c_{(2)}).$$

Of course, if it exists then such a two-sided inverse for f is unique by a standard argument that makes use of associativity.

Notice that when $A = \mathbb{k}$, $C^* = \operatorname{hom}(C, \mathbb{k})$ and $\mathbf{1} = \varepsilon^*$, and the algebra $(C^*, *, \mathbf{1})$ agrees with the algebra $(C^*, \psi^*, \varepsilon^*)$ discussed earlier.

In order to define Hopf algebras, we require some observations on inverses in convolution monoids for bialgebras.

LEMMA 2.17. *Suppose that B is a bialgebra.*

- (a) *If A is an algebra and $f: B \rightarrow A$ is an algebra homomorphism which has a convolution inverse \bar{f} in $\operatorname{hom}(B, A)$, then \bar{f} is an algebra homomorphism $B \rightarrow A^{\operatorname{op}}$.*
- (b) *If C is a coalgebra and $g: C \rightarrow B$ is a coalgebra homomorphism which has a convolution inverse \bar{g} in $\operatorname{hom}(C, B)$, then \bar{g} is a coalgebra homomorphism $C^{\operatorname{op}} \rightarrow B$.*

PROOF. (a) Let $B \otimes B$ with its product $\varphi_{B \otimes B}$ which is also a coalgebra homomorphism with respect to its coproduct $\psi_{B \otimes B}$. This means that $\varphi_{B \otimes B}^*: \operatorname{hom}(B, A) \rightarrow \operatorname{hom}(B \otimes B, A)$ is a monoid homomorphism and in particular $\varphi_{B \otimes B}^*(f) \in \operatorname{hom}(B \otimes B, A)$ has inverse $\varphi_{B \otimes B}^*(\bar{f})$.

Now define $\ell = \varphi_A \circ (\bar{f} \otimes \bar{f}) \circ T: B \otimes B \rightarrow A$, given on elements by

$$\ell(x \otimes y) = \bar{f}(y)\bar{f}(x).$$

We will show that ℓ is also a left inverse for $\varphi_{B \otimes B}^*(f)$ and therefore it agrees with $\varphi_{B \otimes B}^*(\bar{f})$. To verify this we calculate: for $x, y \in B$,

$$\begin{aligned}(\ell * \varphi_{B \otimes B}^*(f))(x \otimes y) &= \sum \sum \ell(x_{(1)} \otimes y_{(1)})\varphi_{B \otimes B}^*(f)(x_{(2)} \otimes y_{(2)}) \\ &= \sum \sum \bar{f}(y_{(1)})\bar{f}(x_{(1)})f(x_{(2)}y_{(2)}) \\ &= \sum \sum \bar{f}(y_{(1)})\bar{f}(x_{(1)})f(x_{(2)})f(y_{(2)}) \\ &= \sum \bar{f}(y_{(1)})(\bar{f} * f)(x)f(y_{(2)}) \\ &= \sum \bar{f}(y_{(1)})\varepsilon(x)f(y_{(2)}) \\ &= \varepsilon(x) \sum \bar{f}(y_{(1)})f(y_{(2)}) \\ &= \varepsilon(x)(\bar{f} * f)(y) \\ &= \varepsilon(x)\varepsilon(y) = \varepsilon(xy).\end{aligned}$$

So $(\ell * \varphi_{B \otimes B}^*(f)) = 1$ and ℓ is the inverse of $\varphi_{B \otimes B}^*(f)$.

The proof of (b) is similar. \square

In particular, when B is a bialgebra, the identity function $\text{Id}_B: B \rightarrow B$ is both an algebra homomorphism and a coalgebra homomorphism, so if it has a convolution inverse $\overline{\text{Id}}_B \in \text{hom}(B, B)$ this is both an algebra isomorphism $B \rightarrow B^{\text{op}}$ and a coalgebra isomorphism $B^{\text{op}} \rightarrow B$.

LEMMA 2.18. *Suppose that $(B, \varphi, \eta, \psi, \varepsilon)$ is a bialgebra which is either commutative or co-commutative and that $\overline{\text{Id}}_B$ exists. Then $\overline{\text{Id}}_B: B \rightarrow B$ is self-inverse, i.e.,*

$$\overline{\text{Id}}_B \circ \overline{\text{Id}}_B = \text{Id}_B.$$

PROOF. We will give the proof when B is commutative, the other case is similar. So Id_B is an isomorphism $B \cong B^{\text{op}}$ and by Lemma 2.17(a), $\overline{\text{Id}}_B: B \rightarrow B$ is an algebra homomorphism, hence $\varphi \circ (\overline{\text{Id}}_B \otimes \overline{\text{Id}}_B) = \overline{\text{Id}}_B \circ \varphi$. To identify $\overline{\text{Id}}_B \circ \overline{\text{Id}}_B$ it is sufficient to show that

$$(\overline{\text{Id}}_B \circ \overline{\text{Id}}_B) * \overline{\text{Id}}_B = \mathbf{1}.$$

We have

$$\begin{aligned} (\overline{\text{Id}}_B \circ \overline{\text{Id}}_B) * \overline{\text{Id}}_B &= \varphi \circ ((\overline{\text{Id}}_B \circ \overline{\text{Id}}_B) \otimes \overline{\text{Id}}_B) \circ \psi \\ &= \varphi \circ (\overline{\text{Id}}_B \otimes \overline{\text{Id}}_B) \circ (\text{Id}_B \otimes \overline{\text{Id}}_B) \circ \psi \\ &= \overline{\text{Id}}_B \circ \varphi \circ (\text{Id}_B \otimes \overline{\text{Id}}_B) \circ \psi \\ &= \overline{\text{Id}}_B \circ (\text{Id}_B * \overline{\text{Id}}_B) \\ &= \overline{\text{Id}}_B \circ \mathbf{1} \\ &= \overline{\text{Id}}_B \circ \eta \circ \varepsilon = \mathbf{1}, \end{aligned}$$

and so $\overline{\text{Id}}_B \circ \overline{\text{Id}}_B = \text{Id}_B$ as required. \square

2.6. Hopf algebras

Finally we are ready to define a Hopf algebra.

DEFINITION 2.19. If $(H, \varphi, \eta, \psi, \varepsilon)$ is a bialgebra for which $\chi = \overline{\text{Id}}_H$ exists then it is called the *antipode* of H and $(H, \varphi, \eta, \psi, \varepsilon, \chi)$ is called a *Hopf algebra*. In many sources χ is denoted by S .

The antipode χ has to satisfy some conditions which we can encode in the following commutative diagram.

(2.9)

$$\begin{array}{ccccc} & & H & & \\ & \swarrow \psi & \downarrow \varepsilon & \searrow \psi & \\ H \otimes H & & \mathbb{k} & & H \otimes H \\ \downarrow \chi \otimes \text{Id} & & \downarrow \eta & & \downarrow \text{Id} \otimes \chi \\ H \otimes H & & H & & H \otimes H \\ & \searrow \varphi & \downarrow \varphi & \swarrow \varphi & \\ & & H & & \end{array}$$

On an element $h \in H$ this expands to give

$$(2.10) \quad \sum \chi(h_{(1)})h_{(2)} = \varepsilon(h) = \sum h_{(1)}\chi(h_{(2)}).$$

In general $\chi: H \rightarrow H$ is not a bijective function, however if it is bijective then its inverse function χ^{-1} fits into the two equivalent commutative diagrams

$$(2.11) \quad \begin{array}{ccc} & H & \\ \psi \swarrow & & \searrow \psi \\ H \otimes H & \xrightarrow{\varepsilon} & H \otimes H \\ \downarrow T & & \downarrow T \\ H \otimes H & \xrightarrow{\chi^{-1} \otimes \text{Id}} & H \otimes H \\ \downarrow \chi^{-1} \otimes \text{Id} & & \downarrow \text{Id} \otimes \chi^{-1} \\ H \otimes H & \xrightarrow{\varphi} & H \otimes H \\ & \searrow \varphi & \swarrow \varphi \\ & H & \end{array} \quad \begin{array}{ccc} & H & \\ \psi \swarrow & & \searrow \psi \\ H \otimes H & \xrightarrow{\varepsilon} & H \otimes H \\ \downarrow \text{Id} \otimes \chi^{-1} & & \downarrow \chi^{-1} \otimes \text{Id} \\ H \otimes H & \xrightarrow{T} & H \otimes H \\ \downarrow T & & \downarrow T \\ H \otimes H & \xrightarrow{\varphi} & H \otimes H \\ & \searrow \varphi & \swarrow \varphi \\ & H & \end{array}$$

which expand to give

$$(2.12) \quad \sum \chi^{-1}(h_{(2)})h_{(1)} = \varepsilon(h) = \sum h_{(2)}\chi^{-1}(h_{(1)}).$$

EXAMPLE 2.20 (The localised Quantum Plane). We can modify the Quantum Plane of Example 2.12 to give a Hopf algebra by forcing x to have an inverse. Let

$$\mathcal{O}_q(\mathbb{k}^2)[x^{-1}] = \mathbb{k}\langle X, Y, Z \rangle / (YX - qXY, XZ - 1, ZX - 1).$$

We will denote the residue class of Z by x^{-1} .

The coproduct and counit of $\mathcal{O}_q(\mathbb{k}^2)$ extend to $\mathcal{O}_q(\mathbb{k}^2)[x^{-1}]$ so that

$$\psi(x^{-1}) = x^{-1} \otimes x^{-1}, \quad \varepsilon(x^{-1}) = 1,$$

and the antipode is given by

$$\chi(x) = x^{-1}, \quad \chi(x^{-1}) = x, \quad \chi(y) = -x^{-1}y.$$

This is a Hopf algebra which is neither commutative nor cocommutative. It has interesting finite dimensional quotient Hopf algebras when q takes special values; these are called *Taft algebras* and are discussed in Chapter 3.

DEFINITION 2.21. A *homomorphism* of Hopf algebras or *Hopf homomorphism*

$$\theta: (H, \varphi, \eta, \psi, \varepsilon, \chi) \rightarrow (H', \varphi', \eta', \psi', \varepsilon', \chi')$$

is a \mathbb{k} -linear mapping $\theta: H \rightarrow H'$ which is both an algebra and a coalgebra homomorphism. A Hopf homomorphism which is invertible is called an *isomorphism*.

Just as a group homomorphism maps inverses to inverses, such a Hopf homomorphism also satisfies

$$\theta \circ \chi = \chi' \circ \theta.$$

The kernel of a Hopf algebra homomorphism θ is both an ideal and a coideal, which is also closed under the restriction of the antipode of the domain. Such an ideal in a Hopf algebra is called a *Hopf ideal*. It is easy to see if $J \triangleleft H$ is a Hopf ideal then there are unique algebra and coalgebra structures on H/J so that the quotient map $H \rightarrow H/J$ is a homomorphism of Hopf algebras; then H/J is called the *quotient Hopf algebra of H with respect to J* .

PROPOSITION 2.22. Let $\theta: (H, \varphi, \eta, \psi, \varepsilon, \chi) \rightarrow (H', \varphi', \eta', \psi', \varepsilon', \chi')$ be a Hopf homomorphism. Then

(a) $\chi' \circ \theta = \theta \circ \chi$;

(b) $\ker \theta \triangleleft H$ is a Hopf ideal and the image of θ is a subHopf algebra of H' is isomorphic to the quotient Hopf algebra $H/\ker \theta$.

PROOF. (a) The idea is to show that in the convolution monoid $\text{hom}(H, H')$ the elements $\chi' \circ \theta$ and $\theta \circ \chi$ satisfy .

$$(\chi' \circ \theta) * \theta = \eta' \circ \varepsilon = \theta * (\chi' \circ \theta)$$

and

$$(\theta \circ \chi) * \theta = \eta' \circ \varepsilon = \theta * (\theta \circ \chi)$$

where $\chi' \circ \theta$ is the identity element. This shows that these elements are both inverses of θ and so must be equal by uniqueness of inverses. Here is a sample, the others follow by similar calculations:

$$\begin{aligned} (\chi' \circ \theta) * \theta &= \varphi' \circ ((\chi' \circ \theta) \otimes \theta) \circ \psi \\ &= \varphi' \circ (\chi' \otimes \text{Id}) \circ (\theta \otimes \theta) \circ \psi \\ &= \varphi' \circ (\chi' \otimes \text{Id}) \circ \psi' \circ \theta \\ &= (\chi' * \text{Id}) \circ \theta \\ &= \eta' \circ \varepsilon' \circ \theta \\ &= \eta' \circ \varepsilon. \end{aligned}$$

(b) This is a consequence of earlier results about homomorphisms of algebras and coalgebras. \square

REMARK 2.23. Of course Hopf algebras over \mathbb{k} and their homomorphisms define a category $\mathbf{HA}_{\mathbb{k}}$ which has the null object \mathbb{k} . There are three obvious full subcategories whose objects are the commutative, the cocommutative and the bicommutative Hopf algebras. In the first two, \otimes is the categorical coproduct and product respectively. The category of bicommutative Hopf algebras (also known as *abelian Hopf algebras*) has many features possessed by an abelian category (for example \otimes is the both the categorical coproduct and product), and indeed appropriate subcategories such as finite dimensional bicommutative Hopf algebras do form abelian categories.

We mention one important example of an isomorphism.

EXAMPLE 2.24. Suppose that $(H, \varphi, \eta, \psi, \varepsilon, \chi)$ is a Hopf algebra whose antipode χ is bijective. Then its *opposite Hopf algebra* is $(H^{\text{op}}, \varphi^{\text{op}}, \eta^{\text{op}}, \psi^{\text{op}}, \varepsilon^{\text{op}}, \chi^{\text{op}})$ where we take the opposite algebra and coalgebra structures and as a function $\chi^{\text{op}} = \chi$. Then the function

$$\tilde{\chi}: H \rightarrow H^{\text{op}}; \quad \tilde{\chi}(h) = (\chi(h))^{\text{op}}$$

is an isomorphism of Hopf algebras with inverse

$$\widetilde{\chi^{\text{op}}}: H^{\text{op}} \rightarrow H; \quad \widetilde{\chi^{\text{op}}}(h^{\text{op}}) = \chi^{-1}(h).$$

A similar result applies if we interchange χ and χ^{-1} .

Later we will see that these isomorphisms allows us to interchange between left and right modules and comodules over H .

PROPOSITION 2.25. Suppose that $(H, \varphi, \eta, \psi, \varepsilon, \chi)$ is a Hopf algebra.

(a) If (H, ψ, ε) is cocommutative then (H, φ, η, χ) is a group object in $\mathbf{Coalg}_{\mathbb{k}}^{\text{co}}$. In particular, φ, η, χ are coalgebra homomorphisms.

(b) If (H, φ, η) is commutative then $(H, \psi, \varepsilon, \chi)$ is a cogroup object in $\mathbf{Alg}_k^{\text{co}}$. In particular, ψ, ε, χ are algebra homomorphisms.

PROOF. This follows from Proposition 2.13 since χ is the inverse map in each case. \square

DEFINITION 2.26. A Hopf algebra which is commutative or cocommutative is called a *classical Hopf algebra*. A Hopf algebra which is both commutative and cocommutative is sometimes called *bicommutative* or *abelian*. A Hopf algebra for which $\chi \circ \chi = \text{Id}$ is called *involutive* or *involutive*.

REMARK 2.27. Notice that involutive Hopf algebras have bijective antipodes; in general the antipode of a Hopf algebra need not be bijective although it often is. We have shown above that for a classical Hopf algebra, $\chi \circ \chi = \text{Id}$ and χ is an (co)algebra isomorphism $H \xrightarrow{\cong} H^{\text{op}}$ to the opposite (co)algebra.

REMARK 2.28. Although in general the antipode χ of a Hopf algebra H need not be either an algebra or a coalgebra homomorphism, its composition square $\chi^2 = \chi \circ \chi$ is by Lemma 2.17 and because χ^2 commutes with χ . This means that $\chi^2 H \subseteq H$ is a subHopf algebra; of course χ is not injective or surjective this might be a proper inclusion of a quotient Hopf algebra.

Some special kinds of elements in a Hopf algebra.

DEFINITION 2.29. If H is a Hopf algebra then its set of *primitive elements* is

$$P(H) = \{h \in H : \psi(h) = 1 \otimes h + h \otimes 1\}.$$

This is a vector subspace of H , but it also has other properties. For $x, y \in P(H)$,

$$\begin{aligned} \psi(xy - yx) &= \psi(x)\psi(y) - \psi(y)\psi(x) \\ &= (x \otimes 1 + 1 \otimes x)(y \otimes 1 + 1 \otimes y) - (y \otimes 1 + 1 \otimes y)(x \otimes 1 + 1 \otimes x) \\ &= (xy \otimes 1 + x \otimes y + y \otimes x + 1 \otimes xy) - (yx \otimes 1 + y \otimes x + x \otimes y + 1 \otimes yx) \\ &= (xy - yx) \otimes 1 + 1 \otimes (xy - yx), \end{aligned}$$

hence $xy - yx \in P(H)$. This shows that $P(H)$ is a Lie subalgebra of H equipped with its commutator bracket. Notice also that if $x \in P(H)$ then

$$x = \varepsilon(1)x + \varepsilon(x) = x + \varepsilon(x)$$

so $\varepsilon(x) = 0$, hence $P(H) \subseteq \ker \varepsilon$.

In fact P defines a functor $\mathbf{HA}_k \rightarrow \mathbf{Lie}_k$ to the category of Lie algebras over k and this has a left adjoint. This will be discussed further in Chapter 3.

DEFINITION 2.30. If H is a Hopf algebra then a non-zero element $g \in H$ is *group-like* if

$$\psi(g) = g \otimes g,$$

and its set of all group-like elements is

$$G(H) = \{g \in H : g \text{ is group-like}\}.$$

If $g, h \in G(H)$ then

$$\psi(gh) = gh \otimes gh$$

and since $\psi(1) = 1 \otimes 1$, $1 \in G(H)$. This show that $G(H)$ is a monoid under multiplication. If $g \in G(H)$ then using the counit we get

$$\varepsilon(g)g = g = g\varepsilon(g)$$

so $\varepsilon(g) = 1$; now using the antipode we also find that

$$\chi(g)g = \varepsilon(g) = g\chi(g)$$

so g is a unit with inverse $g^{-1} = \chi(g)$. Therefore $G(H) \leq H^\times$.

There is a more general notion that combines the group-like and the primitives. If $g \in G(H)$ then the set of g -primitives is

$$P_g(H) = \{h \in H : \psi(h) = g \otimes h + h \otimes g\}.$$

LEMMA 2.31. *Let H be a Hopf algebra. Then the set of group-like elements $G(H)$ is linearly independent. Hence the group-like elements span a cocommutative subHopf algebra isomorphic to the group algebra $\mathbb{k}G(H)$.*

PROOF. Suppose that $G(H)$ is not linearly independent. Then there is a minimal $n \geq 1$ for which there is a subset $\{g_0, g_1, \dots, g_n\} \subseteq G(H)$ with $\{g_1, \dots, g_n\}$ linearly independent and

$$g_0 = \sum_{1 \leq k \leq n} t_k g_k$$

for $t_k \in \mathbb{k}$. Applying ψ we obtain

$$g_0 \otimes g_0 = \sum_{1 \leq k \leq n} t_k g_k \otimes g_k \in H \otimes H$$

and so

$$\sum_{\substack{1 \leq k \leq n \\ 1 \leq \ell \leq n}} t_k t_\ell g_k \otimes g_\ell = \sum_{1 \leq k \leq n} t_k g_k \otimes g_k.$$

Since the basic tensors $g_k \otimes g_\ell \in H \otimes H$ are linearly independent we must have $t_k = 0$. This contradiction shows that no such minimal set exists.

The monoid $G(H)$ spans a subspace with its elements as a basis, and which is closed under multiplication it forms a subalgebra visibly isomorphic to the group algebra $\mathbb{k}G(H)$. Also the coproduct ψ restricts to it and agrees with the coproduct in the group algebra. Finally, it is closed under the action of the antipode. \square

In fact $G(-)$ defines a functor $\mathbf{G}: \mathbf{HA}_{\mathbb{k}} \rightarrow \mathbf{Gp}$ and this has as its left adjoint the group algebra functor $\mathbb{k}(-): \mathbf{Gp} \rightarrow \mathbf{HA}_{\mathbb{k}}$, so there is a natural isomorphism of bifunctors

$$\mathbf{HA}_{\mathbb{k}}(\mathbb{k}(-), -) \cong \mathbf{Gp}(-, G(-)).$$

This will be discussed more in Chapter 3.

2.7. SubHopf algebras, adjoint actions and normal subalgebras

A Hopf algebra H can contain subalgebras, subcoalgebras and subbialgebras. A subbialgebra $K \subseteq H$ where the antipode χ restricts to give an antipode for K is called a *subHopf algebra*; of course K is then a Hopf algebra in its own right.

$$\begin{array}{ccc} K & \xrightarrow{\chi|_K} & K \\ \downarrow & & \downarrow \\ H & \xrightarrow{\chi} & H \end{array}$$

This is analogous to the notion of a subgroup of a group. In fact the group algebra of a subgroup $H \leq G$ is a subHopf algebra $\mathbb{k}H \subseteq \mathbb{k}G$.

PROPOSITION 2.32. *The image of the antipode $\chi H \subseteq H$ is a subHopf algebra. More generally, for $n \geq 2$, $\chi^n H \subseteq H$ is a subHopf algebra.*

PROOF. We know that for H the identities

$$\psi \circ \chi = (\chi \otimes \chi) \circ T \circ \psi, \quad \chi \circ \varphi = \varphi \circ T \circ (\chi \otimes \chi),$$

which imply

$$\varphi(\chi H \otimes \chi H) \subseteq \chi H, \quad \psi \chi H \subseteq \chi H \otimes \chi H,$$

hence χH is a subbialgebra of H .

We also have

$$\begin{aligned} (\chi * \text{Id}) \circ \chi &= \varphi \circ (\chi \otimes \text{Id}) \circ \psi \circ \chi \\ &= \varphi \circ (\chi \otimes \text{Id}) \circ (\chi \otimes \chi) \circ T \circ \psi \\ &= \varphi \circ (\chi \otimes \chi) \circ (\chi \otimes \text{Id}) \circ T \circ \psi \\ &= \varphi \circ (\chi \otimes \chi) \circ T \circ (\text{Id} \otimes \chi) \circ \psi \\ &= \chi \circ \varphi \circ (\chi \otimes \text{Id}) \circ \psi \\ &= \chi \circ \varphi \circ (\chi * \text{Id}) \\ &= \chi \circ \eta \circ \varepsilon = \eta \circ \varepsilon, \end{aligned}$$

and a similar calculation shows that $(\text{Id} * \chi) \circ \chi = \eta \circ \varepsilon$. These identities show that the restriction of χ to χH is an antipode for it, therefore χH is a subHopf algebra of H . \square

Now we will consider the analogue of a *normal* subgroup. There are two approaches which roughly correspond to the two ways of thinking about when a subgroup is normal (i.e., requiring left and right cosets to be equal, or being closed under conjugation).

Let $A \subseteq H$ be a subalgebra and let $A^+ = \ker \varepsilon_A$, the kernel of the counit restricted to A . Then $HA^+ \subseteq H$ is a left ideal and $A^+H \subseteq H$ is a right ideal. If $HA^+ = A^+H$ we can form the quotient algebra H/HA^+ , but this won't always be a Hopf algebra. If $K \subseteq H$ is a subHopf algebra and if $HK^+ = K^+H$, this is also a coideal and H/HK^+ is a quotient Hopf algebra. So this looks like a reasonable way to define a 'normal' subHopf algebra.

The alternative approach requires the two *adjoint actions*.

DEFINITION 2.33. For $h \in H$, the left and right *adjoint actions* $\text{ad}_h^l: H \rightarrow H$ and $\text{ad}_h^r: H \rightarrow H$ are given by

$$\text{ad}_h^l(x) = \sum h_{(1)}x\chi(h_{(2)}), \quad \text{ad}_h^r(x) = \sum \chi(h_{(1)})xh_{(2)}.$$

LEMMA 2.34. *The adjoint actions are left and right actions of H on itself, i.e., for $h', h'' \in H$,*

$$\text{ad}_{h'h''}^l = \text{ad}_{h'}^l \circ \text{ad}_{h''}^l, \quad \text{ad}_{h'h''}^r = \text{ad}_{h''}^r \circ \text{ad}_{h'}^r.$$

Furthermore, for $h, x, y \in H$,

$$\begin{aligned} \text{ad}_h^l(xy) &= \sum \text{ad}_{h_{(1)}}^l(x)\text{ad}_{h_{(2)}}^l(y), & \varepsilon(\text{ad}_h^l(x)) &= \varepsilon(h)\varepsilon(x), & \text{ad}_h^l(1) &= 1, \\ \text{ad}_h^r(xy) &= \sum \text{ad}_{h_{(1)}}^r(x)\text{ad}_{h_{(2)}}^r(y), & \varepsilon(\text{ad}_h^r(x)) &= \varepsilon(h)\varepsilon(x), & \text{ad}_h^r(1) &= 1. \end{aligned}$$

The left/right adjoint actions makes H into a left/right module over itself.

For the Hopf algebra $\mathbb{k}G$ of a group, when $g \in G$, the two adjoint actions correspond to left and right conjugations:

$$\text{ad}_g^l = g(-)g^{-1}, \quad \text{ad}_g^r = g^{-1}(-)g.$$

Now we can define a subalgebra $A \subseteq H$ to be *ad-invariant* if for every $h \in H$, $\text{ad}_h^l A \subseteq A$ and $\text{ad}_h^r A \subseteq A$. Although in general this notion involves two independent conditions, for some Hopf algebras such as group algebras the left and right adjoint actions give equivalent information.

LEMMA 2.35. *Suppose that the coproduct ψ is cocommutative. Then the following conditions are equivalent:*

- A is ad-invariant;
- for every $h \in H$, $\text{ad}_h^l A \subseteq A$;
- for every $h \in H$, $\text{ad}_h^r A \subseteq A$.

PROOF. By Lemma 2.18, $\chi: H \rightarrow H$ is a bijection and indeed $\chi^{-1} = \chi$.

Suppose that for every $h \in H$, $\text{ad}_h^l A \subseteq A$. Then for every $a \in A$ and $h \in H$, let $h' = \chi(h)$ so that $h = \chi(h')$ and

$$\begin{aligned} \text{ad}_h^r(a) &= \sum \chi(h_{(1)})ah_{(2)} \\ &= \sum \chi(\chi(h')_{(1)})a\chi(h')_{(2)} \\ &= \sum \chi(\chi(h'_{(2)}))a\chi(h'_{(1)}) \\ &= \sum h'_{(2)}a\chi(h'_{(1)}) \\ &= \sum h'_{(1)}a\chi(h'_{(2)}) = \text{ad}_{h'}^l(a) \in A, \end{aligned}$$

where we have used cocommutativity in the last step. Therefore

$$\forall h \in H, \text{ad}_h^l A \subseteq A \implies \forall h \in H, \text{ad}_h^r A \subseteq A.$$

Similarly,

$$\forall h \in H, \text{ad}_h^r A \subseteq A \implies \forall h \in H, \text{ad}_h^l A \subseteq A. \quad \square$$

REMARK 2.36. In the next result we need the notion of *faithful flatness*: For a \mathbb{k} -algebra R , a right R -module L is *faithfully flat* if every sequence of left R -modules

$$0 \rightarrow M' \rightarrow M \rightarrow M'' \rightarrow 0$$

is short exact if and only if the induced sequence of \mathbb{k} -vector spaces

$$0 \rightarrow L \otimes_R M' \rightarrow L \otimes_R M \rightarrow L \otimes_R M'' \rightarrow 0$$

is short exact. A similar definition of *faithfully flat* applies to a right module. These notions are of course stronger than *flatness*: a right R -module L is *flat* if $L \otimes_R (-)$ is exact, and similarly for a left module. Free modules are always faithfully flat.

PROPOSITION 2.37. *Let $K \subseteq H$ be a subHopf algebra.*

- (a) *If K is ad-invariant then $HK^+ = K^+H$ and this is a Hopf ideal. Furthermore the quotient mapping $H \rightarrow H/HK^+$ is a homomorphism of Hopf algebras.*
- (b) *If $HK^+ = K^+H$ and H is faithfully flat as a left or right K -module then K is ad-invariant.*
- (c) *If H is finite dimensional then K is ad-invariant if and only if $HK^+ = K^+H$.*

PROOF. Proofs can be found in [Mon93, Rad12].

Since free modules are faithfully flat, part (c) follows from the Nichols-Zoeller Theorem 5.1 that we will meet later. \square

This result leads us to make the following definition.

DEFINITION 2.38. A subHopf algebra $K \subseteq H$ is *normal* if it is ad-invariant and therefore $HK^+ = K^+H$ is a Hopf ideal in H and $H \rightarrow H/HK^+$ is a homomorphism of Hopf algebras. Following Milnor & Moore [MM65] it is common to write

$$H//K = H/HK^+ \cong H \otimes_K \mathbb{k},$$

where the right hand term is defined using the right K -module structure on H and the counit $K \rightarrow \mathbb{k}$ to define the trivial K -module, and this isomorphism is one of left H -modules.

EXAMPLE 2.39. If G is a group then the adjoint actions in $\mathbb{k}G$ are given by $\text{ad}_g^l = g(-)g^{-1}$ and $\text{ad}_g^r = g^{-1}(-)g$ for $g \in G \subseteq \mathbb{k}G$, so $\text{ad}_g^r = \text{ad}_{g^{-1}}^l$. Hence a subalgebra $A \subseteq \mathbb{k}G$ is ad-invariant if and only if for all $g \in G$, $\text{ad}_g^l A = A$.

If $N \triangleleft G$, then $\mathbb{k}N \subseteq \mathbb{k}G$ is a normal subHopf algebra and $\mathbb{k}G//\mathbb{k}N \cong \mathbb{k}G/N$, the group algebra of the quotient group G/N .

EXAMPLE 2.40. Let \mathfrak{g} be a Lie algebra over \mathbb{k} with Lie bracket $[\cdot, \cdot]$ and $U(\mathfrak{g})$ its *universal enveloping algebra* which will be discussed in Chapter 3. This is a cocommutative Hopf algebra in which there is a copy of \mathfrak{g} which generates it as an algebra. Furthermore, each element $x \in \mathfrak{g}$ is primitive and the left adjoint action of x on $y \in \mathfrak{g}$ is given by

$$\text{ad}_x^l(y) = xy - yx = [x, y]$$

so the adjoint action extends the Lie theoretic adjoint action of \mathfrak{g} on itself. In fact the action of ad_x^l on $U(\mathfrak{g})$ is as a *derivation*: for $a, b \in U(\mathfrak{g})$,

$$\text{ad}_x^l(ab) = \text{ad}_x^l(a)b + a\text{ad}_x^l(b),$$

so since every element is a linear combination of monomials in elements of \mathfrak{g} the adjoint action of \mathfrak{g} itself determines it.

If $\mathfrak{h} \subseteq \mathfrak{g}$ is a Lie subalgebra then $U(\mathfrak{h}) \subseteq U(\mathfrak{g})$ is a subHopf algebra. If \mathfrak{h} is *normal* in \mathfrak{g} then $\mathfrak{g}/\mathfrak{h}$ is also a Lie algebra and

$$U(\mathfrak{g})//U(\mathfrak{h}) \cong U(\mathfrak{g}/\mathfrak{h}).$$

By the Poincaré-Birkhoff-Witt Theorem, $U(\mathfrak{g})$ is free as a left or right $U(\mathfrak{h})$ -module.

CHAPTER 3

A cornucopia of examples

3.1. Endomorphism algebras of vector spaces

For a vector space V , its endomorphism algebra is

$$\text{End}_{\mathbb{k}}(V) = \text{hom}(V, V)$$

with composition as its product. If V is finite dimensional then

$$\text{End}_{\mathbb{k}}(V) \cong V \otimes V^*$$

as vector spaces with the obvious pairing

$$(V \otimes V^*) \otimes (V \otimes V^*) \xrightarrow{\cong} V \otimes (V^* \otimes V) \otimes V^* \rightarrow V \otimes \mathbb{k} \otimes V^* \xrightarrow{\cong} V \otimes V^*$$

making this an isomorphism of algebras. Of course if we choose a basis for V and the corresponding dual basis for V^* we can find an isomorphism of algebras with the ring of $\dim_{\mathbb{k}} V$ by $\dim_{\mathbb{k}} V$ matrices

$$\text{End}_{\mathbb{k}}(V) \cong M_{\dim_{\mathbb{k}} V}(\mathbb{k}).$$

3.2. Polynomial algebras and their duals

EXAMPLE 3.1. Let $\mathbb{k}[X]$ be the polynomial ring. We can give it a coproduct by making X primitive,

$$\psi(X) = X \otimes 1 + 1 \otimes X,$$

and the antipode is determined by

$$\chi(X) = -X.$$

This Hopf algebra is bicommutative.

If \mathbb{k} has characteristic 0 this has no ideals which are also coideals, but if the characteristic is $p > 0$ then for $k \geq 1$, (X^{p^k}) is a coideal and $\mathbb{k}[X]/(X^{p^k})$ is a quotient Hopf algebra.

This example can be generalised to a polynomial ring $\mathbb{k}[X_1, \dots, X_n]$ and then there is an isomorphism of Hopf algebras

$$\mathbb{k}[X_1, \dots, X_n] \cong \mathbb{k}[X_1] \otimes \cdots \otimes \mathbb{k}[X_n].$$

EXAMPLE 3.2 (Divided power Hopf algebra). Consider the \mathbb{k} -vector space $\Gamma_{\mathbb{k}}$ with basis consisting of the elements γ_i ($i \geq 0$). Make $\Gamma_{\mathbb{k}}$ into a commutative algebra with product

$$\gamma_i \gamma_j = \binom{i+j}{i} \gamma_{i+j}$$

and unity $1 = \gamma_0$. Make it a cocommutative coalgebra with product

$$\psi(\gamma_k) = \sum_{0 \leq i \leq k} \gamma_i \otimes \gamma_{k-i}$$

and counit

$$\varepsilon(\gamma_0) = 1, \quad \varepsilon(\gamma_k) = 0 \quad (k > 0).$$

Then combining these structures $\Gamma_{\mathbb{k}}$ becomes a bicommutative Hopf algebra with antipode defined recursively using $\chi\gamma_1 = -\gamma_1$ and

$$\sum_{0 \leq i \leq k} \gamma_i \chi \gamma_{k-i} = 0.$$

If the characteristic of \mathbb{k} is 0 then it is easy to show that there is an isomorphism of Hopf algebras $\Gamma_{\mathbb{k}} \cong \mathbb{k}[X]$ under which

$$\gamma_k \leftrightarrow \frac{1}{k!} X^k$$

where X is primitive. In this case $\text{P}\Gamma_{\mathbb{k}} = \mathbb{k}\{\gamma_1\}$ and $\Gamma_{\mathbb{k}}$ is primitively generated.

If the characteristic of \mathbb{k} is $p > 0$ we have relations such as $\gamma_k^p = 0$ when $k > 0$. As an algebra, $\Gamma_{\mathbb{k}}$ is generated by the elements γ_{p^r} with $r \geq 0$. Also $\text{P}\Gamma_{\mathbb{k}} = \mathbb{k}\{\gamma_1\}$, and so $\Gamma_{\mathbb{k}}$ is not primitively generated.

The finite dual $\Gamma_{\mathbb{k}}^\circ$ is familiar: if we define

$$x: \Gamma_{\mathbb{k}} \rightarrow \mathbb{k}; \quad x(\gamma_k) = \begin{cases} 1 & \text{if } k = 1, \\ 0 & \text{otherwise,} \end{cases}$$

then $x \in \Gamma_{\mathbb{k}}^\circ$ and

$$x^n(\gamma_k) = \begin{cases} 1 & \text{if } k = n, \\ 0 & \text{otherwise,} \end{cases}$$

so $x^n \in \Gamma_{\mathbb{k}}^\circ$ for every $n \geq 0$. Furthermore, x is primitive. Then there is an isomorphism of Hopf algebras

$$\mathbb{k}[X] \xrightarrow{\cong} \Gamma_{\mathbb{k}}^\circ; \quad X^k \mapsto x^k.$$

In fact this relationship is symmetric: $\mathbb{k}[X]^\circ \cong \Gamma_{\mathbb{k}}$.

3.3. The free vector space

Let X be a set and recall the free vector space on X , $\mathbb{F}(X)$.

For any non-empty set X , $\mathbb{F}(X \times X) \cong \mathbb{F}(X) \otimes \mathbb{F}(X)$ and the diagonal map $X \rightarrow X \times X$ induces a \mathbb{k} -linear map $\psi: \mathbb{F}(X) \rightarrow \mathbb{F}(X) \otimes \mathbb{F}(X)$ with $\psi(x) = x \otimes x$ for $x \in X$.

$$\begin{array}{ccccc} & & \psi & & \\ & \searrow & \xrightarrow{\quad} & \searrow & \\ \mathbb{F}(X) & \longrightarrow & \mathbb{F}(X \times X) & \longrightarrow & \mathbb{F}(X) \otimes \mathbb{F}(X) \end{array}$$

Since there is a bijection $X \times (X \times X) \cong (X \times X) \times X$ this is coassociative. If we take any set $\mathbf{1}$ with a single element it is a terminal object and there are bijections

$$\mathbf{1} \times X \cong X \cong X \times \mathbf{1}.$$

Also, $\mathbb{F}(\mathbf{1}) \cong \mathbb{k}$. Now the unique function $X \rightarrow \mathbf{1}$ induces a counit $\varepsilon: \mathbb{F}(X) \rightarrow \mathbb{k}$. Putting all this together we find that $(\mathbb{F}(X), \psi, \varepsilon)$ is a coalgebra. In fact the switch map gives a commutative diagram

$$\begin{array}{ccc} & X & \\ \swarrow & & \searrow \\ X \times X & \xleftarrow[\cong]{\tau} & X \times X \end{array}$$

and using this we can show that $(\mathbb{F}(X), \psi, \varepsilon)$ is a cocommutative coalgebra.

If X is a monoid it has a product $X \times X \rightarrow X$ and a unit $\mathbf{1} \rightarrow X$. By functoriality, these induce maps

$$\varphi: \mathbb{F}(X) \otimes \mathbb{F}(X) \xrightarrow{\cong} \mathbb{F}(X \times X) \rightarrow \mathbb{F}(X), \quad \eta: \mathbb{k} \rightarrow \mathbb{F}(X),$$

so that $(\mathbb{F}(X), \varphi, \eta)$ is an algebra which is commutative if and only if the monoid X is commutative.

Now if X is a monoid we can put together the coalgebra and algebra structures to obtain a cocommutative bialgebra $(\mathbb{F}(X), \varphi, \eta, \psi, \varepsilon)$ which is commutative if and only if the monoid is commutative. With this structure, $\mathbb{k}X = \mathbb{F}(X)$ is called the *monoid algebra* of X .

There is also a dual object, namely

$$\mathbb{k}^X = \text{the set of all functions } X \rightarrow \mathbb{k} \text{ constant a.e.,}$$

i.e., functions which are constant except for a finite number of exceptional values. This has a basis consisting of the Dirac functions $\delta_x: X \rightarrow \mathbb{k} (x \in X)$ together with the constant function 1 taking value 1, where

$$(3.1) \quad \delta_x(y) = \begin{cases} 1 & \text{if } y = x, \\ 0 & \text{otherwise.} \end{cases}$$

Notice that the basis elements δ_x are idempotents as are the elements $1 - \delta_x$; in fact when $x \neq y$, $\delta_x \delta_y = 0$ so δ_x and δ_y are *orthogonal idempotents*. For two sets X, Y , there is an isomorphism of algebras

$$\mathbb{k}^{X \times Y} \cong \mathbb{k}^X \otimes \mathbb{k}^Y.$$

The diagonal map $X \rightarrow X \times X$ induces a multiplication

$$\mathbb{k}^X \otimes \mathbb{k}^X \xrightarrow{\cong} \mathbb{k}^{X \times X} \rightarrow \mathbb{k}^X$$

which is ‘pointwise product’ of functions. This makes into a commutative algebra. In fact

$$\mathbb{k}^X \cong \text{hom}(\mathbb{k}X, \mathbb{k}).$$

The Dirac functions satisfy

$$\delta_x \delta_y = \begin{cases} \delta_x & \text{if } x = y, \\ 0 & \text{otherwise.} \end{cases}$$

When X is a finite monoid, there is a coproduct and \mathbb{k}^X is then a commutative bialgebra.

3.4. Group algebras and dual group algebras

If G is a group, the inverse map $G \rightarrow G$ induces a coalgebra map $\chi: \mathbb{F}(G) \rightarrow \mathbb{F}(G)$. Then $(\mathbb{F}(G), \varphi, \eta, \psi, \varepsilon, \chi)$ is a cocommutative Hopf algebra. The algebra $\mathbb{k}G = \mathbb{F}(G)$ is called the *group algebra* of G , and we know that it is also Hopf algebra. The dual \mathbb{k}^G is also a commutative Hopf algebra, the *dual group algebra*; when G is finite \mathbb{k}^G is the \mathbb{k} -linear dual of $\mathbb{k}G$.

Let’s make these structures explicit. For $\mathbb{k}G$ the coproduct, counit and antipode are given on a group elements $g \in G$ by

$$(3.2) \quad \psi(g) = g \otimes g, \quad \varepsilon(g) = 1, \quad \chi(g) = g^{-1}.$$

For \mathbb{k}^G we have

$$(3.3) \quad \psi(\delta_g) = \sum_{h \in G} \delta_h \otimes \delta_{h^{-1}g}, \quad \varepsilon(\delta_g) = \delta_g(1) = \begin{cases} 1 & \text{if } g = 1, \\ 0 & \text{otherwise,} \end{cases} \quad \chi(\delta_g) = \delta_{g^{-1}}.$$

This construction of the (Hopf) algebra $\mathbb{k}G$ for each group defines two left adjoints. Recall that every ring has a group of units and in particular every \mathbb{k} -algebra A has a group of units A^\times ; we can think of this as defining a functor $(-)^\times: \mathbf{Alg}_{\mathbb{k}} \rightarrow \mathbf{Gp}$. Of course every Hopf algebra is also an algebra so there is a restriction to a functor $(-)^\times: \mathbf{HA}_{\mathbb{k}} \rightarrow \mathbf{Gp}$.

PROPOSITION 3.3. *The functor $\mathbb{F}: \mathbf{Gp} \rightarrow \mathbf{Alg}_{\mathbb{k}}$ is a left adjoint to the unit functor, i.e., there is natural isomorphism of bifunctors*

$$\mathbf{Alg}_{\mathbb{k}}(\mathbb{F}(-), -) \cong \mathbf{Gp}(-, (-)^{\times}).$$

Similarly, the functor $\mathbb{F}: \mathbf{Gp} \rightarrow \mathbf{HA}_{\mathbb{k}}$ is a left adjoint to the group-like element functor G , i.e., there is natural isomorphism of bifunctors

$$\mathbf{HA}_{\mathbb{k}}(\mathbb{F}(-), -) \cong \mathbf{Gp}(-, G(-)).$$

For finite groups, we can do something similar with $\mathbb{M}(G)$, this time obtaining a commutative Hopf algebra contravariantly functorial in G . It is common to set $\mathbb{k}^G = \mathbb{M}(G)$ and call this the dual group algebra of G .

3.5. Poset coalgebras and algebras

Let $(\mathcal{P}, \preccurlyeq)$ be a *locally finite poset*, i.e., each interval

$$[x, y] = \{t \in \mathcal{P} : x \preccurlyeq t \preccurlyeq y\}$$

is finite. We define a vector space $C(\mathcal{P}, \preccurlyeq)$ with basis the symbols $[x, y]$ with $x \preccurlyeq y$. Then

$$\psi: C(\mathcal{P}, \preccurlyeq) \rightarrow C(\mathcal{P}, \preccurlyeq) \otimes C(\mathcal{P}, \preccurlyeq); \quad \psi([x, y]) = \sum_{t \in [x, y]} [x, t] \otimes [t, y]$$

is a coproduct and

$$\varepsilon: C \rightarrow \mathbb{k}; \quad \varepsilon([x, y]) = \begin{cases} 1 & \text{if } x = y, \\ 0 & \text{otherwise,} \end{cases}$$

is its counit. There is a dual *incidence algebra* $A(\mathcal{P}, \preccurlyeq)$ which consists of the finitely supported functions $f: \{[x, y] : x \preccurlyeq y\} \rightarrow \mathbb{k}$ with the product given by convolution,

$$(f * g)([x, y]) = \sum_{t \in [x, y]} f([x, t])g([t, y]),$$

and the unit is given by the constant functions.

3.6. Free algebras, bialgebras and Hopf algebras

The forgetful functor $\mathbf{Alg}_{\mathbb{k}} \rightarrow \mathbf{Vect}_{\mathbb{k}}$ which forgets the multiplication has a left adjoint. Its construction involves the tensor powers of a vector space V : set $T^0(V) = \mathbb{k}$ and for each $n \geq 1$,

$$T^n(V) = V \otimes T^{n-1}(V) = V^{\otimes n}.$$

Then

$$T(V) = \bigoplus_{n \geq 0} T^n(V) = \bigoplus_{n \geq 0} V^{\otimes n}.$$

There are obvious linear mappings $T^m(V) \otimes T^n(V) \rightarrow T^{m+n}(V)$ and these make $T(V)$ into a \mathbb{k} -algebra. It is easy to see that for any \mathbb{k} -linear mapping $f: U \rightarrow V$ there is a unique algebra homomorphism $T(f): T(U) \rightarrow T(V)$ which extends $T^1(f) = f: T^1(U) \rightarrow T^1(V)$. Then $T(V)$ is called the *tensor algebra* or the *free algebra* on V .

PROPOSITION 3.4. *The functor $T: \mathbf{Vect}_{\mathbb{k}} \rightarrow \mathbf{Alg}_{\mathbb{k}}$ is left adjoint to the forgetful functor $\mathbf{Alg}_{\mathbb{k}} \rightarrow \mathbf{Vect}_{\mathbb{k}}$, i.e., there is a natural isomorphism of bifunctors*

$$\mathbf{Alg}_{\mathbb{k}}(T(-), (-)) \cong \mathbf{Vect}_{\mathbb{k}}((-), (-)).$$

We can modify this to the case of commutative algebras. The free algebra $T(V)$ has a 2-sided ideal $I(V)$ generated by all the elements of form

$$u \otimes v - v \otimes u \in T^2(V) \quad (u, v \in V).$$

The quotient algebra

$$S(V) = T(V)/I(V)$$

is commutative since we have implicitly killed all commutators (exercise!), and $S(V)$ is called the *symmetric algebra* or the *free commutative algebra* on V .

PROPOSITION 3.5. *The functor $S: \mathbf{Vect}_k \rightarrow \mathbf{Alg}_k^{\text{co}}$ is left adjoint to the forgetful functor $\mathbf{Alg}_k^{\text{co}} \rightarrow \mathbf{Vect}_k$, i.e., there is a natural isomorphism of bifunctors*

$$\mathbf{Alg}_k^{\text{co}}(S(-), (-)) \cong \mathbf{Vect}_k((-), (-)).$$

Notice that both $T(V)$ and $S(V)$ are naturally \mathbb{N} -graded algebras: the degree n part of $T(V)$ is $T^n(V)$ and its image in $S(V)$ is $S^n(V)$. As a vector space,

$$S(V) = \bigoplus_{n \geq 0} S^n(V).$$

3.7. Free bialgebras and free Hopf algebras

There is also a functor which forgets the algebra structure:

$$\mathbf{HA}_k \rightarrow \mathbf{Coalg}_k; \quad (H, \varphi, \eta, \psi, \varepsilon, \chi) \mapsto (H, \psi, \varepsilon, \chi).$$

This also has a left adjoint, but we have to construct it in stages.

We first form the composition

$$\mathbf{Coalg}_k \rightarrow \mathbf{Vect}_k \xrightarrow{T} \mathbf{Bialg}_k$$

into the category of bialgebras, where the first map is the forgetful functor. Then for a coalgebra C , $T(C)$ is the *free bialgebra* on C . Its elements are sums of monomials in elements of $C \cong T^1(C)$ so the coproduct is obtained using

$$\psi(c_1 c_2 \cdots c_\ell) = \psi(c_1) \psi(c_2) \cdots \psi(c_\ell).$$

There is a similar construction forming the free *free commutative bialgebra* on C , $S(C)$.

There are variants of these for (co)augmented coalgebras which form a category k/\mathbf{Coalg}_k (i.e., coalgebras under k). Given a coaugmented coalgebra $\eta: k \rightarrow C$ we form $T(C)$ then pass to the quotient bialgebra

$$T(C)/(\eta(1) - 1).$$

This of course identifies $\eta(1) \in T(C)$ with $1 \in T^0(C)$. We can do a similar thing with the commutative version.

To get the free algebra functor into \mathbf{HA}_k we take the direct sum of coalgebras $C \oplus C^{\text{op}}$, form the free algebra $T(C \oplus C^{\text{op}})$ and then impose relations to identify each element c^{op} with an antipode applied to c , i.e., quotient by the ideal generated by all the expressions

$$\sum c_{(1)} \otimes c_{(2)}^{\text{op}} - \varepsilon(c) \otimes 1, \quad \sum c_{(1)}^{\text{op}} \otimes c_{(2)} - \varepsilon(c) \otimes 1$$

where $c \in C$.

To get a free commutative Hopf algebra we can use S instead of T . In fact

$$S(C \oplus C^{\text{op}}) \cong S(C) \otimes S(C^{\text{op}}).$$

Since a Hopf algebra is naturally a coaugmented coalgebra we can also do this by first applying the free bialgebra functors for coaugmented coalgebras.

3.8. Enveloping algebras of Lie algebras

Recall that a *Lie algebra over \mathbb{k}* is a vector space L equipped with a linear mapping called the *Lie bracket*

$$[-, -]: L \otimes L \rightarrow L$$

which satisfies the following conditions for all $x, y, z \in L$:

$$\begin{aligned} \text{(Jacobi identity)} \quad & [x, [y, z]] + [y, [z, x]] + [z, [x, y]] = 0; \\ \text{(Anticommutativity)} \quad & [y, x] + [x, y] = 0; \\ \text{(Alternativity)} \quad & [x, x] = 0. \end{aligned}$$

If the characteristic of \mathbb{k} is not 2 then anticommutativity implies alternativity so then the last condition is redundant. Care is also required when the characteristic is 3 but we will ignore this subtlety.

A Lie algebra with trivial bracket $[x, y] = 0$ is called an *abelian Lie algebra*; abelian Lie algebras are essentially the same thing as vector spaces.

Lie algebras over \mathbb{k} form an abelian category $\mathbf{Lie}_{\mathbb{k}}$ with homomorphisms preserving brackets.

For any algebra A , its elements form a Lie algebra with the usual commutator $[x, y] = xy - yx$ as its bracket. Of course this Lie algebra is abelian if and only if the algebra is commutative. This construction defines a functor $\mathbf{Alg}_{\mathbb{k}} \rightarrow \mathbf{Lie}_{\mathbb{k}}$. We will see that it has a left adjoint. But in fact there is another functor $P: \mathbf{HA}_{\mathbb{k}} \rightarrow \mathbf{Lie}_{\mathbb{k}}$ which also has a left adjoint.

To construct the adjoint in the algebra case we first recall the free algebra functor T . We can apply this to a Lie algebra L but the linear mapping $L = T^1(L) \hookrightarrow T(L)$ is not a homomorphism of Lie algebras if we make $T(L)$ a Lie algebra using the commutator. To correct this we have to force relations by passing to a quotient algebra. We consider the 2-sided ideal $J(L) \triangleleft T(L)$ generated by all the elements

$$x \otimes y - y \otimes x - [x, y] \quad (x, y \in L).$$

Notice that $x \otimes y, y \otimes x \in T^2(L)$ but $[x, y] \in T^1(L)$. The resulting quotient algebra

$$U(L) = T(L)/J(L)$$

is called the *universal enveloping algebra* of L . It can be verified that the mapping $L \rightarrow U(L)$ is a Lie algebra homomorphism where $U(L)$ is given the commutator as its Lie bracket (it is injective except possibly when the characteristic of \mathbb{k} is 3).

PROPOSITION 3.6. *The functor $U: \mathbf{Lie}_{\mathbb{k}} \rightarrow \mathbf{Alg}_{\mathbb{k}}$ is left adjoint to the functor $\mathbf{Alg}_{\mathbb{k}} \rightarrow \mathbf{Lie}_{\mathbb{k}}$ sending each algebra to its Lie algebra with the commutator bracket, i.e., there is a natural isomorphism of bifunctors*

$$\mathbf{Alg}_{\mathbb{k}}(U(-), (-)) \cong \mathbf{Lie}_{\mathbb{k}}((-), (-)).$$

The Poincaré-Birkhoff-Witt Theorem is an important result which describes the vector space structure of $U(L)$ at least given a certain kind of basis of L . Here is a version when L is of finite or countable dimension with a basis x_1, x_2, \dots and we denote the image of an element $\ell \in L$ in $U(L)$ by $\tilde{\ell}$.

THEOREM 3.7 (Poincaré-Birkhoff-Witt Theorem). *The distinct monomials*

$$\tilde{x}_1^{k_1} \tilde{x}_2^{k_2} \cdots \tilde{x}_{\ell}^{k_{\ell}} \quad (k_i \geq 0)$$

form a basis for $U(L)$. In particular the linear map $L \rightarrow U(L)$ sending x to \tilde{x} is injective.

Since the map $L \rightarrow U(L)$ is injective, it is usual to omit the tildes and write x for the image of $x \in L$ in $U(L)$.

Of course we have chosen a particular ordering here; for example to express x_2x_1 we note that in $U(L)$ we have

$$x_2x_1 = (x_1x_2 - x_2x_1) + x_1x_2 = [x_1, x_2] + x_1x_2$$

where $[x_1, x_2] \in L \subseteq U(L)$ is a linear combination of the x_i .

For any Lie algebra L we can make $U(L)$ into a Hopf algebra by defining $L \subseteq U(L)$ to be contained in $PU(L)$. Then $U(L)$ is generated as an algebra by $PU(L)$. Of course for any Hopf algebra H the inclusion $P(H) \hookrightarrow H$ is a Lie homomorphism so it induces a Hopf algebra homomorphism $UP(H) \rightarrow H$; if this is surjective then H is called *primitively generated*. Primitively generated Hopf algebras are cocommutative and in a sense the ‘easy’ ones to understand.

PROPOSITION 3.8. *The functor $U: \mathbf{Lie}_{\mathbb{k}} \rightarrow \mathbf{HA}_{\mathbb{k}}$ is left adjoint to the functor $P: \mathbf{HA}_{\mathbb{k}} \rightarrow \mathbf{Lie}_{\mathbb{k}}$, i.e., there is a natural isomorphism of bifunctors*

$$\mathbf{HA}_{\mathbb{k}}(U(-), (-)) \cong \mathbf{Lie}_{\mathbb{k}}((-), P(-)).$$

Here are some examples.

EXAMPLE 3.9. Let p be a prime number and \mathbb{k} a field of characteristic p . Let

$$H = \mathbb{k}[X]/(X^p)$$

and write $x = X + (X^p) \in H$. Then the coproduct $\psi(x) = 1 \otimes x + x \otimes 1 + x \otimes x$ and counit $\varepsilon(x) = 0$ make H a bicommutative Hopf algebra.

It is easy to see that $PH = 0$, so H is not primitively generated.

This is a disguised version of the group algebra $\mathbb{k}C_p$. If the characteristic of \mathbb{k} is not equal to p and \mathbb{k} contains a primitive p -th root of unity then $\mathbb{k}C_p$ is not primitively generated.

EXAMPLE 3.10. The polynomial ring $H = \mathbb{k}[X]$ given the coproduct

$$\psi(X^n) = \sum_{0 \leq i \leq n} \binom{n}{i} X^i \otimes X^{n-i}$$

is a commutative and cocommutative Hopf algebra which is primitively generated. If the characteristic of \mathbb{k} is 0 then $PH = \mathbb{k}\{x\}$, but if it is a prime number p then

$$PH = \mathbb{k}\{x^{p^k} : k \geq 0\}.$$

3.9. Restricted Lie algebras

In this section we will assume that \mathbb{k} has positive characteristic p . Then there is a variation of the notion of Lie algebra called a *restricted Lie algebra* or a *p-Lie algebra*. For details see Jacobson [Jac79, section V.7] or Milnor & Moore [MM65, section 6].

For any Hopf algebra H over \mathbb{k} , there is a Frobenius mapping

$$PH \rightarrow PH; \quad x \mapsto x^p.$$

Of course this is not linear over \mathbb{k} but if $t \in \mathbb{k}$, then $(tx)^p = t^p x^p$. If $x, y \in PH$ commute then $(x + y)^p = x^p + y^p$, but in general there is a more complicated formula.

A Lie algebra over \mathbb{k} is called a *restricted Lie algebra* if there is an additive homomorphism $(-)^{[p]}: L \rightarrow L$ (the *restriction*) such that

- for $x \in L$ and $t \in \mathbb{k}$, $(tx)^{[p]} = t^p x^{[p]}$;

- $\text{ad}_{x^{[p]}} = \text{ad}_x^p = \text{ad}_x \circ \text{ad}_x \circ \cdots \circ \text{ad}_x$, where $\text{ad}_x: L \rightarrow L$ is the linear mapping given by $\text{ad}_x(y) = [x, y]$;
- for $x, y \in L$,

$$(x + y)^{[p]} = x^{[p]} + y^{[p]} + \sum_{i=1}^{p-1} s_i(x, y)$$

where for an indeterminate Z , the coefficient of Z^{i-1} in $(\text{ad}_{tx+y})^{p-1}(x)$ is $is_i(x, y)$.

For a Hopf algebra H over \mathbb{k} , its primitives form a restricted Lie algebra with the Frobenius map being its retraction. Then there is a functor from Hopf algebras to restricted Lie algebras and this has a left adjoint given on a restricted Lie algebra by

$$V(L) = U(L)/(\widetilde{x^{[p]} - \tilde{x}^p} : x \in L),$$

a quotient Hopf algebra of the usual enveloping algebra; this is called the *restricted enveloping algebra* of L . When L is finite dimensional, so is $V(L)$ whereas $U(L)$ is infinite dimensional. There is also a version of the PBW Theorem for $V(L)$.

3.10. Affine group schemes

Motivated by Alexander Grothendieck's insights, much of Algebraic Geometry is now centred on *representable functors* on the category of commutative algebras over a base ring. A commutative algebra $R \in \mathbf{Alg}_{\mathbb{k}}^{\text{co}}$ defines a functor

$$\text{Spec}(R): \mathbf{Alg}_{\mathbb{k}}^{\text{co}} \rightarrow \mathbf{Set}; \quad \text{Spec}(R)(A) = \mathbf{Alg}_{\mathbb{k}}^{\text{co}}(R, A).$$

This is called an *affine scheme*. Its space of *geometric points* is given by its value on an algebraic closure $\bar{\mathbb{k}}$, $\text{Spec}(R)(\bar{\mathbb{k}})$.

In practise such a functor often has a factorisation through a functor into a concrete category \mathbf{C} such as the category of groups; in this case we say that it is a *\mathbf{C} -scheme*.

$$\begin{array}{ccc} & \text{Spec}(R) & \rightarrow \mathbf{C} \\ & \curvearrowright & \downarrow \\ \mathbf{Alg}_{\mathbb{k}}^{\text{co}} & \longrightarrow & \mathbf{Set} \end{array}$$

Let's suppose that $\text{Spec}(R)$ takes values in the category of groups \mathbf{Gp} so it is *group scheme*. Now the coproduct in $\mathbf{Alg}_{\mathbb{k}}^{\text{co}}$ is given by \otimes and \mathbb{k} is an initial object, so for any commutative algebra A ,

$$\text{Spec}(R \otimes R)(A) \cong \text{Spec}(R)(A) \times \text{Spec}(R)(A)$$

and $\text{Spec}(\mathbb{k})(A)$ contains only the unit homomorphism $\mathbb{k} \rightarrow A$. The multiplication is a natural transformation

$$\text{Spec}(R \otimes R) \cong \text{Spec}(R) \times \text{Spec}(R) \rightarrow \text{Spec}(R)$$

so if we evaluate on $R \otimes R$ we get

$$\text{Spec}(R \otimes R)(R \otimes R) \rightarrow \text{Spec}(R)(R \otimes R)$$

which sends $\text{Id}_{R \otimes R}$ to a homomorphism $\psi: R \rightarrow R \otimes R$. Similarly the identity evaluated on \mathbb{k} gives

$$\text{Spec}(\mathbb{k})(\mathbb{k}) \rightarrow \text{Spec}(R)(\mathbb{k})$$

which sends $\text{Id}_{\mathbb{k}}$ to an element $\varepsilon: R \rightarrow \mathbb{k}$. Finally the inverse map gives a natural transformation $\text{Spec}(R) \rightarrow \text{Spec}(R)$ which when evaluated on R sends Id_R to $\chi: R \rightarrow \mathbb{k}$. All of these structure maps are algebra homomorphisms by definition and make $(R, \psi, \varepsilon, \chi)$ a cogroup object in $\mathbf{Alg}_{\mathbb{k}}^{\text{co}}$,

in other words we have a commutative Hopf algebra; if the group scheme takes values in abelian groups then it will be cocommutative. Here are some examples.

Each commutative algebra A has a group of units A^\times . To specify a unit means to pick an element and another element which is its inverse. We can do this with the affine scheme $\text{Spec}(\mathbb{k}[U, V]/(UV - 1))$ where

$$\psi(U) = U \otimes U, \quad \psi(V) = V \otimes V, \quad \varepsilon(U) = 1 = \varepsilon(V), \quad \chi(U) = V, \quad \chi(V) = U.$$

It is usual to set $V = U^{-1}$ and write $\mathbb{k}[U, U^{-1}] = \mathbb{k}[U, V]/(UV - 1)$. This is the *multiplicative group scheme* which is often denoted \mathbb{G}_m .

For each natural number $n \geq 1$, there is a natural transformation $[n]: \mathbb{G}_m \rightarrow \mathbb{G}_m$ induced by the Hopf algebra homomorphism $\mathbb{k}[U, U^{-1}] \rightarrow \mathbb{k}[U, U^{-1}]$ which maps U to U^n . This corresponds to the n -th power map when evaluated on an algebra A .

$$\begin{array}{ccc} \mathbb{G}_m(A) & \xrightarrow{[n]} & \mathbb{G}_m(A) \\ \parallel & & \parallel \\ A^\times & \xrightarrow{(-)^n} & A^\times \end{array}$$

In fact $\mathbb{G}_m[n] = \ker[n]$ is also a scheme, given by

$$\mathbb{G}_m[n] = \text{Spec}(\mathbb{k}[U, U^{-1}]/(U^n - 1)),$$

represented by the quotient Hopf algebra $\mathbb{k}[U, U^{-1}]/(U^n - 1) \cong \mathbb{k}[U, U^{-1}]/\mathbb{k}[U^n, U^{-n}]$ where $\mathbb{k}[U^n, U^{-n}] \subseteq \mathbb{k}[U, U^{-1}]$ is the evident subHopf algebra.

The multiplicative group scheme can be generalised to a non-abelian group scheme GL_n for $n \geq 2$. For example, when $n = 2$,

$$\text{GL}_2 = \text{Spec}(\mathbb{k}[A, B, C, D, E]/((AD - BC)E - 1))$$

with coproduct induced by matrix multiplication

$$\begin{aligned} \psi(A) &= A \otimes A + B \otimes C, & \psi(B) &= A \otimes B + B \otimes D, \\ \psi(C) &= C \otimes A + D \otimes C, & \psi(D) &= C \otimes B + D \otimes D, \\ \psi(E) &= E \otimes E. \end{aligned}$$

The antipode is induced by the formula for finding the entries in inverse of a 2 by 2 matrix:

$$\begin{aligned} \chi(A) &= DE, & \chi(B) &= -BE, \\ \chi(C) &= -CE, & \chi(D) &= AE, \\ \chi(E) &= AD - BC. \end{aligned}$$

There is a normal subgroup scheme $\text{SL}_2 \triangleleft \text{GL}_2$ given by

$$\text{SL}_2 = \text{Spec}(\mathbb{k}[A, B, C, D]/((AD - BC) - 1))$$

where $\mathbb{k}[A, B, C, D]/((AD - BC) - 1)$ is a quotient Hopf algebra of $\mathbb{k}[A, B, C, D, E]/((AD - BC)E - 1)$.

Natural transformations of group schemes. A natural transformation $\Theta: \text{Spec}(R) \rightarrow \text{Spec}(S)$ between two group schemes represented by Hopf algebras R and S is determined by its effect on $\text{Id}_R \in \text{Spec}(R)(R)$, i.e., by $\Theta(\text{Id}_R) \in \text{Spec}(S)(R)$ which is an algebra homomorphism $\theta: S \rightarrow R$. In order for Θ to give group homomorphisms it turns out that θ must also be a coalgebra homomorphism, hence it is a Hopf algebra homomorphism. This gives rise to a bijection between natural transformations $\text{Spec}(R) \rightarrow \text{Spec}(S)$ and $\mathbf{HA}_{\mathbb{k}}(S, R)$.

3.11. Combinatorial Hopf algebras

The symmetric function Hopf algebra can be defined over any commutative ring \mathbb{k} . It is bicommutative and

$$\text{Symm}(\mathbb{k}) = \mathbb{k}[e_n : n \geq 1]$$

with coproduct given by

$$\psi(e_n) = \sum_{0 \leq i \leq n} e_i \otimes e_{n-i}$$

where $e_0 = 1$. Its vector space of primitives is spanned by the elements s_n defined by $s_1 = e_1$ and the Newton recursion formula

$$s_n = e_1 s_{n-1} - e_2 s_{n-2} + e_3 s_{n-3} - \cdots + (-1)^{n-2} e_{n-1} s_1 + (-1)^{n-1} n e_n.$$

If the characteristic of \mathbb{k} is zero then

$$\text{Symm}(\mathbb{k}) = \mathbb{k}[s_n : n \geq 1]$$

but if it is a prime $p > 0$ then for any k ,

$$s_{pk} = s_k^p.$$

The e_n are essentially the elementary symmetric functions in infinitely many indeterminates while the s_n are the power sums. The antipode is given by

$$\chi(e_n) = h_n$$

where the h_n are the total symmetric functions. There is another set of polynomial generators that occurs, namely the w_n defined recursively by

$$p_n = \sum_{k|n} k w_k^{n/k}.$$

If the characteristic of \mathbb{k} is $p > 0$ then for each m with $p \nmid m$, there is a subHopf algebra

$$B[m] = \mathbb{k}[w_{mp^r} : r \geq 0] \subseteq \text{Symm}(\mathbb{k})$$

and a Hopf algebra splitting

$$\text{Symm}(\mathbb{k}) = \bigotimes_{p \nmid m} B[m].$$

This is related to Witt vectors and also the Necklace Algebra of Rota and Metropolis [MR83].

3.12. Taft algebras

For $n \geq 2$ and $\zeta \in \mathbb{k}$ a primitive n -th root of unity (so the characteristic of \mathbb{k} does not divide n) there is a *Taft Hopf algebra* $H_{n,\zeta}$. As an algebra,

$$H_{n,\zeta} = \mathbb{k}\langle u, v \rangle / (u^n - 1, v^n, vu - \zeta uv).$$

The coproduct is given by

$$\psi(u) = u \otimes u, \quad \psi(v) = v \otimes u + 1 \otimes v$$

and the antipode by

$$\chi(u) = u^{-1}, \quad \chi(v) = -vu^{-1}.$$

The Hopf algebra $H_{n,\zeta}$ is neither commutative nor cocommutative. In fact

$$\chi^2(u) = u, \quad \chi^2(v) = uvu^{-1} = \zeta^{-1}v.$$

The elements $u^i v^j$ ($0 \leq i, j < n$) form a basis of $H_{n,\zeta}$ so $\dim_{\mathbb{k}} H_{n,\zeta} = n^2$. Also, these basis elements are eigenvectors for the linear mapping χ^2 which can be diagonalised with respect to that basis.

The ideal generated by v is a Hopf ideal and the quotient is

$$H_{n,\zeta}/(v) \cong \mathbb{k}C_n,$$

the group algebra of the cyclic group C_n .

Finally, the dual Hopf algebra $H_{n,\zeta}^*$ and $H_{n,\zeta}$ are isomorphic to as Hopf algebras, so $H_{n,\zeta}$ is self-dual.

3.13. Frobenius algebras

Frobenius algebras are commonly encountered, and we will see later that every finite dimensional Hopf algebra is a Frobenius algebra. The book of Kock [Koc04] is a good general introduction.

DEFINITION 3.11. A finite dimensional \mathbb{k} -algebra A is a *Frobenius algebra* if it has a *Frobenius form* $\lambda \in A^* = \text{hom}(A, \mathbb{k})$ which is non-trivial on every simple left submodule.

A left submodule is of course a left ideal; it is *simple* if it has no non-trivial proper submodules so it is *minimal*. A given Frobenius algebra can have many different Frobenius forms.

A Frobenius form λ has an associated non-degenerate \mathbb{k} -bilinear *Frobenius form*

$$\beta: A \times A \rightarrow \mathbb{k}; \quad \beta(x, y) = \lambda(xy)$$

which satisfies

$$\beta(xy, z) = \beta(x, yz).$$

This can be used to show that λ is non-trivial on every simple right submodule.

The Frobenius form induces two \mathbb{k} -linear mappings

$$A \rightarrow A^*; \quad a \mapsto a \cdot \lambda, \quad a \mapsto \lambda \cdot a$$

where

$$a \cdot \lambda(x) = \lambda(xa), \quad \lambda \cdot a(x) = \lambda(ax).$$

If we make A^* a left or right A -module by premultiplying on the right or the left these become left and right A -module isomorphisms. In particular this means that A is injective as a left or right A -module, i.e., it is *self-injective*. This has lots of implications: for example, A is a *Kasch algebra*, i.e., every simple left or right module is isomorphic to a submodule of A .

As well as the algebra (A, φ, η) structure, λ also gives rise a coalgebra (A, ψ, ε) for which the counit is $\varepsilon = \lambda$. Note that (A, φ, η) and (A, ψ, ε) do not interact appropriately to form a bialgebra, but instead the two structures interact through the *Frobenius condition* which says that the following diagram commutes.

$$(3.4) \quad \begin{array}{ccccc} & & A \otimes A & & \\ \psi \otimes \text{Id} \swarrow & & \downarrow \varphi & & \text{Id} \otimes \psi \searrow \\ A \otimes A \otimes A & & A & & A \otimes A \otimes A \\ \text{Id} \otimes \varphi \searrow & & \downarrow \psi & & \varphi \otimes \text{Id} \swarrow \\ & & A \otimes A & & \end{array}$$

In Sweedler notation this becomes for $x, y \in A$,

$$(3.5) \quad \sum x_{(1)} \otimes x_{(2)} y = \psi(xy) = \sum xy_{(1)} \otimes y_{(2)}.$$

Later we will see that every finite dimensional Hopf algebra is a Frobenius algebra, but the coproduct associated to the Frobenius form is not the same as that of the coalgebra structure of the Hopf algebra.

3.14. Quivers and their path algebras

A *quiver* is a (finite) directed/oriented graph $Q = (Q_0, Q_1)$ where Q_0 is the set vertices and Q_1 is the set of edges/arrows. An arrow a has a source vertex $s(a)$ and a target vertex $t(a)$. An ordered pair of arrows (a, b) is *composable* if $t(a) = s(b)$. A *path in Q of length $\ell \geq 0$* is a sequence $\mathbf{a} = (a_1, a_2, \dots, a_\ell)$ of composable arrows. A path of length 0 is just a vertex and we allow it to be pre/post composed with any arrow with it as target or source.

The *path algebra of Q* is the vector space with paths as its basis, i.e.,

$$\mathcal{P}(Q) = \bigoplus_{\mathbf{a}} \mathbb{k} \mathbf{a},$$

and we make this into algebra by extending

$$\mathbf{a} \otimes \mathbf{b} \mapsto \mathbf{ab}$$

where \mathbf{ab} is the path obtained by splicing the composable paths (\mathbf{a}, \mathbf{b}) . If \mathbf{a} and \mathbf{b} are not composable the product is 0. Two paths of length 0/vertices have product 0 if they are distinct, and \mathbf{a} if they are equal. The unity is


$$1 = \sum_{a \in Q_0} a.$$

In $\mathcal{P}(Q)$ the vertices/paths of length 0 are primitive orthogonal idempotents which sum up to 1.

If we now specify a subset R of paths of length at least 2 then these generate an ideal $\mathcal{I}(R) \triangleleft \mathcal{P}(Q)$ so we may form the quotient algebra $\mathcal{P}(Q)/\mathcal{I}(R)$. The pair (Q, R) is called a *quiver with relations*.

Here are some examples.

- The quiver



has exactly one path of length 1, the path algebra has dimension 3 and basis a, b, p and relations


$$a^2 = a, \quad b^2 = b, \quad ap = p = pb, \quad p^2 = 0.$$

- The quiver



is infinite dimensional and its path algebra is the polynomial algebra $\mathbb{k}[p]$.

- The path algebra of the quiver



is also infinite dimensional and is additively isomorphic to the product

$$a\mathbb{k}[pq] \times b\mathbb{k}[qp] \times p\mathbb{k}[qp] \times q\mathbb{k}[pq].$$

3.15. Graded (co)algebra

We can work with graded vector spaces as discussed in Section 1.3. We will view \mathbb{k} as a graded vector space concentrated in degree 0.

A *graded \mathbb{k} -algebra* A_* is a graded vector space together with a product $\varphi: A_* \otimes A_* \rightarrow A_*$ and a unit $\eta: \mathbb{k} \rightarrow A_*$ (which is really a linear map $\mathbb{k} \rightarrow A_0$) so that the analogues of (2.1) are commutative. Of course φ restricts to give maps $A_m \otimes A_n \rightarrow A_{m+n}$ so this can be pulled apart to give statements about maps between ungraded vector spaces.

A graded algebra A_* is (*graded*) *commutative* if the analogue of (2.2) commutes where we use the switch map with built in signs. This means that on elements $a \in A_m$ and $b \in A_n$, if we set $xy = \varphi(x \otimes y)$ on basic tensors,

$$ab = (-1)^{mn}ba.$$

This leads to the result $a^2 = 0$ whenever a has odd degree and $\text{char } \mathbb{k} \neq 2$.

Similarly we can define a *graded \mathbb{k} -coalgebra* C_* to be a graded vector space with a coproduct $\psi: C_* \rightarrow C_* \otimes C_*$ and counit $\varepsilon: C_* \rightarrow \mathbb{k}$ where the analogues of (2.4) commute. Graded cocommutativity is also defined in the obvious way. Here the coproduct gives rise to maps

$$C_n \rightarrow \bigoplus_{k \in \mathbb{Z}} C_k \otimes C_{n-k}.$$

Bialgebras and Hopf algebras can now be defined in the graded setting. But beware: the commutative diagrams required for the interactions between product and coproduct involve using the switch map and this needs to be interpreted using the Koszul sign convention! There are also signs appearing in the formula for the antipode of a Hopf algebra.

CHAPTER 4

Modules and comodules

Rings have modules, and so do algebras. The dual notion for a coalgebra is that of a comodule. As special cases, bialgebras and Hopf algebras have both!

4.1. Modules over an algebra

Algebras are rings with additional structure, so they have modules; in particular a module over a \mathbb{k} -algebra is automatically a \mathbb{k} -vector space. But in keeping with our viewpoint of working with vector spaces, we will define a module as a vector space with additional structure.

DEFINITION 4.1. Given a \mathbb{k} -algebra (A, φ, η) , a *left A -module* (M, μ) is a \mathbb{k} -vector space M and a \mathbb{k} -linear map $\mu: A \otimes M \rightarrow M$ for which the following diagrams commute.

$$\begin{array}{ccc} A \otimes A \otimes M & \xrightarrow{\text{Id} \otimes \mu} & A \otimes M \\ \varphi \otimes \text{Id} \downarrow & & \downarrow \mu \\ A \otimes M & \xrightarrow{\mu} & M \end{array} \quad \begin{array}{ccc} \mathbb{k} \otimes M & \xrightarrow{\cong} & M \\ \eta \otimes \text{Id} \downarrow & \nearrow \mu & \\ A \otimes M & & \end{array}$$

A similar definition applies to a *right A -module*, but we can also view it as a *left* module over the opposite algebra A^{op} so won't discuss right modules explicitly. The action of the algebra for a right module can be thought of either as a map $A^{\text{op}} \otimes M \rightarrow M$ or as a map $M \otimes A \rightarrow M$.

An *A -module homomorphism* $\theta: (M, \mu) \rightarrow (M', \mu')$ is a \mathbb{k} -linear mapping $\theta: M \rightarrow M'$ that makes the following diagram commute.

$$\begin{array}{ccc} A \otimes M & \xrightarrow{\text{Id} \otimes \theta} & A \otimes M' \\ \mu \downarrow & & \downarrow \mu' \\ M & \xrightarrow{\theta} & M' \end{array}$$

Of course a homomorphism has a kernel, an image and a cokernel, all of which are easily seen to be A -modules. Furthermore, the set of all homomorphisms $M \rightarrow N$ between two A -modules is a subspace $\text{Hom}_A(M, M') \subseteq \text{hom}(M, M')$.

EXAMPLE 4.2. An A -module M is *cyclic* if it is generated by one element, so for some $m_0 \in M$,

$$M = Am_0 = \{am_0 : a \in A\}.$$

Such an element x_0 is called a (*cyclic*) *generator*. The *annihilator* of an element $m \in M$,

$$\text{ann}_A(m) = \{a \in A : am = 0\} \subseteq A$$

is always a left ideal of A . For the cyclic module above there is an isomorphism

$$A / \text{ann}_A(m_0) \xrightarrow{\cong} M; \quad a + \text{ann}_A(m_0) \mapsto am_0.$$

More generally, an A -module M is *finitely generated* if there are elements m_1, \dots, m_k such that

$$M = Am_1 + Am_2 + \dots + Am_k = \{a_1m_1 + \dots + a_km_k : a_i \in A\}.$$

So a finitely generated module is quotient of a finite direct sum of free cyclic modules, i.e., there is an epimorphism

$$A^k = \overbrace{A \oplus A \oplus \cdots \oplus A}^k \twoheadrightarrow M$$

under which each standard basis element $e_i = (0, \dots, 0, 1, 0, \dots, 0)$ maps to m_i .

EXAMPLE 4.3. Recall Example 2.3. A left module M over the enveloping algebra $A^e = A \otimes A^{\text{op}}$ is sometimes called a A - A -bimodule because it is simultaneously a left and a right A -module and the two actions commute, i.e., if $a', a'' \in A$ and $m \in M$, then

$$(a'm)a'' = a'(ma'').$$

An important example of such a module is A itself acted on by A through left and right multiplication. This gives rise to an algebra homomorphism $A^e \rightarrow \text{End}_{\mathbb{k}}(A)$; when A is finite dimensional this homomorphism need not be injective, but if $A^e \cong \text{End}_{\mathbb{k}}(A)$ then A is called an *Azumaya algebra*. Examples include matrix rings of central simple algebras over \mathbb{k} and they give rise to the *Brauer group* of the field which appears in Galois Theory and Class Field Theory.

The multiplication map

$$A^e = A \otimes A^{\text{op}} \rightarrow A; \quad x \otimes y^{\text{op}} \mapsto xy$$

is a surjective homomorphism of A^e -modules. If A is a projective A^e -module (or equivalently if this is a split surjection) then A is called *separable*. For the case where A is a field extension of \mathbb{k} this is equivalent to the notion of separability met in Galois Theory.

For a vector space W , the tensor product $A \otimes W$ becomes a left A -module where the composition

$$A \otimes (A \otimes W) \xrightleftharpoons[\cong]{} (A \otimes A) \otimes W \xrightarrow[\varphi]{} A \otimes W$$

is the multiplication. The left module $A \otimes W$ is called an *extended* A -module.

The set of A -module homomorphisms $\theta: (M, \mu) \rightarrow (M', \mu')$ is a subspace

$$\text{Hom}_A(M, M') \subseteq \text{hom}(M, M').$$

If A is commutative it is also an A -module. When $M' = M$, $\text{End}_A(M) = \text{Hom}_A(M, M)$ is an algebra with composition as its product.

There is an adjunction

$$(4.1) \quad \text{Hom}_A(A \otimes W, M) \cong \text{hom}(W, M)$$

under which $\theta \in \text{Hom}_A(A \otimes W, M)$ corresponds to

$$W \xrightleftharpoons[\cong]{} \mathbb{k} \otimes W \xrightarrow[\eta \otimes \text{Id}]{} A \otimes W \xrightarrow[\theta]{} M$$

and $f \in \text{hom}(W, M)$ corresponds to $A \otimes W \rightarrow M$ given on basic tensors by

$$a \otimes w \mapsto af(w).$$

The vector space $\text{hom}(A, W)$ becomes a left A -module with the multiplication of $a \in A$ and $f \in \text{hom}(A, W)$ given by

$$(af)(x) = f(xa).$$

Notice that if $b \in A$,

$$(a(bf))(x) = (bf)(xa) = f((xa)b) = f(x(ab)) = ((ab)f)(x),$$

so $a(bf) = (ab)f$ as required for this to define a left module structure.

This A -module fits into another important adjunction. For any A -module L , there is an isomorphism

$$(4.2) \quad \text{hom}(L, W) \xrightarrow{\cong} \text{Hom}_A(L, \text{hom}(A, W)); \quad f \mapsto (\ell \mapsto (a \mapsto af(\ell))).$$

The inverse sends $g \in \text{Hom}_A(L, \text{hom}(A, W))$ to the composition

$$L \xrightarrow{g} \text{hom}(A, W) \xrightarrow{\eta^*} \text{hom}(\mathbb{k}, W) \xrightarrow{\cong} W$$

induced by the unit $\eta: \mathbb{k} \rightarrow A$.

Before stating the next lemma, we give some definitions that generalise those for modules over rings. In this we always work with left modules but analogous notions apply to right modules.

DEFINITION 4.4. Let A be a \mathbb{k} -algebra.

- An A -module P is *projective* if given a diagram of solid arrows consisting of homomorphisms of A -modules with exact column

$$\begin{array}{ccc} & & U \\ & \nearrow f' & \downarrow p \\ P & \xrightarrow{f} & V \\ & & \downarrow \\ & & 0 \end{array}$$

there is a homomorphism $f': P \rightarrow U$ making the resulting diagram commute.

- An A -module I is *injective* if given a diagram of solid arrows consisting of homomorphisms of A -modules with exact column

$$\begin{array}{ccc} 0 & & \\ \downarrow & & \\ U & \xrightarrow{g} & I \\ \downarrow i & \nearrow g' & \\ V & & \end{array}$$

there is a homomorphism $g': V \rightarrow I$ making the resulting diagram commute.

LEMMA 4.5. Let A be a \mathbb{k} -algebra.

- For any \mathbb{k} -vector space W , the extended A -module $A \otimes W$ is a free module.
- For any A -module M , let M_0 denote its underlying vector space. Then there is a surjective A -module homomorphism $A \otimes M_0 \rightarrow M$.
- If P is a projective A -module, then there is an isomorphism of A -modules $A \otimes P_0 \cong P \oplus Q$ where Q is another projective module. So P is a summand/retract of a free module.
- For any \mathbb{k} -vector space W , $\text{hom}(A, W)$ is an injective A -module.
- If I is an injective A -module then there is an isomorphism of A -modules $\text{hom}(A, I_0) \cong I \oplus J$ where J is also an injective module. So I is a summand/retract of an injective module of the form $\text{hom}(A, I_0)$.

PROOF. (a) Choose a basis of W and use it to give a basis for the A -module $A \otimes W$.

(b) Use the isomorphism (4.1).

- (c) This is a standard argument: use (b) and projectivity.
(d) Suppose that we have a diagram of A -modules with exact row

$$\begin{array}{ccccc} 0 & \longrightarrow & U & \longrightarrow & V \\ & & \downarrow & & \\ & & \text{hom}(A, W) & & \end{array}$$

Now apply $\text{Hom}_A(-, \text{hom}(A, W))$ to the row to obtain a commutative diagram where we use (4.2) to get the vertical isomorphisms.

$$\begin{array}{ccc} \text{Hom}_A(U, \text{hom}(A, W)) & \longleftarrow & \text{Hom}_A(V, \text{hom}(A, W)) \\ \uparrow \cong & & \uparrow \cong \\ \text{hom}(U, W) & \longleftarrow & \text{hom}(V, W) \end{array}$$

But the original \mathbb{k} -linear map $U \rightarrow V$ is split injection, so the linear map in the bottom row is surjective, hence so is the one in the top. It follows that the original diagram of A -modules can be extended with the dotted arrow to

$$\begin{array}{ccccc} 0 & \longrightarrow & U & \longrightarrow & V \\ & & \downarrow & \nearrow \text{dotted} & \\ & & \text{hom}(A, W) & & \end{array}$$

and so $\text{hom}(A, W)$ is injective.

- (e) This is proved in a similar way to (c) using (4.2). □

Now we can summarise all of this in categorical language.

THEOREM 4.6. *There is an abelian category \mathbf{Mod}_A whose objects are the left A -modules and whose morphisms are given by $\mathbf{Mod}_A(M, N) = \text{Hom}_A(M, N)$. The usual $\oplus = \times$ is the coproduct and product; more generally, in this category arbitrary coproducts and coproducts exist. This category has enough projectives and injectives.*

REMARK 4.7. In general there is no obvious monoidal structure on \mathbf{Mod}_A since although we can form the tensor product $M \otimes_A N$ of a right A -module M and a left A -module N , it is only an A -module when A is commutative. An alternative approach is possible when A is a bialgebra such as a Hopf algebra: in that case we can give \mathbf{Mod}_A a monoidal structure using $\otimes_{\mathbb{k}}$. This will be discussed more later.

EXAMPLE 4.8. When G is a group, a $\mathbb{k}G$ -module is the same thing as a \mathbb{k} -representation or \mathbb{k} -linear representation of G .

EXAMPLE 4.9. A module over the polynomial algebra $\mathbb{k}[X]$ is the same thing as a vector space V together with a given linear endomorphism $f: V \rightarrow V$. Since $\mathbb{k}[X]$ is a Euclidean domain there is a structure theory for such modules which are finite dimensional: every such module is isomorphic to a direct sum of cyclic modules $\mathbb{k}[X]/(p(X))$.

An non-trivial A -module M is called *simple* if it contains no non-zero proper submodules. For example, every finite dimensional module contains a simple submodule.

EXAMPLE 4.10. A finite dimensional algebra A is called *semi-simple* if every finite dimensional A -module M is isomorphic to a direct sum of simple submodules. This notion is studied in *Artin-Wedderburn theory*. For a finite group G , $\mathbb{k}G$ is semi-simple if and only if $\text{char } \mathbb{k} \nmid |G|$.

The algebra A is called *simple* if it contains no non-zero proper two sided ideals. Every semi-simple algebra is isomorphic to a product of simple algebras. Every finite dimensional simple algebra is isomorphic to a matrix algebra $M_n(D)$ over a division algebra D whose centre is \mathbb{k} .

EXAMPLE 4.11. In Example 2.2 we described the infinite dimensional Weyl algebra: it is simple if and only if $\text{char } \mathbb{k} = 0$.

EXAMPLE 4.12. For a Lie algebra \mathfrak{g} , a representation (or \mathfrak{g} -module) means a Lie algebra homomorphism $\mathfrak{g} \rightarrow \text{End}_{\mathbb{k}}(V)$ for a vector space V . This is equivalent to making V into a $U(\mathfrak{g})$ -module where $U(\mathfrak{g})$ is the universal enveloping algebra of \mathfrak{g} . Even when \mathfrak{g} is finite dimensional, $U(\mathfrak{g})$ is not so this leads to consideration of infinite dimensional modules. However in many cases the simple \mathfrak{g} -modules are most conveniently described as cyclic quotients of $U(\mathfrak{g})$.

4.2. Comodules over a coalgebra

Dually, a coalgebra has comodules. Their basic theory is very similar to that of modules over an algebra with ‘arrows reversed’.

DEFINITION 4.13. Given a \mathbb{k} -coalgebra (C, ψ, ε) , a *left comodule* (N, ν) is a \mathbb{k} -vector space N and a \mathbb{k} -linear map $\nu: N \rightarrow C \otimes N$ called the *coaction* or *comultiplication* which makes the following diagrams commute.

$$\begin{array}{ccc} C \otimes C \otimes N & \xleftarrow{\text{Id} \otimes \nu} & C \otimes N \\ \psi \otimes \text{Id} \uparrow & & \uparrow \nu \\ C \otimes N & \xleftarrow{\nu} & N \end{array} \quad \begin{array}{ccc} \mathbb{k} \otimes N & \xleftarrow{\cong} & N \\ \varepsilon \otimes \text{Id} \uparrow & \swarrow \nu & \\ C \otimes N & & \end{array}$$

A *right C -comodule* is the same thing as a left comodule over the opposite coalgebra C^{op} .

Sweedler notation is often used for the coproduct of a comodule, one version is

$$\nu(n) = \sum n_{(1)} \otimes n_{(0)}$$

where $n_{(1)} \in C$ and $n_{(0)} \in N$, so the index (0) is reserved for elements in the comodule.

A *C -comodule homomorphism* $\rho: (N, \nu) \rightarrow (N', \nu')$ is a \mathbb{k} -linear mapping $\theta: N \rightarrow N'$ that makes the following diagram commute.

$$\begin{array}{ccc} C \otimes N & \xrightarrow{\text{Id} \otimes \rho} & C \otimes N' \\ \nu \uparrow & & \uparrow \nu' \\ N & \xrightarrow{\rho} & N' \end{array}$$

It is easy to see that the image and the cokernel of a homomorphism ρ are comodules. To see that kernels exist, let $\rho: N \rightarrow N'$ be a C -comodule homomorphism. As a linear mapping ρ has a kernel and there is an exact sequence of linear mappings

$$0 \longrightarrow \ker \rho \longrightarrow N \xrightarrow{\rho} N'$$

and we can extend this to a commutative diagram of solid arrows

$$\begin{array}{ccccccc} 0 & \longrightarrow & \ker \rho & \xrightarrow{\text{inc}} & N & \xrightarrow{\rho} & N' \\ & & \downarrow \text{dotted} & & \downarrow \nu & & \downarrow \nu' \\ 0 & \longrightarrow & C \otimes \ker \rho & \xrightarrow{\text{Id} \otimes \text{inc}} & C \otimes N & \xrightarrow{\rho} & C \otimes N' \end{array}$$

in which the bottom row is exact because tensoring over a field is an exact functor. Now a diagram chase shows that $\nu \circ \text{inc}$ factors through $C \otimes \ker \rho$ hence we can fill in the dotted arrow and more diagram chasing shows that it is a comultiplication making $\ker \rho$ a comodule and a kernel for ρ .

For a vector space W , the tensor product $C \otimes W$ becomes a left C -comodule where the composition

$$C \otimes W \xrightarrow[\psi]{\quad} (C \otimes C) \otimes W \xleftarrow[\cong]{\quad} C \otimes (C \otimes W)$$

is the comultiplication; $C \otimes W$ is called an *extended C -comodule*. The set of C -comodule homomorphisms $\rho: (N, \nu) \rightarrow (N', \nu')$ is a vector subspace $\text{Cohom}_C(N, N') \subseteq \text{hom}(N, N')$. There is an adjunction isomorphism

$$(4.3) \quad \text{Cohom}_C(N, C \otimes W) \cong \text{hom}(N, W)$$

under which $\rho \in \text{Cohom}_C(N, C \otimes W)$ corresponds to

$$N \xrightarrow[\rho]{\quad} C \otimes W \xrightarrow[\varepsilon \otimes \text{Id}]{\quad} \mathbb{k} \otimes W \xleftarrow[\cong]{\quad} W$$

and $g \in \text{hom}(N, W)$ corresponds to the following composition.

$$N \xrightarrow[\nu]{\quad} C \otimes N \xrightarrow[\text{Id} \otimes g]{\quad} C \otimes W$$

PROPOSITION 4.14. *An extended comodule $C \otimes W$ is an injective comodule. Hence a comodule which is a direct summand of an extended comodule is injective.*

PROOF. Suppose given the following commutative diagram of comodule homomorphisms with an exact row.

$$\begin{array}{ccccc} 0 & \longrightarrow & N & \xrightarrow{\rho} & N' \\ & & \downarrow & & \\ & & C \otimes W & & \end{array}$$

Applying $\text{Cohom}_C(-, C \otimes W) \cong \text{hom}(-, W)$ to the row we get a diagram of vector spaces

$$\begin{array}{ccccc} 0 & \longleftarrow & \text{Cohom}_C(N, C \otimes W) & \xleftarrow{\rho^*} & \text{Cohom}_C(N', C \otimes W) \\ & & \uparrow \cong & & \uparrow \cong \\ 0 & \longleftarrow & \text{hom}(N, W) & \xleftarrow{\rho^*} & \text{hom}(N', W) \end{array}$$

with exact bottom row. Therefore the top row is exact so we can fill in the dotted arrow with a comodule homomorphism.

$$\begin{array}{ccccc} 0 & \longrightarrow & N & \xrightarrow{\rho} & N' \\ & & \downarrow & \swarrow & \\ & & C \otimes W & & \end{array}$$

This shows that $C \otimes W$ is injective and any summand of such a comodule is as well by a standard argument. \square

The next result implies that the abelian category of C -comodules has enough injectives and identifies the injectives.

PROPOSITION 4.15. *For any C -comodule N there is a comodule monomorphism $N \rightarrow I$ where I is an extended comodule and so injective. Hence every injective comodule is a direct summand of an extended comodule.*

PROOF. We can view N as just a vector space and then using the isomorphism (4.3) we obtain $\text{Cohom}_C(N, C \otimes N) \cong \text{hom}(N, N)$ and $\text{Id}_N \in \text{hom}(N, N)$ corresponds to a comodule homomorphism $N \rightarrow C \otimes N$ and the commutative diagram

$$\begin{array}{ccc} N & \xrightarrow{\nu} & C \otimes N \\ & \searrow & \downarrow \varepsilon \otimes \text{Id} \\ & & \mathbb{k} \otimes N \\ & & \uparrow \cong \\ & & N \end{array}$$

shows that it is injective, so N embeds into the extended comodule $I = C \otimes N$ which is an injective comodule.

It follows that every injective comodule J embeds into the extended comodule $C \otimes J$ and by injectivity it must be a direct summand. \square

We can summarise this information in a statement about the category of comodules.

THEOREM 4.16. *For a \mathbb{k} -coalgebra C , its comodules and comodule homomorphisms form an abelian category \mathbf{Comod}_C with enough injectives. This category has \oplus as coproduct and product.*

If C is finite dimensional then \mathbf{Comod}_C also has enough projectives.

In general the comodule category of a coalgebra may not have enough projectives, although in many cases it does. This asymmetry leads to slight differences in their homological algebra compared to that of algebras. The finite dimensional case can be verified using ideas in the discussion that follows, see Proposition 4.23.

Now recall that a coalgebra C has an associated algebra C^* . A left C -comodule has an action $\nu^\dagger: C^* \otimes N \rightarrow N$ defined by

$$\gamma n = \nu^\dagger(\gamma \otimes n) = \sum \gamma(n_{(1)})n_{(0)},$$

where of course $\gamma(n_{(1)}) \in \mathbb{k}$. If $\alpha, \beta \in C^*$,

$$\begin{aligned} \alpha(\beta n) &= \sum \alpha(\beta(n_{(1)})n_{(0)}) \\ &= \sum \beta(n_{(1)})\alpha((n_{(0)})_{(1)})(n_{(0)})_{(0)} \end{aligned}$$

while

$$\begin{aligned} (\alpha\beta)n &= \sum (\alpha\beta)(n_{(1)})n_{(0)} \\ &= \sum \alpha((n_{(1)})_{(0)})\beta((n_{(1)})_{(1)})n_{(0)} = \sum \beta((n_{(1)})_{(1)})\alpha((n_{(1)})_{(0)})n_{(0)} \\ &= \sum \beta(n_{(1)})\alpha((n_{(0)})_{(1)})(n_{(0)})_{(1)}, \end{aligned}$$

so $\alpha(\beta n) = (\alpha\beta)n$. Another argument shows that $\varepsilon^*n = n$. So with this multiplication, N becomes a left C^* -module.

DEFINITION 4.17. Let A be a \mathbb{k} -algebra and M a left A -module. Then M is *locally finite* if every element $m \in M$ is contained in a submodule which is a finite dimensional subspace.

In particular this means that for each $m \in M$, the cyclic submodule $Am \subseteq M$ is a finite dimensional subspace. The locally finite A -modules form a full abelian subcategory $\mathbf{Mod}_A^{\text{l.f.}}$ of the full category \mathbf{Mod}_A of all A -modules.

DEFINITION 4.18. Let C be a \mathbb{k} -coalgebra and N a left C -comodule. Then N is *locally finite* if every element $m \in M$ is contained in a subcomodule which is a finite dimensional subspace.

In fact this notion is redundant!

LEMMA 4.19. *Let C be a coalgebra. Then every C -comodule is locally finite.*

PROOF. Let N be a C -comodule. The idea of the proof is that for $n \in N$, the coproduct

$$\nu(n) = \sum n_{(1)} \otimes n_{(0)}$$

gives rise to a finite dimensional subspace spanned by the elements $n_{(0)} \in N$. Now using coassociativity of ν , this can be shown to be a subcomodule. \square

LEMMA 4.20. *Let C be a coalgebra and C^* its dual algebra. Let N be a left C -comodule which we also view as a left C^* -module. Then N is a locally finite C^* -module.*

PROOF. It is sufficient to show that for $n \in N$, the cyclic submodule $C^*n \subseteq N$ is finite dimensional. Lemma 4.19 tells us that n is contained in a finite dimensional subcomodule $W \subseteq N$ and by definition of the action of C^* on n , $C^*n \subseteq W$. \square

Dualising from an algebra to a coalgebra is more problematic unless the finite dual is used. Details can be found in Montgomery [Mon93] or Radford [Rad12]. We summarise the main results.

LEMMA 4.21. *Let A be an algebra and A° its finite dual coalgebra. Let M be a locally finite left A -module. Then M can be given the structure of a left A° -comodule.*

PROPOSITION 4.22. *There is an isomorphism of abelian categories*

$$\mathbf{Mod}_A^{\text{l.f.}} \xrightarrow{\sim} \mathbf{Comod}_{A^\circ}.$$

Of course when A is finite dimensional, $A^\circ = A^*$, and locally finite is equivalent to every element being in a finitely generated submodule. If we restrict attention to finite dimensional modules and comodules we obtain an important related result.

PROPOSITION 4.23. *There is an isomorphism of abelian categories*

$$\mathbf{Mod}_A^{\text{f.d.}} \xrightarrow{\sim} \mathbf{Comod}_{A^*}^{\text{f.d.}}.$$

In particular, projective/injective modules correspond to projective/injective comodules.

Of course the finite dimensional projective A -modules are summands of direct sums of copies of A . Also A is an A^* -comodule through the adjunction

$$\text{hom}(A \otimes A, A) \cong \text{hom}(A, A^* \otimes A).$$

under which the product correspond to a coaction $A \rightarrow A^* \otimes A$ making it an A^* -comodule, and in fact it is projective.

EXAMPLE 4.24. Recall Section 3.4. Let G be a finite group. Then the dual group algebra $\mathbb{k}^G = (\mathbb{k}G)^*$ is a coalgebra with coproduct given in (3.3). Now a $\mathbb{k}G$ -module M gives rise to a \mathbb{k}^G -comodule $\rho: M \rightarrow \mathbb{k}^G \otimes M$ where

$$\rho(m) = \sum_{g \in G} \delta_g \otimes g^{-1}m.$$

4.3. Tensor and cotensor products

Suppose that A is an algebra, (M, μ) is a right A -module and (N, ν) is a left A -module. One definition of the *tensor product* $M \otimes_A N$ makes it the cokernel of the \mathbb{k} -linear mapping $(\mu \otimes \text{Id} - \text{Id} \otimes \nu): M \otimes A \otimes N \rightarrow M \otimes N$. In other words there is an exact sequence

$$(4.4) \quad M \otimes A \otimes N \xrightarrow{\mu \otimes \text{Id} - \text{Id} \otimes \nu} M \otimes N \longrightarrow M \otimes_A N \longrightarrow 0$$

of \mathbb{k} -linear mappings. This can also be defined as the *coequalizer* of the maps $\mu \otimes \text{Id}$ and $\text{Id} \otimes \nu$. Unless A is commutative $M \otimes_A N$ is not an A -module in an obvious way. For this construction there are formulae such as

$$A \otimes_A N \cong N, \quad M \otimes_A A \cong M.$$

Notice also that if $B \subseteq A$ is a subalgebra we can also define $M \otimes_B N$ and there is a linear surjection $M \otimes_B N \rightarrow M \otimes_A N$. If we fix M or N then $M \otimes_A (-)$ and $(-) \otimes_A N$ define additive functors.

PROPOSITION 4.25. *For a fixed right A -module M , the functor $M \otimes_A (-): \mathbf{Mod}_A \rightarrow \mathbf{Vect}_{\mathbb{k}}$ is right exact, i.e., it sends every short exact sequence of left A -modules*

$$0 \rightarrow N' \rightarrow N \rightarrow N'' \rightarrow 0$$

to an exact sequence

$$M \otimes_A N' \rightarrow M \otimes_A N \rightarrow M \otimes_A N'' \rightarrow 0.$$

The left derived functors of $M \otimes_A (-)$ are denoted $\text{Tor}_^A(M, -)$; these can be computed using projective resolutions.*

REMARK 4.26. A left A -module P is called *flat* if for every right A -module M , $\text{Tor}_s^A(M, P) = 0$ for $s > 0$. In fact we can calculate $\text{Tor}_*^A(M, N)$ by using any *flat resolution* $P_{\bullet} \rightarrow N \rightarrow 0$, i.e., a resolution consisting of flat modules P_s . Then $\text{Tor}_*^A(M, N)$ is the homology of the chain complex $N \otimes_A P_{\bullet}$. Free and projective modules are flat, and so are colimits of flat modules.

Now we dualise to comodules. Suppose that C is a coalgebra, (M, μ) is a right C -comodule and (N, ν) is a left C -comodule. We define the cotensor product $M \square_C N$ as the kernel of $(\mu \otimes \text{Id} - \text{Id} \otimes \nu): M \otimes N \rightarrow M \otimes C \otimes N$, so there is an exact sequence

$$(4.5) \quad 0 \longrightarrow M \square_C N \longrightarrow M \otimes N \xrightarrow{\mu \otimes \text{Id} - \text{Id} \otimes \nu} M \otimes C \otimes N$$

and $M \square_C N$ can be viewed as the *equalizer* of $\mu \otimes \text{Id}$ and $\text{Id} \otimes \nu$. Here $M \square_C N$ is only a C -comodule if C is cocommutative. We have

$$C \square_C N \cong N, \quad M \square_C C \cong M.$$

A surjection of coalgebras $C \rightarrow D$ induces an injective linear mapping $M \square_C N \rightarrow M \square_D N$.

We identified injective comodules in Proposition 4.15; they are direct summands of extended comodules. For such a comodule we have

$$M \square_C (C \otimes W) \cong M \otimes W$$

PROPOSITION 4.27. *For a fixed right C -comodule M , the functor $M \square_C (-): \mathbf{Comod}_C \rightarrow \mathbf{Vect}_{\mathbb{k}}$ is left exact, i.e., it sends every short exact sequence of left C -comodules*

$$0 \rightarrow N' \rightarrow N \rightarrow N'' \rightarrow 0$$

to an exact sequence

$$0 \rightarrow M \square_C N' \rightarrow M \square_C N \rightarrow M \square_C N''.$$

The right derived functors of $M \square_C (-)$ are denoted $\text{Cotor}_C^*(M, -)$; these can be computed using injective resolutions.

PROOF. This is a routine exercise in homological algebra. The main thing to note is that a short exact sequence

$$0 \rightarrow I \rightarrow N \rightarrow N'' \rightarrow 0$$

with I an injective comodule splits and so

$$0 \rightarrow M \square_C I \rightarrow M \square_C N \rightarrow M \square_C N'' \rightarrow 0$$

is short exact. □

4.4. Modules over a Hopf algebra

For a Hopf algebra we have both modules and comodules. We will focus on (left) modules but similar things apply to comodules.

From now on, let $(H, \varphi, \eta, \psi, \varepsilon, \chi)$ be a Hopf algebra which we will assume has an invertible antipode; this condition holds if the Hopf algebra is classical since then $\chi \circ \chi = \text{Id}$. We will often indicate the multiplication in a module (M, μ) by writing $hx = \mu(h \otimes x)$.

There are two obvious left modules. First we can let H act on itself by left multiplication, so for μ we just take φ . This is sometimes called the *left regular representation* of H ; this module is free of rank 1. At the other extreme we can let H act on \mathbb{k} using the counit ε , so μ is the map

$$H \otimes \mathbb{k} \rightarrow \mathbb{k}; \quad h \otimes 1 \mapsto \varepsilon(h).$$

In fact for any vector space W we can let H act on W by

$$H \otimes W \rightarrow W; \quad h \otimes w \mapsto \varepsilon(h)w.$$

Such representations are called *trivial representations*, and the one with $W = \mathbb{k}$ is often called *the* trivial representation and it is *simple* or *irreducible*.

Now we come to an important property of the category of modules over a Hopf algebra: it forms a closed monoidal category. Let (M_1, μ_1) and (M_2, μ_2) be two left H -modules. Their tensor product $M_1 \otimes M_2$ is a \mathbb{k} -vector space which also admits a multiplication $\tilde{\mu}$ which is defined to make the diagram

$$\begin{array}{ccc} H \otimes (M_1 \otimes M_2) & \xrightarrow{\tilde{\mu}} & M_1 \otimes M_2 \\ \psi \otimes \text{Id} \otimes \text{Id} \downarrow & & \uparrow \mu_1 \otimes \mu_2 \\ (H \otimes H) \otimes (M_1 \otimes M_2) & \xrightarrow[\cong]{\text{Id} \otimes \text{T} \otimes \text{Id}} & (H \otimes M_1) \otimes (H \otimes M_2) \end{array}$$

commute and making it an H -module $(M_1 \otimes M_2, \tilde{\mu})$. Using Sweedler notation we can write this explicitly as

$$\tilde{\mu}(h \otimes m_1 \otimes m_2) = \sum h_{(1)} m_1 \otimes h_{(2)} m_2.$$

If H is cocommutative then the switch map

$$M_1 \otimes M_2 \xrightarrow[\cong]{\text{T}} M_2 \otimes M_1.$$

is an isomorphism of H -modules, but when H is *not* cocommutative this need not be true.

If W is any vector space with the trivial H -module structure, there is an isomorphism of H -modules

$$W \otimes M \cong M \otimes W.$$

In particular,

$$\mathbb{k} \otimes M \cong M \cong M \otimes \mathbb{k}.$$

For any H -module M we can consider the subspace of H -invariants

$$M^H = \{x \in M : \forall h \in H, hx = \varepsilon(h)x\} \subseteq M.$$

What about $M^* = \text{hom}(M, \mathbb{k})$? There is a natural *right* H -module structure on this given by taking for $h \in H$ and $f \in M^*$,

$$(f \cdot h)(x) = f(hx).$$

We can twist this into a *left* action by defining

$$(h \cdot f)(x) = f(\chi(h)x).$$

More generally, for two H -modules M, N , $\text{hom}(M, N)$ becomes a module with the action given in Sweedler notation by

$$(h \cdot g)(x) = \sum h_{(1)}g(\chi(h_{(2)})x).$$

By an interesting calculation, the subspace of H -invariants of $\text{hom}(M, N)$ turns out to be

$$(4.6) \quad \text{hom}(M, N)^H = \text{Hom}_H(M, N).$$

In particular,

$$M^H \cong \{f(1) \in M : f \in \text{hom}(\mathbb{k}, M)^H\} \cong \text{Hom}_H(\mathbb{k}, M).$$

In fact taking invariants gives a functor $(-)^H : \mathbf{Mod}_H \rightarrow \mathbf{Vect}_{\mathbb{k}}$ which is *left exact*, i.e., it sends every short exact sequence

$$0 \rightarrow L \rightarrow M \rightarrow N \rightarrow 0$$

to an exact sequence

$$0 \rightarrow L^H \rightarrow M^H \rightarrow N^H.$$

This means it has right derived functors denoted by $\text{Ext}_H^*(\mathbb{k}, -)$ and also called the *cohomology of H with coefficients in M* . When $H = \mathbb{k}G$ is a group algebra this is the *cohomology of G relative to \mathbb{k}* .

We can also define the H -coinvariants of an H -module M to be

$$M_H = M / \text{span}\{hm - \varepsilon(h)m : h \in H, m \in M\},$$

where $\text{span}(X)$ means the subspace spanned by X . This can be shown to be isomorphic to the tensor product $\mathbb{k} \otimes_H M$ where we view \mathbb{k} as a right H -module. Taking coinvariants gives a functor $(-)_H : \mathbf{Mod}_H \rightarrow \mathbf{Vect}_{\mathbb{k}}$ which is *right exact*, i.e., it sends every short exact sequence

$$0 \rightarrow L \rightarrow M \rightarrow N \rightarrow 0$$

to an exact sequence

$$L_H \rightarrow M_H \rightarrow N_H \rightarrow 0.$$

The left derived functors are $\text{Tor}_*^H(\mathbb{k}, -)$ and $\text{Tor}_*^H(\mathbb{k}, M)$ is also known as the *homology of H with coefficients in M* . When $H = \mathbb{k}G$ for a group G , this is the homology of G .

When M and N are two left H -modules,

$$(M \otimes N)_H \cong \mathbb{k} \otimes_H (M \otimes N)$$

is also isomorphic to the quotient $(M \otimes N)/T$ where T is the subspace spanned by the elements

$$hm \otimes n - m \otimes hn \quad (h \in H, m \in M, n \in N).$$

As a special case of this, suppose that L is a right H -module; we can make this into a left H -module by defining the action to be

$$h \cdot \ell = \ell \chi(h).$$

Then with this left H -module L and a left H -module N ,

$$(L \otimes N)_H \cong L \otimes_H N$$

where the latter is the right-left tensor product over H .

We can assemble all of these ideas into an important categorical result which we will make use of later.

THEOREM 4.28. *The category of left H -modules \mathbf{Mod}_H under \otimes and $\text{hom}(-, -)$ is closed monoidal. So for H -modules L, M, N there is a functorial adjunction isomorphism*

$$(4.7) \quad \mathbf{Mod}_H(L \otimes M, N) \xleftarrow{\cong} \mathbf{Mod}_H(L, \text{hom}(M, N)).$$

If H is cocommutative \mathbf{Mod}_H is symmetric monoidal.

If M is finite dimensional then (4.7) gives rise to a functorial isomorphism

$$(4.8) \quad \mathbf{Mod}_H(L \otimes M, N) \xleftarrow{\cong} \mathbf{Mod}_H(L, N \otimes M^*).$$

We will return to the issue of the lack of symmetry for non-cocommutative Hopf algebras when we discuss *quantum groups*. We mention one general observation that shows care is need in such situations.

Suppose that M is a finite dimensional H -module. Then the dual space $M^* = \text{hom}(M, \mathbb{k})$ and hence the double dual space $M^{**} = (M^*)^* = \text{hom}(M^*, \mathbb{k})$ admit left H -modules structures as described above.

LEMMA 4.29. *The canonical linear isomorphism $M \rightarrow M^{**}$ need not be an isomorphism of H -modules, but does induce an isomorphism of H -modules*

$$(\chi^2)^* M \xrightarrow{\cong} M^{**},$$

where $(\chi^2)^* M$ is the vector space M given the H -module structure with

$$h \cdot m = \chi^2(h)m.$$

Of course if H is involutory (i.e., $\chi^2 = \text{Id}_H$) then $(\chi^2)^* M = M$, but in general these need not even be isomorphic H -modules.

If W is a vector space which we view as a trivial H -module, then the extended H -module on W is the left H -module $H \otimes W$ with action on basic tensors

$$h(k \otimes w) = (hk) \otimes w.$$

More generally, if $K \subseteq H$ is a subalgebra then for a left K -module N there is an *induced H -module*

$$\text{ind}_K^H N = H \otimes_K N$$

where the tensor product is formed using right K -module structure on H . There is also the *coinduced H -module*

$$\text{coind}_K^H N = \text{Hom}_K(H, N)$$

where the left H -multiplication is induced by *right* multiplication on the domain, i.e.,

$$h \cdot f(-) = f((-)h).$$

These constructions give rise to additive functors

$$\mathrm{ind}_K^H : \mathbf{Mod}_K \rightarrow \mathbf{Mod}_H, \quad \mathrm{coind}_K^H : \mathbf{Mod}_K \rightarrow \mathbf{Mod}_H$$

which have adjoints.

If M is a left H -module, by restriction to K it is also a left K -module which we will denote by $\mathrm{res}_K^H M$; this defines an additive functor

$$\mathrm{res}_K^H : \mathbf{Mod}_H \rightarrow \mathbf{Mod}_K.$$

If M is a left H -module and N is a left K -module, there are isomorphisms

$$(4.9) \quad \mathbf{Mod}_H(\mathrm{ind}_K^H N, M) \cong \mathbf{Mod}_K(N, \mathrm{res}_K^H M),$$

$$(4.10) \quad \mathbf{Mod}_H(M, \mathrm{coind}_K^H N) \cong \mathbf{Mod}_K(\mathrm{res}_K^H M, N),$$

functorial in M, N . Of course this says that there are adjoint pairs

$$\mathrm{ind}_K^H \dashv \mathrm{res}_K^H, \quad \mathrm{res}_K^H \dashv \mathrm{coind}_K^H.$$

Notice that the functor res_K^H both a left and a right adjoint.

When $H = \mathbb{k}G$ and $K = \mathbb{k}H$ for a subgroup $H \leq G$, these adjunctions are the source of *Frobenius reciprocity* in the representation theory of finite groups.

If M is an H -module it is useful to forget its module structure and take its underlying vector space with the trivial H -module structure which we will denote ${}_\epsilon M = \mathrm{res}_{\mathbb{k}}^H M$. The next result is really important and useful when doing homological algebra over a Hopf algebra.

PROPOSITION 4.30. *Suppose that H is a Hopf algebra. For a left H -module M there are isomorphisms of left H -modules*

$$H \otimes M \xleftarrow{\cong} H \otimes {}_\epsilon M \xleftarrow{\cong} M \otimes H,$$

where $H \otimes {}_\epsilon M$ is the extended module for the vector space M . Hence $H \otimes M$ is always a free H -module.

PROOF. The following \mathbb{k} -linear maps are inverse H -module maps:

$$H \otimes M \rightarrow H \otimes {}_\epsilon M; \quad h \otimes x \mapsto \sum h_{(1)} \otimes \chi(h_{(2)})x,$$

$$H \otimes {}_\epsilon M \rightarrow H \otimes M; \quad h \otimes x \mapsto \sum h_{(1)} \otimes h_{(2)}x.$$

In the first map, $\chi(h_{(2)})x$ really does mean we multiply using the original module structure of M not the trivial one! Verifying it is an H -module homomorphism is an exercise in using Sweedler notation (or better, working with commutative diagrams).

A similar argument works for $M \otimes H$. □

If H is finite dimensional it is also true that for any H -module M , there is an isomorphism of H -modules

$$\mathrm{hom}(H, M) \cong H^* \otimes {}_\epsilon M$$

where $H^* = \mathrm{hom}(H, \mathbb{k})$ is injective; later we will see that $H^* \cong H$ so $\mathrm{hom}(H, M)$ is also a free H -module.

An important example: representations of a finite group. A representation of a finite group G over \mathbb{k} is equivalent to a $\mathbb{k}G$ -module. For a vector space V , the induced $\mathbb{k}G$ -module

$$V \uparrow_1^G = \text{ind}_1^G V \cong \mathbb{k}G \otimes V$$

is free and for a $\mathbb{k}G$ -module M it is well known that

$$\mathbb{k}G \otimes M \cong M \uparrow_1^G.$$

If M, N are two $\mathbb{k}G$ -modules then so is $M \otimes N$ with $g \in G$ acting on basic tensors by

$$g \cdot (m \otimes n) = gm \otimes gn.$$

Similarly, $M^* = \text{hom}(M, \mathbb{k})$ is a $\mathbb{k}G$ -module with action of $g \in G$ on $f \in M^*$ given by

$$(g \cdot f)(m) = f(g^{-1}m) \quad (m \in M).$$

This is sometimes called the *dual* or *contragredient* module of M .

If $H \leq G$ then $\mathbb{k}H \subseteq \mathbb{k}G$ is a subHopf algebra and for any $\mathbb{k}H$ -module L , there is an induced module $L \uparrow_H^G$; in particular,

$$\mathbb{k}G/H \cong \mathbb{k}G \otimes_{\mathbb{k}H} \mathbb{k}.$$

If M is a $\mathbb{k}G$ -module we can view it as a $\mathbb{k}H$ -module and then as $\mathbb{k}G$ -modules,

$$\mathbb{k}G \otimes_{\mathbb{k}H} M \cong \mathbb{k}G/H \otimes M.$$

In fact the only subHopf algebras of $\mathbb{k}G$ are the $\mathbb{k}H$. If $N \triangleleft G$ the $\mathbb{k}G/N$ is a quotient Hopf algebra of $\mathbb{k}G$.

In the representation theory of a finite group it is well known that the tensor product of two G -modules M and N is a G -module $M \otimes N$ with the action of $g \in G$ on basic tensors given by

$$g(x \otimes y) = gx \otimes gy;$$

this of course is equivalent to the Hopf algebra definition since in $\mathbb{k}G$ the coproduct on an element $g \in G \subseteq \mathbb{k}G$ is given by $\psi(g) = g \otimes g$.

All of the above structure is important in studying representations and it can be viewed as having its origins in the Hopf algebra structure of $\mathbb{k}G$. Actually it can be thought of in terms of group actions: Suppose that G acts on a set X ; for a subgroup $H \leq G$ there is also a transitive action on G/H . Then there is a diagonal action of G on $G/H \times X$. Also, using the right H -action on G and the restricted H -action on X we can form $G \times_H X$ which has a left G -action. Then there is a G -equivariant bijection

$$G/H \times X \xrightarrow{\cong} G \times_H X.$$

If we apply the free vector space functor to this isomorphism we obtain an isomorphism like the one above.

Representations of Lie algebras. Another important kind of example is provided by Lie algebras.

A *representation* of a Lie algebra L is a Lie algebra homomorphism $\rho: L \rightarrow \text{End}_{\mathbb{k}}(V)$ where V is a vector space and $\text{End}_{\mathbb{k}}(V)$ is given its algebra commutator bracket. Using Proposition 3.6 we see that this data is equivalent to considering the corresponding algebra homomorphism $\tilde{\rho}: U(L) \rightarrow \text{End}_{\mathbb{k}}(V)$ which makes V a $U(L)$ -module.

Since $U(L)$ is a cocommutative Hopf algebra, given two L -representations U, V , their tensor product $U \otimes V$ becomes a $U(L)$ -module, where for $\ell \in L \subseteq U(L)$ and $u \in U, v \in V$.

$$\ell(u \otimes v) = (\ell u) \otimes v + u \otimes (\ell v).$$

Similarly, the dual V^* has a natural $U(L)$ -module structure satisfying

$$\ell f = -f(\ell(-)).$$

If $K \subseteq L$ is a subLie algebra then $U(K) \subseteq U(L)$ is a subHopf algebra so we can define induction coinduction and restriction functors $\text{ind}_{U(K)}^{U(L)}$, $\text{coind}_{U(K)}^{U(L)}$ and $\text{res}_{U(K)}^{U(L)}$. All of this is familiar structure in Lie theory. When L is a *semisimple* Lie algebra we can induce up 1-dimensional representations of *Borel subalgebras* to get irreducible representations.

4.5. Hopf module algebras and coalgebras

A Hopf algebra $(H, \varphi, \eta, \psi, \varepsilon)$ can act or coact on other things such as algebras and coalgebras.

DEFINITION 4.31. An H -module algebra is a \mathbb{k} -algebra (A, φ_A, η_A) which is an H -module with multiplication denoted by $h \cdot a$ for $h \in H$ and $a \in A$, which satisfies

$$h \cdot (ab) = \sum (h_{(1)} \cdot a)(h_{(2)} \cdot b), \quad h \cdot 1 = \varepsilon(h) \quad (h \in H, a, b \in A).$$

An H -module coalgebra is a \mathbb{k} -coalgebra $(C, \psi_C, \varepsilon_C)$ which is an H -module with multiplication denoted by $h \cdot a$ for $h \in H$ and $a \in A$, which satisfies

$$\psi_C(h \cdot c) = \sum (h_{(1)} \cdot c_{(1)}) \otimes (h_{(2)} \cdot c_{(2)}), \quad \varepsilon_C(h \cdot c) = \varepsilon(h)\varepsilon_C(c), \quad (h \in H, c \in C).$$

An H -module bialgebra/Hopf algebra is a bialgebra/Hopf algebra that is both an H -module algebra and a H -module coalgebra.

EXAMPLE 4.32. An important example is provided by the left adjoint action of H on itself: for $h, x \in H$,

$$h \cdot x = \text{ad}_h^l(x) = \sum h_{(1)}x\chi(h_{(2)}).$$

This makes H into an H -module Hopf algebra. To see that the product formula holds, let $a, b, h \in H$. Using a modified version of Sweedler notation where $h_{(ij)} = (h_{(i)})_{(j)}$, we have

$$\begin{aligned} h \cdot (ab) &= \sum h_{(1)}ab\chi(h_{(2)}) \\ &= \sum h_{(11)}(\varepsilon(h_{(12)})1)ab\chi(h_{(2)}) \\ &= \sum h_{(11)}a(\varepsilon(h_{(12)})1)b\chi(h_{(2)}) \\ &= \sum h_{(11)}a\chi(h_{(121)})h_{(122)}b\chi(h_{(2)}) \\ &= \sum h_{(11)}a\chi(h_{(12)})h_{(21)}b\chi(h_{(22)}) \\ &= \sum (h_{(1)} \cdot a)(h_{(2)} \cdot b), \end{aligned}$$

where we have used coassociativity to rewrite the penultimate sum. Verifying the formula $h \cdot 1 = \varepsilon(h)1$ requires a simpler calculation.

If H is commutative then

$$h \cdot x = \sum h_{(1)}\chi(h_{(2)})x = \varepsilon(h)x$$

so in this case the action is trivial.

If $A \subseteq H$ is a subalgebra which is closed under the left adjoint action (i.e., for all $h \in H$, $\text{ad}_h^l A \subseteq A$) then the adjoint action restricted to A makes it into an H -module subalgebra.

EXAMPLE 4.33. Let $\mathbb{k}G$ be the group algebra of a group G and $N \triangleleft G$. Then for $g \in G \subseteq \mathbb{k}G$ and $n \in N \subseteq \mathbb{k}N \subseteq \mathbb{k}G$, the adjoint action is given by

$$g \cdot n = gng^{-1}$$

so $\mathbb{k}N$ is a $\mathbb{k}G$ -module Hopf algebra. This case is very important in representation theory and cohomology of finite groups.

EXAMPLE 4.34. Suppose that K/\mathbb{k} is a Galois extension with Galois group $G = \text{Gal}(K/\mathbb{k})$. Then we can view K as a \mathbb{k} -algebra and each $g \in G$ acts on K as an algebra automorphism. The group algebra $\mathbb{k}G$ also acts on K making it a $\mathbb{k}G$ -module algebra since for $g \in G$ and $x, y \in K$,

$$g \cdot (xy) = (g \cdot x)(g \cdot y).$$

EXAMPLE 4.35 (Cross product algebras). Given a Hopf algebra H and an H -module algebra A , we can form a new algebra $A \sharp H$ called the *cross product algebra* of A and H as follows. The underlying vector space is $A \otimes H$ and we multiply basic tensors using the rule

$$(a' \otimes h')(a'' \otimes h'') = \sum (a'(h'_{(1)} \cdot a'')) \otimes h'_{(2)}h'' = \sum (a'\rho(h'_{(1)} \otimes a'')) \otimes h'_{(2)}h'',$$

where $\rho: H \otimes A \rightarrow A$ is the action map. So the product is the composition shown in the following commutative diagram.

$$\begin{array}{ccc}
 & & A \otimes H \otimes H \otimes A \otimes H \\
 & \nearrow \text{Id} \otimes \psi \otimes \text{Id} \otimes \text{Id} & \uparrow \text{Id} \otimes \text{Id} \otimes \tau \otimes \text{Id} \\
 (A \otimes H) \otimes (A \otimes H) & = A \otimes H \otimes A \otimes H & A \otimes H \otimes A \otimes H \otimes H \\
 & \searrow & \downarrow \text{Id} \otimes \rho \otimes \text{Id} \otimes \text{Id} \\
 & & A \otimes A \otimes H \otimes H \\
 & & \downarrow \varphi_A \otimes \text{Id} \otimes \varphi \\
 & & A \otimes H
 \end{array}$$

CHAPTER 5

Finite dimensional Hopf algebras

In this chapter we survey some of the main results in the theory of finite dimensional Hopf algebras, some of which are recognisable as generalisations of the special case of group algebras of finite groups. The proofs tend to be quite technical so we will sometimes refer to sources such as Radford [Rad12] for details.

5.1. The Nichols-Zoeller Theorem

Here is an important result about finite dimensional Hopf algebras. Earlier versions of this for arbitrary graded connected Hopf algebras were due to Milnor & Moore [MM65]. More general results are known, for example when K is finite dimensional.

THEOREM 5.1 (Nichols & Zoeller). *Let H be a finite dimensional Hopf algebra and let K be a subHopf algebra. Then when viewed as a left or right K -module, H is free. Hence*

$$\dim_{\mathbb{k}} H = (\dim_{\mathbb{k}} K)(\text{rank}_K H).$$

PROOF. Proofs seem to require considerable background in module theory and can be found in [Rad12, theorem 9.3.3] or [Mon93, chapter 3]. \square

The dimension formulae is of course a generalisation of *Lagrange's Theorem*: For a finite group G and $H \leq G$, $\mathbb{k}G$ is a finite dimensional Hopf algebra and $\mathbb{k}H$ is a subHopf algebra, with $\dim_{\mathbb{k}} \mathbb{k}G = |G|$ and $\dim_{\mathbb{k}} \mathbb{k}H = |H|$. Here is another nice generalisation.

COROLLARY 5.2. *Let H be a finite dimensional Hopf algebra. Then the grouplike elements form a finite subgroup $G(H) \leq H^\times$ and $|G(H)|$ divides $\dim_{\mathbb{k}} H$.*

PROOF. By Lemma 2.31, $G(H)$ is linearly independent so it must be finite with $|G(H)| \leq \dim_{\mathbb{k}} H$. In fact $G(H)$ spans the subHopf algebra $\mathbb{k}G(H) \subseteq H$, so $|G(H)| \mid \dim_{\mathbb{k}} H$ by Theorem 5.1. \square

5.2. Antipodes and finite dimensionality

In general the antipode of a Hopf algebra need not be bijective. But it often is, for example when the Hopf algebra is commutative or cocommutative. Here is another important case.

THEOREM 5.3. *Let H be a finite dimensional Hopf algebra. Then its antipode $\chi: H \rightarrow H$ is bijective.*

The proof will require a lemma which does not require H to be finite dimensional.

LEMMA 5.4. *Let H be a Hopf algebra and suppose that $K = \chi H \subseteq H$. If the restriction $\chi|_K: K \rightarrow K$ is a bijection then χ is a bijection.*

PROOF. The linear mapping $H \rightarrow K$ given by χ is surjective and $\chi|_K: K \rightarrow K$ is injective, so $H = \ker \chi \oplus K$ as vector spaces. Let $\pi: H \rightarrow K$ be projection onto the second factor; then $\ker \chi = \ker \pi$ and $\pi|_K = \text{Id}_K$.

By Proposition 2.32, $K \subseteq H$ is a subHopf algebra and $\ker \chi$ is a Hopf ideal of H . Hence $\varepsilon \ker \pi = \{0\}$ and

$$\psi \ker \chi \subseteq \ker \chi \otimes H + H \otimes \ker \chi = \ker \pi \otimes H + H \otimes \ker \chi,$$

so for $h \in \ker \chi$ and working with the convolution in $\text{hom}(H, H)$,

$$(\pi * \chi)(h) = 0 = \varepsilon(h)1.$$

For $k \in K$ we have

$$(\pi * \chi)(k) = \sum \pi(k_{(1)})\chi(k_{(2)}) = \sum k_{(1)}\chi(k_{(2)}) = \varepsilon(k)1.$$

It follows that π is the convolution inverse of χ , but this is Id_H . So in fact $\pi = \text{Id}_H$ and χ is surjective with $\ker \chi = 0$, hence χ is a bijection. \square

PROOF OF THEOREM 5.3. We will prove this in stages using a ‘downward induction’ argument. See Radford [Rad12, theorem 7.1.14] for more details on this.

Since $\chi: H \rightarrow H$ is a linear mapping and H is finite dimensional, *Fitting’s Lemma* implies that for some large enough n ,

$$H = \text{im } \chi^n \oplus \ker \chi^n.$$

where $K = \text{im } \chi^n = \chi^n H \subseteq H$ is a subHopf algebra on which the restriction of χ is injective. Since $K = \chi K = \chi(\chi^{n-1}H)$, we can apply Lemma 5.4 to the subHopf algebra $\chi^{n-1}H \subseteq H$ to deduce that χ is bijective on $\chi^{n-1}H$. Now we can repeat this argument to show that χ is bijective on each $\chi^k H$ with $1 \leq k \leq n-1$ and then show that it is bijective on H itself. \square

A finite submonoid of a group is always a subgroup (this is a routine consequence of the Pigeonhole Principle). A related result holds for Hopf algebras although the proof seems to require a more involved argument.

PROPOSITION 5.5. *Let H be a Hopf algebra and let $B \subseteq H$ be a subbialgebra that is finite dimensional. Then the antipode of H restricts to an antipode B , therefore B is a subHopf algebra.*

PROOF. Consider the convolution monoids $\text{hom}(B, B)$, $\text{hom}(B, H)$ and $\text{hom}(H, H)$; since B is a subbialgebra of H , $\text{hom}(B, B) \subseteq \text{hom}(B, H)$ is a submonoid and the inclusion $\text{inc}_B: B \rightarrow H$ induces a monoid homomorphism $\text{inc}_B: \text{hom}(H, H) \rightarrow \text{hom}(B, H)$. Let $\chi' = \chi \circ \text{inc}_B \in \text{hom}(B, H)$ be the restriction of the antipode $\chi: H \rightarrow H$ to B . The restriction of the identity Id_H to B is just the inclusion $\text{inc}_B: B \rightarrow H$, and

$$\chi' * \text{inc}_B = \text{inc}_B^*(\chi * \text{Id}_H) = \text{inc}_B^*(1_H)$$

which is the identity in $\text{hom}(B, H)$. So χ' is the $*$ -inverse of $\text{inc}_B \in \text{hom}(B, H)$.

Now $\text{hom}(B, B) \subseteq \text{hom}(B, H)$ is a submonoid and $(\text{inc}_B)_*(\text{Id}_B) = \text{inc}_B \circ \text{Id}_B = \text{inc}_B$. But $\text{hom}(B, B)$ and $\text{hom}(B, H)$ are also algebras with $\text{hom}(B, B) \subseteq \text{hom}(B, H)$ a subalgebra. Let $\Lambda: \text{hom}(B, B) \rightarrow \text{hom}(B, B)$ be the \mathbb{k} -linear endomorphism given by left multiplication by Id_B . Since $(\text{inc}_B)_*(\text{Id}_B) = \text{inc}_B \in \text{hom}(B, H)$ has a left inverse, it is injective, hence so is Λ . As B is finite dimensional so is $\text{hom}(B, B)$ and therefore Λ must be invertible. It follows that Id_B is invertible in $\text{hom}(B, B)$ under $*$, hence B has antipode χ_B making it a Hopf algebra. By construction, $\chi_B = \chi \circ \text{inc}_B$ so B is a subHopf algebra. \square

5.3. Hopf modules

The theory of *Hopf modules* is nowadays central in the study of finite dimensional Hopf algebras and its Fundamental Theorem provides a key ingredient for proofs of results such as the Larson-Sweedler Theorem.

Let $(H, \varphi, \eta, \psi, \varepsilon, \chi)$ be a Hopf algebra (not necessarily finite dimensional). Then H and $H \otimes H$ are both left H -modules and the coproduct $\psi: H \rightarrow H \otimes H$ is a module homomorphism since H is a bialgebra. Similarly, if M is a left H -module $H \otimes M$ is also an H -module.

DEFINITION 5.6. Suppose that M is a left H -module which is also a left H -comodule (M, μ) . Then (M, μ) is a (left) *H-Hopf module* if $\mu: M \rightarrow H \otimes M$ is an H -module homomorphism.

A homomorphism of *H-Hopf modules* $\theta: (M, \mu) \rightarrow (N, \nu)$ is a \mathbb{k} -linear mapping $\theta: M \rightarrow N$ which is both an H -module homomorphism and an H -comodule homomorphism.

If W is any vector space then $H \otimes W$ is both a left H -module and a left H -comodule and it is easy to check it is a Hopf module.

For a Hopf module (M, μ) we define its subspace of *coinvariants* to be

$$M_{\text{coinv}} = \{m \in M : \mu(m) = 1 \otimes m\} \subseteq M.$$

This vector subspace of M can be identified with the cotensor product $\mathbb{k} \square_H M$ where we view \mathbb{k} as a right H -comodule.

Here is the main result about Hopf modules, again we do not assume finite dimensionality.

THEOREM 5.7 (Fundamental Theorem of Hopf Modules). *Let M be an H -Hopf module. Then there is an isomorphism of Hopf modules*

$$H \otimes M_{\text{coinv}} \xrightarrow{\cong} M.$$

Hence every *H-Hopf module* is a free *H*-module.

PROOF. We start by defining the linear mapping

$$\Theta: H \otimes M_{\text{coinv}} \xrightarrow{\cong} M; \quad \Theta(h \otimes m) = hm.$$

Since M_{coinv} is just a vector space, this is a homomorphism of H -modules.

Let $h \in H$ and $m \in M_{\text{coinv}}$. The coaction applied to the element $hm \in M$ gives

$$\mu(hm) = \sum h_{(1)} 1 \otimes h_{(2)} m = \left(\sum h_{(1)} \otimes h_{(2)} \right) (1 \otimes m) = h \mu(m),$$

so this is a homomorphism of H -comodules.

Now for $m \in M$, let

$$\mu(m) = \sum m_{(1)} \otimes m_{(2)} \in H \otimes M.$$

Then

$$\begin{aligned} \mu\left(\sum \chi(m_{(1)})m_{(2)}\right) &= \sum \chi(m_{(1)})\mu(m_{(2)}) \\ &= \sum \chi(m_{(1)})_{(1)}(m_{(2)})_{(1)} \otimes \chi(m_{(1)})_{(2)}(m_{(2)})_{(2)} \\ &= \sum \chi(m_{(12)})(m_{(21)}) \otimes \chi(m_{(11)})(m_{(22)}) \\ &= \sum \chi(m_{(121)})(m_{(122)}) \otimes \chi(m_{(11)})(m_{(2)}) \\ &= \sum \varepsilon(m_{(12)}) \otimes \chi(m_{(11)})(m_{(2)}) \\ &= \sum 1 \otimes \varepsilon(m_{(12)})\chi(m_{(11)})(m_{(2)}) \\ &= \sum 1 \otimes \chi(m_{(1)})(m_{(2)}), \end{aligned}$$

hence $\sum \chi(m_{(1)})m_{(2)} \in M_{\text{coinv}}$.

Now consider the \mathbb{k} -linear map $\Psi: M \rightarrow H \otimes M_{\text{coinv}}$ given by

$$\Psi(m) = \sum m_{(1)} \otimes \chi((m_{(2)})_{(1)})(m_{(2)})_{(2)}.$$

Since H acts trivially on M_{coinv} , this is an H -module homomorphism and an H -comodule homomorphism by coassociativity. Also,

$$\begin{aligned} \Theta\Psi(m) &= \Theta\left(\sum m_{(1)} \otimes \chi((m_{(2)})_{(1)})(m_{(2)})_{(2)}\right) \\ &= \Theta\left(\sum (m_{(1)})_{(1)} \otimes \chi((m_{(1)})_{(2)})m_{(2)}\right) \\ &= \sum (m_{(1)})_{(1)} \chi((m_{(1)})_{(2)})m_{(2)} \\ &= \sum \varepsilon(m_{(1)})m_{(2)} \\ &= m, \end{aligned}$$

and when $\mu(m) = 1 \otimes m$,

$$\begin{aligned} \Psi\Theta(h \otimes m) &= \Psi(hm) = \sum h_{(1)} \otimes \chi((h_{(2)})_{(1)})(h_{(2)})_{(2)}m \\ &= \sum h_{(1)} \otimes \varepsilon(h_{(2)})m \\ &= \sum \varepsilon(h_{(2)})h_{(1)} \otimes m \\ &= h \otimes m. \end{aligned}$$

Therefore Ψ and Θ are inverse functions.

If we choose a \mathbb{k} -basis for M_{coinv} then we get an H basis for $H \otimes M_{\text{coinv}}$, hence $M \cong H \otimes M_{\text{coinv}}$ is a free H -module. \square

This result tells us that for a non-trivial Hopf module M , M_{coinv} is also non-trivial. But we can say more when we have appropriate finiteness conditions.

COROLLARY 5.8. *If M_{coinv} is finite dimensional then*

$$\text{rank}_H M = \dim_{\mathbb{k}} M_{\text{coinv}},$$

and if H is also finite dimensional then

$$\dim_{\mathbb{k}} M = \dim_{\mathbb{k}} H \dim_{\mathbb{k}} M_{\text{coinv}}.$$

Here is another interesting application. For a Hopf algebra H , a subspace $L \subseteq H$ is a *left coideal* if the image of the coproduct applied to L satisfies $\psi L \subseteq H \otimes L$, so L is a subcomodule of H .

COROLLARY 5.9. *If H is finite dimensional and a non-zero left ideal $I \subseteq H$ is also a left coideal, then $I = H$.*

PROOF. The conditions imply that I is a Hopf module which is a subHopf module of H . By the Fundamental Theorem,

$$I \cong H \otimes I_{\text{coinv}}$$

as H -modules, so $\dim_{\mathbb{k}} I \geq \dim_{\mathbb{k}} H$ which is only possible if $I = H$. \square

5.4. Applications to finite dimensional Hopf algebras

Now we will apply our results on Hopf modules to study finite dimensional Hopf algebras.

THEOREM 5.10. *If H is a finite dimensional Hopf algebra then its dual H^* is an H -Hopf module which is free of rank 1 as an H -module, i.e., $H^* \cong H$ as left H -modules.*

PROOF. The dual $H^* = \text{hom}(H, \mathbb{k})$ is both a left H -module where for $h \in H$ and $f \in H^*$,

$$h \cdot f = f(\chi(h)-).$$

It is also an algebra where the product is obtained by dualising the coproduct of H , i.e., it is the composition

$$H^* \otimes H^* \xleftarrow[\cong]{\psi^\dagger} (H \otimes H)^* \xrightarrow[\psi^*]{} H^*$$

In fact this is a homomorphism of left H -modules where we use the antipode and the left multiplication on the domains of $H^* = \text{hom}(H, \mathbb{k})$ and $(H \otimes H)^* = \text{hom}(H \otimes H, \mathbb{k})$ to define their module structures.

Now we make H^* into a Hopf module over H by defining the coaction $\mu: H^* \rightarrow H \otimes H^*$ as follows: for $f \in H^*$,

$$\mu(f) = \sum f_{(1)} \otimes f_{(2)} \in H \otimes H^*$$

where the terms $f_{(1)}$ are characterised by requiring that for all $g \in H^*$, the product $fg \in H^*$ satisfies

$$fg = \sum g(f_{(1)})f_{(2)}.$$

A verification that this is an H -module homomorphism can be found in the proof of [Lor18, theorem 10.9].

The Fundamental Theorem tells us that H^* is a free module and since $\dim_{\mathbb{k}} H^* = \dim_{\mathbb{k}} H$ it must have rank 1, i.e., $H^* \cong H$ as H -modules. \square

Since H^* is an injective H -module this result says that H is also injective as well as projective, i.e., it is *self-injective*.

We can also give another proof of Theorem 5.3. For if $z \in \ker \chi \subseteq H$ then for any $f \in H^*$,

$$z \cdot f = f(\chi(z)-) = 0,$$

but since $H^* \cong H$, this is only possible if $z = 0$.

We now have an important result on finite dimensional Hopf algebras. A graded analogue of this was proved by Browder & Spanier [BS62], then the ungraded case was proved by Larson & Sweedler [LS69].

THEOREM 5.11 (Larson & Sweedler). *If H is a finite dimensional Hopf algebra then it is a Frobenius algebra.*

PROOF. The existence of a left H -module isomorphism $H \xrightarrow{\cong} H^*$ gives us an element $\lambda \in H^*$ which is the image of $1 \in H$. By definition of the module structure on H^* , the image of $h \in H$ is then $h\lambda \in H^*$ where

$$(h\lambda)(x) = \lambda(xh).$$

If this λ is trivial on some simple left submodule $S \subseteq H$ then for any non-zero element $s \in S$, $s\lambda = 0$, contradicting the definition of λ . It follows that λ is a Frobenius form and so H is a Frobenius algebra. \square

This result has many interesting consequences. An algebra A is called a *Kasch algebra* if every left or right simple A -module is isomorphic to a minimal left or right ideal of A (so these are its simple submodules).

If S is a (non-trivial) simple left or right A -module, then by Schur's Lemma its endomorphism algebra $\text{End}_A(S)$ is a central division algebra over \mathbb{k} . It is easy to see that the sum of all the submodules of A isomorphic to S is actually a finite direct sum

$$S_1 \oplus S_2 \oplus \cdots \oplus S_m,$$

where for each i , $S_i \cong S$. The number m is well-defined and is called the *multiplicity* of S in A .

PROPOSITION 5.12. *Every Frobenius algebra A is a Kasch algebra. In particular, if S is a simple A -module then its multiplicity in A is equal to $\dim_{\text{End}_A(S)} S$.*

PROOF. When S is a non-trivial simple left A -module, the opposite division algebra $E = \text{End}_A(S)^{\text{op}}$ acts on $\text{Hom}_A(S, A)$ by precomposition making it a left E -module. There are isomorphisms of E -vector spaces

$$\begin{aligned} \text{Hom}_A(S, A) &\cong \text{Hom}_A(S, A^*) = \text{Hom}_A(S, \text{hom}(A, \mathbb{k})) \\ &\cong \text{Hom}_{\mathbb{k}}(A \otimes_A S, \mathbb{k}) \\ &\cong \text{Hom}_{\mathbb{k}}(S, \mathbb{k}). \end{aligned}$$

Every non-trivial A -module homomorphism $S \rightarrow A$ must be injective by simplicity, so the multiplicity of S is

$$\dim_E \text{Hom}_A(S, A) = \dim_{\text{End}_A(S)} S \neq 0. \quad \square$$

If $\dim_{\mathbb{k}} S = 1$, then $\text{End}_A(S) = \mathbb{k}$ and $\dim_{\mathbb{k}} \text{Hom}_A(S, A) = 1$, so S occurs with multiplicity 1, i.e., there is a unique submodule of A isomorphic to S .

DEFINITION 5.13. Of course Proposition 5.12 applies to any finite dimensional Hopf algebra. In particular the counit $\varepsilon: H \rightarrow \mathbb{k}$ gives us a 1-dimensional simple left or right module and each of these occurs as a unique submodule. These 1-dimensional subspaces are called the spaces of left or right *integrals* of H :

$$\int_H^l = \{z \in H : \forall h \in H, hz = \varepsilon(h)z\}, \quad \int_H^r = \{z \in H : \forall h \in H, zh = \varepsilon(h)z\}.$$

In general, $\int_H^l \neq \int_H^r$, but if $\int_H^l = \int_H^r$ then H is called *unimodular* and we set $\int_H = \int_H^l = \int_H^r$. In general,

$$\chi \int_H^l = \int_H^r, \quad \chi \int_H^r = \int_H^l,$$

so when H is unimodular,

$$\chi \int_H = \int_H.$$

We can generalise Proposition 5.12.

PROPOSITION 5.14. *For a Frobenius algebra A , the following are true.*

- (a) *If M is a cyclic left A -module then its \mathbb{k} -dual M^* is a right A -module isomorphic to a submodule of A viewed as a right A -module.*
- (b) *If $N \subseteq A$ is a right A -submodule then its \mathbb{k} -dual N^* is a cyclic left A -module.*

Furthermore these results hold with left and right interchanged throughout.

PROOF. (a) The right module structure on $M^* = \text{hom}(M, \mathbb{k})$ is given by

$$(f \cdot a)(x) = f(ax) \quad (x \in M)$$

for $a \in A$ and $f \in \text{hom}(M, \mathbb{k})$.

Since M is cyclic there is an epimorphism of A -modules $A \rightarrow M$ and on applying $\text{hom}(-, \mathbb{k})$ we obtain a homomorphism of right A -modules $M^* \rightarrow A^*$ which is an injective \mathbb{k} -linear mapping. (b) This is similar to (b) and is left as an exercise. \square

REMARK 5.15. Recall that the antipode of a Hopf algebra H is a homomorphism $H \rightarrow H^{\text{op}}$ which must be an isomorphism when H is finite dimensional. This allows us to treat a right module as a left module and vice versa. In particular we can use the antipode to make the dual of a left H -module into a left H -module, so we can apply this result to deduce that the dual of a cyclic left module is isomorphic to a left submodule of H , and the dual of a left submodule of H is cyclic.

We mention that for infinite dimensional Hopf algebras things are very different.

PROPOSITION 5.16. *Let H be an infinite dimensional Hopf algebra. Then H has no non-trivial finite dimensional left or right ideals. In particular, H is not a Kasch algebra.*

PROOF. See Lorenz [Lor18, proposition 10.6]. Since the trivial module is simple and does not embed in H , hence H is not a Kasch algebra. \square

Although we know that a finite dimensional Hopf algebra H has a Frobenius form we have not yet explained how to find a suitable element of H^* .

LEMMA 5.17. *Let H a finite dimensional Hopf algebra. Then in the dual Hopf algebra H^* , any non-zero right integral $\lambda \in \int_{H^*}^r$ is a Frobenius form for H .*

PROOF. See Radford [Rad12, theorem 10.2.2(e)] or the proof of Montgomery [Mon93, theorem 2.1.3(4)]. \square

EXAMPLE 5.18. Let $\mathbb{k}G$ be the group algebra of a finite group. Every element can be uniquely written as $\sum_{g \in G} t_g g$ where t_g . The element $z_0 = \sum_{g \in G} g$ satisfies

$$\left(\sum_{g \in G} t_g g\right) z_0 = \left(\sum_{g \in G} t_g\right) z_0 = \varepsilon\left(\sum_{g \in G} t_g g\right) z_0 = z_0 \left(\sum_{g \in G} t_g g\right),$$

so $\mathbb{k}G$ is unimodular and

$$\int_{\mathbb{k}G} = \{tz_0 : t \in \mathbb{k}\}.$$

Define the form $\lambda \in (\mathbb{k}G)^*$ by

$$\lambda\left(\sum_{g \in G} t_g g\right) = t_1,$$

so $\lambda = \delta_1$, the Dirac function of (3.1). For any $g \in G$,

$$\delta_g \lambda = \delta_g \delta_1 = \begin{cases} \delta_1 = \lambda & \text{if } g = 1, \\ 0 & \text{if } g \neq 1, \end{cases}$$

which shows that λ is a left and a right integral for the dual Hopf algebra $(\mathbb{k}G)^*$, hence it is a Frobenius form for $\mathbb{k}G$.

EXAMPLE 5.19 (Taft algebras). Recall the Taft Hopf algebra $H_{n,\zeta}$ of Section 3.12:

$$H_{n,\zeta} = \mathbb{k}\langle u, v \rangle / (u^n - 1, v^n, vu - \zeta uv).$$

The element

$$\sum_{0 \leq i \leq n-1} u^i v^{n-1}$$

is a left integral and

$$\sum_{0 \leq i \leq n-1} \zeta^i u^i v^{n-1} = \sum_{0 \leq i \leq n-1} v^{n-1} u^i$$

is a right integral.

Any linear map $\lambda: H_{n,\zeta} \rightarrow \mathbb{k}$ for which

$$\lambda\left(\sum_{0 \leq i \leq n-1} a_i u^i v^{n-1}\right) = a_0$$

for $a_i \in \mathbb{k}$ is a Frobenius form – of course there are many choices!

5.5. Semisimplicity for finite dimensional Hopf algebras

For a finite dimensional algebra A it is important to know whether it is semisimple and so its modules are completely reducible (i.e., direct sums of simple modules). Semisimplicity is equivalent to the triviality of the Jacobson radical of the algebra, and more generally, for any finite dimensional algebra A , the quotient algebra $A/\text{rad } A$ is semisimple and so it is isomorphic to a product of matrix rings over division algebras; this is the main content of *Artin-Wedderburn Theory*. Before stating a version of *Maschke's Theorem* for finite dimensional Hopf algebras we give a preliminary result which contains the main ingredients.

LEMMA 5.20 (Maschke Lemma). *Let H be a finite dimensional Hopf algebra and suppose that the space of left integrals satisfies $\varepsilon \int_H^1 \neq 0$.*

(a) *The space of left integrals \int_H^1 contains a unique non-zero idempotent element e which also has $\varepsilon(e) = 1$.*

(b) *Let M be a left H -module. Then the space of H -invariants satisfies*

$$M^H = eM = \{em : m \in M\}$$

and there is a direct sum splitting of vector spaces

$$M = eM \oplus (1 - e)M = M^H \oplus (1 - e)M$$

where

$$(1 - e)M = \{(1 - e)m : m \in M\}.$$

(c) *Let*

$$0 \rightarrow L \xrightarrow{f} M \xrightarrow{g} N \rightarrow 0$$

be a short exact sequence of H -modules. Then there is an induced short exact sequence

$$0 \rightarrow L^H \xrightarrow{f'} M^H \xrightarrow{g'} N^H \rightarrow 0.$$

PROOF. (a) Let $z \in \int_H^1$ with $\varepsilon(z) \neq 0$. Then $z^2 = \varepsilon(z)z$ and so

$$(\varepsilon(z)^{-1}z)^2 = \varepsilon(z)^{-2}(z^2) = \varepsilon(z)^{-2}(\varepsilon(z)z) = \varepsilon(z)^{-1}z.$$

So we may take $e = \varepsilon(z)^{-1}z$. Uniqueness involves a routine calculation.

(b) If $m \in M$ then

$$h(em) = (he)m = \varepsilon(h)em$$

so $em \in M^H$. On the other hand, if $n \in M^H$ then

$$en = \varepsilon(e)n = n,$$

so $n \in eM$. Together these containments show that $M^H = eM$.

Since e is an idempotent, so is $1 - e$, and a standard linear algebra argument gives the direct sum decomposition.

(c) Since f and g are H -module homomorphism the images of their restrictions to $L^H = eL$ and $M^H = eM$ are contained in $eM = M^H$ and $eN = N^H$ respectively. There is a natural isomorphism of functors

$$(-)^H \cong \text{Hom}_H(\mathbb{k}, -)$$

where the latter is left exact, so we obtain an exact sequence

$$0 \rightarrow L^H \xrightarrow{f'} M^H \xrightarrow{g'} N^H.$$

Let $n \in N^H$. Then by short exactness of the original sequence, there is an element $m \in M$ with $g(m) = n$. But then

$$g'(em) = g(em) = eg(m) = en = n.$$

Therefore g' is also surjective. \square

Here is a result which is a generalisation of a fundamental result in group representation theory.

THEOREM 5.21 (Maschke's Theorem). *Let H be a finite dimensional Hopf algebra. The following conditions are equivalent:*

- (a) *Every left H -module is completely reducible, so H is left semisimple.*
- (b) *For any non-zero left integral $z \in \int_H^1$, $\varepsilon(z) \neq 0$.*
- (c) *For any non-zero right integral $z \in \int_H^r$, $\varepsilon(z) \neq 0$.*
- (d) *Every right H -module is completely reducible, so H is right semisimple.*

If these conditions hold then H is semisimple and unimodular.

PROOF. For detailed discussions see Radford [**Rad12**, theorem 10.8.2] or Montgomery [**Mon93**, theorem 2.2.1].

The underlying idea is to use our Maschke Lemma 5.20 to show that given a short exact sequence of H -modules

$$0 \rightarrow L \xrightarrow{f} M \xrightarrow{g} N \rightarrow 0$$

the induced (split) short exact sequence of vector spaces

$$0 \rightarrow \text{hom}(N, L) \xrightarrow{f_*} \text{hom}(N, M) \xrightarrow{g_*} \text{hom}(N, N) \rightarrow 0$$

is actually a short exact sequence of H -modules, where we defined the module structure in Section 4.4 and in (4.6) showed that for H -modules U, V ,

$$\text{hom}(U, V)^H = \text{Hom}_H(U, V).$$

Applying part (b) of the Maschke Lemma we get a short exact sequence

$$0 \rightarrow \text{Hom}_H(N, L) \xrightarrow{f_*} \text{Hom}_H(N, M) \xrightarrow{g_*} \text{Hom}_H(N, N) \rightarrow 0.$$

So if we take any $\ell \in \text{hom}(N, M)$ with $g_*(\ell) = \text{Id}_N \in \text{Hom}_H(N, N)$ we find that in the last sequence, $e \cdot \ell \in \text{Hom}_H(N, M)$ and $g_*(e \cdot \ell) = \text{Id}_N$. This provides an H -module splitting homomorphism $e \cdot \ell: N \rightarrow M$ for the original sequence of H -modules. \square

A finite dimensional Hopf algebra which is semisimple has a representation theory very similar to that of a finite group over a field whose characteristic does not divide its order. In the setting of Example 5.18,

$$\varepsilon(z_0) = \varepsilon\left(\sum_{g \in G} g\right) = |G| \in \mathbb{k},$$

so $\mathbb{k}G$ is semisimple if and only if the characteristic of \mathbb{k} does not divide $|G|$ and in that case the idempotent e is given by

$$e = \frac{1}{|G|} \sum_{g \in G} g.$$

Of course this is a well-known fact in the representation theory of finite groups and is a consequence of Maschke's Theorem; the action of e on a linear section is used to give a formula for the splitting which is equivalent the one above.

For a semisimple finite dimensional Hopf algebra H , every simple submodule of H is a direct summand and $1 \in H$ can be expressed as a unique sum of central primitive idempotents.

Of course not all finite dimensional Hopf algebras are semisimple.

EXAMPLE 5.22. An important class of non-semisimple examples is provided by Hopf algebras which are *local*. Such an algebra has a unique maximal left/right ideal which must agree with its Jacobson radical. A local algebra also has a unique simple left/right module. For a local finite dimensional Hopf algebra H , the unique simple module is the trivial module \mathbb{k} and this occurs in H with multiplicity 1. Examples of this are provided by group algebras of finite p -groups over fields of characteristic p .

EXAMPLE 5.23. Let p be a prime number and $r \geq 1$. For a field \mathbb{k} of characteristic p the cyclic group $C_{p^r} = \langle \gamma \rangle$ of order p^r has group algebra $\mathbb{k}C_{p^r}$. We can replace the usual basis of powers of γ by the powers of $\tilde{\gamma} = \gamma - 1$,

$$1, \tilde{\gamma}, \tilde{\gamma}^2, \dots, \tilde{\gamma}^{p^r-1}.$$

Then as an algebra,

$$\mathbb{k}C_{p^r} \cong \mathbb{k}[X]/(X^{p^r}); \quad \tilde{\gamma}^k \leftrightarrow X^k + (X^{p^r}).$$

If we set $x = X + (X^{p^r})$ the coproduct on $\mathbb{k}[X]/(X^{p^r})$ is given by

$$\psi(x) = x \otimes 1 + 1 \otimes x + x \otimes x.$$

Of course this algebra is commutative and its unique maximal ideal is $(x) \triangleleft \mathbb{k}[X]/(X^{p^r})$. The space of integrals is the 1-dimensional subspace

$$\int_{\mathbb{k}[X]/(X^{p^r})} = \mathbb{k}\{x^{p^r-1}\}.$$

EXAMPLE 5.24. For the Taft algebra $H_{n,\zeta}$, the left and right integrals identified in Example 5.19 satisfy

$$\varepsilon\left(\sum_{0 \leq i \leq n-1} u^i v^{n-1}\right) = 0 = \varepsilon\left(\sum_{0 \leq i \leq n-1} \zeta^i u^i v^{n-1}\right)$$

so $H_{n,\zeta}$ is not semisimple; its Jacobson radical is the ideal generated by v .

CHAPTER 6

A brief introduction to quantum groups

This chapter is meant to provide some indicators of what the study of quantum groups involves and it is certainly very incomplete. The literature on the subject is enormous and certainly extends outside of purely algebraic contexts.

As Hopf algebras, quantum groups are the opposite of classical Hopf algebras: they are neither commutative nor cocommutative. Many examples arise as ‘deformations’ of classical ones, typically obtained by inserting a parameter q so that in the limit as $q \rightarrow 1$ the classical case is recovered. Alternatively, a parameter \hbar is used so that $q = e^{\hbar}$ and passing to the limit $\hbar \rightarrow 0$ recovers the classical version. All of this is of course suggested by phenomena in Quantum Mechanics which is itself involves a kind of ‘deformation’ of Classical Mechanics.

6.1. q -combinatorics

Suppose that we are working in a non-commutative ring R where $q, x, y \in R$ with $q \neq 1$ in the centre of R and the other elements satisfy

$$yx = qxy.$$

What is the analogue of the usual binomial expansion of $(x + y)^n$ when $n \geq 1$?

To describe the answer we introduce analogues of standard combinatorial expressions. We will set

$$[n]_q = \frac{q^n - 1}{q - 1} = q^{n-1} + q^{n-2} + \cdots + q + 1,$$

so

$$\lim_{q \rightarrow 1} [n]_q = n.$$

First we have the q -factorials; these are defined recursively for $n \geq 0$:

$$[0]_q! = 1, \quad [n]_q! = [n]_q([n-1]_q!) = \frac{q^n - 1}{q - 1} [n-1]_q!,$$

so

$$[n]_q! = \prod_{1 \leq k \leq n} (q^{k-1} + q^{k-2} + \cdots + q + 1).$$

Notice that

$$\lim_{q \rightarrow 1} [n]_q! = n!.$$

Next we have the q -binomial coefficients for $0 \leq k \leq n$:

$$\begin{bmatrix} n \\ k \end{bmatrix}_q = \frac{[n]_q!}{[k]_q! [n-k]_q!}.$$

These satisfy

$$\lim_{q \rightarrow 1} \begin{bmatrix} n \\ k \end{bmatrix}_q = \binom{n}{k}.$$

as well as two generalisations of Pascal's Triangle which are easily verified:

$$(6.1) \quad \begin{bmatrix} n \\ k \end{bmatrix}_q = \begin{bmatrix} n-1 \\ k-1 \end{bmatrix}_q + q^k \begin{bmatrix} n-1 \\ k \end{bmatrix}_q,$$

$$(6.2) \quad \begin{bmatrix} n \\ k \end{bmatrix}_q = q^{n-k} \begin{bmatrix} n-1 \\ k-1 \end{bmatrix}_q + \begin{bmatrix} n-1 \\ k \end{bmatrix}_q.$$

Now working in the non-commutative ring R we have

PROPOSITION 6.1. For $n \geq 1$,

$$(x+y)^n = \sum_{0 \leq k \leq n} \begin{bmatrix} n \\ k \end{bmatrix}_q x^k y^{n-k}.$$

PROOF. This can be proved by induction on n using one of the identities

$$(x+y)(x+y)^{n-1} = (x+y)^n = (x+y)^{n-1}(x+y)$$

and one of the identities (6.1) or (6.2). □

Here is an application. Suppose that $z \in R$ is nilpotent; the q -exponential of z is

$$\exp_q(z) = \sum_{k \geq 0} \frac{1}{[k]_q!} z^k$$

which is of course a finite sum. Notice that

$$\lim_{q \rightarrow 1} \exp_q(z) = \exp(z).$$

Now suppose that $x, y \in R$ as above are also nilpotent; then for some large enough m , for any $0 \leq i \leq m$ we have $x^i y^{m-i}$. So we have

$$(6.3) \quad \exp_q(x+y) = \exp_q(x) \exp_q(y).$$

To see this, expand out the left hand side to obtain

$$\begin{aligned} \exp_q(x+y) &= \sum_{k \geq 0} \frac{1}{[k]_q!} (x+y)^k \\ &= \sum_{k \geq 0} \frac{1}{[k]_q!} \left(\sum_{0 \leq i \leq k} \begin{bmatrix} k \\ i \end{bmatrix}_q x^i y^{k-i} \right) \\ &= \sum_{k \geq 0} \sum_{0 \leq i \leq k} \frac{1}{[i]_q!} x^i \frac{1}{[k-i]_q!} y^{k-i} \\ &= \sum_{i \geq 0} \sum_{j \geq 0} \frac{1}{[i]_q!} x^i \frac{1}{[j]_q!} y^j \\ &= \exp_q(x) \exp_q(y), \end{aligned}$$

where the sums are really finite.

For polynomials in a variable X , there is a q -derivative ∂_q given by

$$\partial_q f(X) = \frac{f(qX) - f(X)}{(q-1)X},$$

so for example,

$$\partial_q X^n = [n]_q X^{n-1}.$$

For nilpotent z , we have

$$\partial_q \exp_q(z) = \exp_q(z).$$

6.2. The Quantum Plane

Recall the Quantum Plane of Example 2.12, the non-commutative bialgebra $\mathcal{O}_q(\mathbb{k}^2)$ where $q \neq 1$, generated by two elements x, y satisfying $yx = qxy$. The coproduct ψ and counit ε are given by

$$\psi(x) = x \otimes x, \quad \psi(y) = y \otimes 1 + x \otimes y, \quad \varepsilon(x) = 1, \quad \varepsilon(y) = 0.$$

The quantum version of the general linear group for $1 \neq q \in \mathbb{k}^\times$ is a Hopf algebra $\mathcal{GL}_q(2)$ which we will now define. As an algebra, $\mathcal{GL}_q(2)$ is generated by a, b, c, d, e satisfying the relations

$$\begin{aligned} ca &= qac, & ba &= qab, & db &= qbd, \\ dc &= qcd, & cb &= bc, & da - ad &= (q - q^{-1})bc, \\ (ad - q^{-1}bc)e &= 1. \end{aligned}$$

It turns out that $(ad - q^{-1}bc)$ is in the centre of $\mathcal{GL}_q(2)$, hence so is

$$e = (ad - q^{-1}bc)^{-1}.$$

We can also define the quotient Hopf algebra $\mathcal{SL}_q(2)$ where we have the additional relations

$$ad - q^{-1}bc = 1 = e.$$

This is called the *quantum special linear group*. The coproduct, counit and antipode are given by

$$\begin{aligned} \psi(a) &= a \otimes a + b \otimes c, & \psi(b) &= b \otimes d + a \otimes b, & \psi(c) &= c \otimes a + d \otimes b, & \psi(d) &= d \otimes d + c \otimes b, \\ \varepsilon(a) &= 1 = \varepsilon(d), & \varepsilon(b) &= 0 = \varepsilon(c), \\ \chi(a) &= d, & \chi(d) &= a, & \chi(b) &= c, & \chi(c) &= b. \end{aligned}$$

Just as the special linear group acts linearly on the plane, so the quantum special linear group coacts on the quantum plane, i.e., there is a coaction $\rho: \mathcal{O}_q(\mathbb{k}^2) \rightarrow \mathcal{SL}_q(2) \otimes \mathcal{O}_q(\mathbb{k}^2)$. This is given on the generators by

$$\rho(x) = a \otimes x + b \otimes y, \quad \rho(y) = c \otimes x + d \otimes y.$$

6.3. Quasitriangular Hopf algebras

For a non-cocommutative Hopf algebra H , its module category \mathbf{Mod}_H is monoidal under tensor product but not always *symmetric* monoidal since in general $M \otimes N$ need not be isomorphic to $N \otimes M$ as H -modules. One way to ‘correct’ this is to impose extra structure. There are detailed discussions of this and examples in the books [BG02, Maj95, Maj02] with the last being particularly suitable for a quick introduction.

DEFINITION 6.2. A *quasitriangular Hopf algebra* (H, \mathcal{R}) is a Hopf algebra H with an element $\mathcal{R} \in H \otimes H$ satisfying the following conditions.

- \mathcal{R} is a unit in the algebra $H \otimes H$ and for all $h \in H$,

$$\mathrm{T} \circ \psi(h) = \mathcal{R}(\psi(h))\mathcal{R}^{-1}.$$

- In the algebra $H \otimes H \otimes H$ we have the identities

$$(\psi \otimes \mathrm{Id}_H)(\mathcal{R}) = \mathcal{R}_{13}\mathcal{R}_{23}, \quad (\mathrm{Id}_H \otimes \psi)(\mathcal{R}) = \mathcal{R}_{13}\mathcal{R}_{12},$$

where $\mathcal{R}_{ij} \in H \otimes H \otimes H$ means the image of \mathcal{R} under the algebra homomorphism $H \otimes H \rightarrow H \otimes H \otimes H$ obtained by including the i and j factors (so $\mathcal{R}_{12} = \mathcal{R} \otimes 1$ and $\mathcal{R}_{23} = 1 \otimes \mathcal{R}$ for example).

LEMMA 6.3. Suppose that (H, \mathcal{R}) is a quasitriangular Hopf algebra.

(a) We have

$$\begin{aligned}(\varepsilon \otimes \text{Id}_H)(\mathcal{R}) &= 1 = (\text{Id}_H \otimes \varepsilon)(\mathcal{R}), \\(\chi \otimes \text{Id}_H)(\mathcal{R}) &= \mathcal{R}^{-1}, \\(\text{Id}_H \otimes \chi)(\mathcal{R}^{-1}) &= \mathcal{R},\end{aligned}$$

and therefore

$$(\chi \otimes \chi)(\mathcal{R}) = \mathcal{R},$$

so \mathcal{R} is χ -invariant.

(b) $(H, \mathcal{R}_{21}^{-1})$ is also a quasitriangular Hopf algebra where

$$\mathcal{R}_{21}^{-1} = T(\mathcal{R}^{-1}).$$

(c) The Yang-Baxter identity holds in $H \otimes H \otimes H$, i.e.,

$$\mathcal{R}_{12}\mathcal{R}_{13}\mathcal{R}_{23} = \mathcal{R}_{23}\mathcal{R}_{13}\mathcal{R}_{12}.$$

Notice that the Yang-Baxter equation is similar to the following identity in the symmetric group S_3 :

$$(1\ 2)(1\ 3)(2\ 3) = (1\ 3) = (2\ 3)(1\ 3)(1\ 2).$$

It is also a relation in the 3-rd braid group so is sometimes called the *braid relation*.

THEOREM 6.4. Suppose that (H, \mathcal{R}) is a quasitriangular Hopf algebra. Then

- (a) the antipode of H is a bijection;
- (b) there is a unit $u \in H^\times$ such that

$$\chi^2 = u(-)u^{-1},$$

and moreover

$$\psi(u) = (T(\mathcal{R})\mathcal{R})^{-1}(u \otimes u).$$

PROOF. See Majid [Maj02, theorem 5.7]. □

COROLLARY 6.5. The element $v = \chi(u)$ has the following properties.

- For all $h \in H$, $\chi^{-2}(h) = v h v^{-1}$.
- The coproduct on v is $\psi(v) = (T(\mathcal{R})\mathcal{R})^{-1}(v \otimes v)$.
- The elements u and v commute, $uv = vu$, and this element is central with coproduct

$$\psi(uv) = (T(\mathcal{R})\mathcal{R})^{-2}(uv \otimes uv).$$

- The element $uv^{-1} = v^{-1}u$ and for all $h \in H$, $\chi^4(h) = (uv^{-1})h(uv^{-1})^{-1}$.

PROOF. See Majid [Maj02, corollary 6.1]. □

Given this result we make a definition.

DEFINITION 6.6. A quasitriangular Hopf algebra (H, \mathcal{R}) is called *ribbon* if uv has a central square root ν called a *ribbon element* such that

$$\nu^2 = uv, \quad \psi(\nu) = (T(\mathcal{R})\mathcal{R})^{-1}(\nu \otimes \nu), \quad \varepsilon(\nu) = 1, \quad \chi(\nu) = \nu.$$

To illustrate the impact of having a quasitriangular structure on a Hopf algebra, recall from Lemma 4.29 that for a finite dimensional H -module M , $M^{**} \cong (\chi^2)^*M$. Using (b) it is easy to see that for a quasitriangular Hopf algebra we have $(\chi^2)^*M \cong M$ and therefore $M^{**} \cong M$. We will see far more is true.

Writing down explicit examples takes work and is the subject of a lot of literature on Quantum Groups, so rather than go into details we refer the interested reader to other sources. A good introduction is provided by Majid [Maj02], while [Maj95] and Chari & Pressley [CP94] have even more detail. More traditional algebraic aspects of the subject can be found in Brown & Goodearl [BG02].

6.4. Braidings on module categories of quasitriangular Hopf algebras

In the following we assume that (H, \mathcal{R}) is a quasitriangular Hopf algebra. We will often write \mathcal{R} using Sweedler-style notation as a sum

$$\mathcal{R} = \sum \mathcal{R}_1 \otimes \mathcal{R}_2.$$

The module category \mathbf{Mod}_H is monoidal under \otimes . We define

$$M \overset{\text{op}}{\otimes} N = N \otimes M$$

with the usual H -action given by multiplication by $T \circ \psi(h)$:

$$h(m \overset{\text{op}}{\otimes} n) = h(n \otimes m) = \sum h_{(1)}n \otimes h_{(2)}m = \sum h_{(2)}m \overset{\text{op}}{\otimes} h_{(1)}n.$$

LEMMA 6.7. *For two left H -modules M and N ,*

$$\Psi_{M,N}: M \otimes N \rightarrow M \overset{\text{op}}{\otimes} N = N \otimes M; \quad \Psi_{M,N}(m \otimes n) = \sum \mathcal{R}_1 m \overset{\text{op}}{\otimes} \mathcal{R}_2 n = \sum \mathcal{R}_2 n \otimes \mathcal{R}_1 m$$

defines an isomorphism of H -modules.

PROOF. Notice that

$$\Psi_{M,N} = T \circ \mathcal{R}$$

where \mathcal{R} means the multiplication by \mathcal{R} function on $H \otimes H$. By the first part of Definition 6.2, for $h \in H$,

$$\psi(h) \circ \mathcal{R} = T \circ \mathcal{R} \circ \psi(h).$$

We have for $h \in H$, $m \in M$ and $n \in N$,

$$\begin{aligned} \Psi_{M,N}(h(m \otimes n)) &= T(\mathcal{R}\psi(h)(m \otimes n)) \\ &= T \circ \mathcal{R} \circ \psi(h)(m \otimes n) \\ &= \psi(h) \circ \mathcal{R}(m \otimes n) \\ &= \psi(h) \circ T \circ T \circ \mathcal{R}(m \otimes n) \\ &= T \circ \psi(h) \circ \Psi_{M,N}(m \otimes n) \\ &= h\Psi_{M,N}(m \otimes n). \end{aligned}$$

It is clear that $\Psi_{M,N}$ does has an inverse, namely $\mathcal{R}^{-1} \circ T$. □

The existence of \mathcal{R} makes the monoidal category $(\mathbf{Mod}_H, \otimes)$ into a *braided monoidal category*. This involves $\Psi_{-, -}$ as well as functorial isomorphisms

$$\Phi_{U,V,W}: (U \otimes V) \otimes W \rightarrow U \otimes (V \otimes W)$$

which obey the *Pentagon Condition* making the following diagram commute for all U, V, W, Z .

$$\begin{array}{ccccc}
& & (U \otimes V) \otimes (W \otimes Z) & & \\
& \nearrow \Phi & & \nwarrow \Phi & \\
((U \otimes V) \otimes W) \otimes Z & & & & U \otimes (V \otimes (W \otimes Z)) \\
& \searrow \Phi \otimes \text{Id} & & \nearrow \text{Id} \otimes \Phi & \\
& (U \otimes (V \otimes W)) \otimes Z & \xrightarrow{\Phi} & U \otimes ((V \otimes W) \otimes Z) &
\end{array}$$

Furthermore, Φ and Ψ must obey the *Hexagon Conditions* making the following diagrams commute.

$$\begin{array}{ccc}
& U \otimes (V \otimes W) & \\
\text{id} \otimes \Psi \swarrow & & \searrow \Phi^{-1} \\
U \otimes (W \otimes V) & & (U \otimes V) \otimes W \\
\Phi^{-1} \downarrow & & \downarrow \Psi \\
(U \otimes W) \otimes V & & W \otimes (U \otimes V) \\
\Psi \otimes \text{Id} \searrow & & \swarrow \Phi^{-1} \\
& (W \otimes U) \otimes V &
\end{array}
\quad
\begin{array}{ccc}
& (U \otimes V) \otimes W & \\
\Phi \swarrow & & \searrow \Psi \otimes \text{Id} \\
U \otimes (V \otimes W) & & (V \otimes U) \otimes W \\
\Psi \downarrow & & \downarrow \Phi \\
(V \otimes W) \otimes U & & V \otimes (U \otimes W) \\
\Phi \searrow & & \swarrow \text{Id} \otimes \Psi \\
& V \otimes (W \otimes U) &
\end{array}$$

Notice that we do not assume that $\Psi_{V,U} = \Psi_{U,V}^{-1}$ as it would if the tensor product were symmetric. This is related to the fact that \mathcal{R}^2 may not be $1 \otimes 1$, and this means that the group of functorial isomorphisms acting on a tensor product of H -modules $M_1 \otimes M_2 \otimes \cdots \otimes M_n$ is not the symmetric group S_n but rather the n -th *braid group* Br_n (or the *braid group on n -strings*) which admits an epimorphism $\pi_n: \text{Br}_n \rightarrow S_n$ with infinite kernel.

The group Br_n has a presentation with generators b_1, b_2, \dots, b_{n-1} and relations

$$b_i b_j = b_j b_i \quad (|i - j| \geq 2),$$

and the *Yang-Baxter equation*

$$b_i b_{i+1} b_i = b_{i+1} b_i b_{i+1}.$$

Similarly, S_n has a presentation with generators s_1, s_2, \dots, s_{n-1} and relations

$$s_i s_j = s_j s_i \quad (|i - j| \geq 2),$$

and

$$s_i s_{i+1} s_i = s_{i+1} s_i s_{i+1},$$

as well as

$$s_i^2 = 1.$$

Here $s_i = (i \ i + 1)$ and $\pi_n(b_i) = s_i$.

6.5. Some examples of quasitriangular structures

We take $\mathbb{k} = \mathbb{C}$ but any field containing a primitive n -th root of unity will do. Throughout we set $q = e^{2\pi i/n} \in \mathbb{C}$.

Let C_n be the cyclic group of order n and $\mathbb{C}C_n$ its group algebra. If $g \in C_n$ is a generator then $1, g, \dots, g^{n-1}$ is a basis of group-like elements for $\mathbb{C}C_n$. Of course $\chi(g) = g^{-1}$ and $\varepsilon(g) = 1$. This cocommutative Hopf algebra has the trivial quasitriangular structure with $\mathcal{R} = 1 \otimes 1$. But we can give it a different quasitriangular structure.

EXAMPLE 6.8. Take $(\mathbb{C}_q C_n, \mathcal{R})$ to be the Hopf algebra $\mathbb{C}_q C_n = \mathbb{C} C_n$ with

$$\mathcal{R} = \frac{1}{n} \sum_{a,b=0}^{n-1} q^{-ab} g^a \otimes g^b \in \mathbb{C}_q C_n \otimes \mathbb{C}_q C_n.$$

Then for $n > 2$ odd, $(\mathbb{C}_q C_n, \mathcal{R})$ is a ribbon quasitriangular Hopf algebra with

$$T(\mathcal{R})\mathcal{R} = \frac{1}{n} \sum_{a,b=0}^{n-1} q^{-ab} g^{2a} \otimes g^b$$

and

$$u = v = \nu = \frac{1}{n} \sum_{c=0}^{n-1} \vartheta(c) g^c,$$

where

$$\vartheta(c) = \sum_{d=0}^{n-1} q^{-(c+d)d} \in \mathbb{C}$$

is known as a C_n theta function.

See Majid [Maj02, Example 6.3] for the gory details.

The Lie algebra $\mathfrak{sl}_2 = \mathfrak{sl}_2(\mathbb{C})$ consists of all 2×2 traceless complex matrices,

$$\mathfrak{sl}_2 = \left\{ \begin{bmatrix} a & b \\ c & -a \end{bmatrix} : a, b, c \in \mathbb{C} \right\},$$

with the usual commutator as its Lie bracket. It is usual to take as a basis

$$h = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}, \quad e = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}, \quad f = \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}$$

with Lie brackets

$$[e, f] = h, \quad [h, e] = 2e, \quad [h, f] = -2f.$$

We can of course form the universal enveloping Hopf algebra $U(\mathfrak{sl}_2)$ which is primitively generated by the images of these basis elements.

Now we describe the quantum version.

EXAMPLE 6.9. Let $n > 2$ be odd and q a primitive complex n -th root of unity. We define the algebra $U_q(\mathfrak{sl}_2)$ by

$$U_q(\mathfrak{sl}_2) = \mathbb{C}\langle E, F, g \rangle / I$$

where $I \triangleleft \mathbb{C}\langle E, F, g \rangle$ is the two sided ideal generated by

$$g^n - 1, \quad E^n, \quad F^n, \quad [E, F] - \frac{g - g^{-1}}{q - q^{-1}}, \quad gEg^{-1} - q^2E, \quad gFg^{-1} - q^{-2}F.$$

This becomes a Hopf algebra with coalgebra structure given by

$$\begin{aligned} \psi(g) &= g \otimes g, \quad \psi(E) = E \otimes g + 1 \otimes E, \quad \psi(F) = F \otimes 1 + g^{-1} \otimes F, \\ \varepsilon(g) &= 1, \quad \varepsilon(E) = \varepsilon(F) = 0, \quad \chi(g) = g^{-1}, \quad \chi(E) = -Eg^{-1}, \quad \chi(F) = -gF. \end{aligned}$$

There is a quasitriangular structure

$$\mathcal{R} = \left(\frac{1}{n} \sum_{a,b=0}^{n-1} q^{-2ab} g^a \otimes g^b \right) \left(\sum_{c=0}^{n-1} \frac{(q - q^{-1})^c}{[c]_{q^{-2}}!} E^c \otimes F^c \right).$$

Then $(U_q(\mathfrak{sl}_2), \mathcal{R})$ is a quasitriangular Hopf algebra.

See Majid [Maj02, Example 6.4] for yet more gory details.

Bibliography

- [BG02] K. A. Brown and K. R. Goodearl, *Lectures on Algebraic Quantum Groups*, Advanced Courses in Mathematics, CRM Barcelona, Birkhäuser, 2002.
 - [CP21] P. Cartier and F. Patras, *Classical Hopf Algebras and their Applications*, Algebra and Applications, vol. 29, Springer, 2021. (pdf file downloadable through GU Library).
 - [CP94] V. Chari and A. Pressley, *A Guide to Quantum Groups*, Cambridge University Press, 1994.
 - [Kas23] R. Kashaev, *A Course on Hopf Algebras*, Universitext, Springer, 2023.
 - [Lan02] S. Lang, *Algebra*, 3rd ed., Graduate Texts in Mathematics, vol. 211, Springer-Verlag, 2002.
 - [Lor18] M. Lorenz, *A Tour of Representation Theory*, Graduate Studies in Mathematics, vol. 193, American Mathematical Society, 2018. (pdf file downloadable through GU Library).
 - [ML98] S. Mac Lane, *Categories for the Working Mathematician*, 2nd ed., Graduate Texts in Mathematics, vol. 5, Springer-Verlag, 1998.
 - [Maj95] S. Majid, *Foundations of Quantum Group Theory*, Cambridge University Press, 1995.
 - [Maj02] ———, *A Quantum Groups Primer*, London Mathematical Soc. Lect. Note Ser., vol. 292, Cambridge University Press, 2002.
 - [MP12] J. P. May and K. Ponto, *More Concise Algebraic Topology*, Chicago Lectures in Mathematics, University of Chicago Press, 2012. Localization, completion, and model categories.
 - [Mil17] J. S. Milne, *Algebraic Groups*, Cambridge Studies in Advanced Mathematics, vol. 170, Cambridge University Press, 2017. (pdf file downloadable through GU Library).
 - [MM65] J. W. Milnor and J. C. Moore, *On the structure of Hopf algebras*, Ann. of Math. (2) **81** (1965), 211–264.
 - [Mon93] S. Montgomery, *Hopf Algebras and their Actions on Rings*, CBMS Regional Conference Series in Mathematics, vol. 82, 1993.
 - [Rad12] D. E. Radford, *Hopf Algebras*, Series on Knots and Everything, vol. 49, World Scientific Publishing, 2012. (pdf file downloadable through GU Library).
 - [Rie16] E. Riehl, *Category Theory in Context*, Grad. Texts in Math., Dover Publications, 2016. (pdf file downloadable from <https://math.jhu.edu/~eriehl/context/>).
 - [Und11] R. G. Underwood, *An Introduction to Hopf Algebras*, Springer-Verlag, 2011. (pdf file downloadable through GU Library).
 - [Und15] ———, *Fundamentals of Hopf Algebras*, Springer, 2015. (pdf file downloadable through GU Library).
 - [Wat79] W. C. Waterhouse, *Introduction to Affine Group Schemes*, Grad. Texts in Math., vol. 66, Springer-Verlag, 1979.
-
- [AFS09] N. Andruskiewitsch and W. Ferrer Santos, *The beginnings of the theory of Hopf algebras*, Acta Appl. Math. **108** (2009), 3–17, available at [arXiv:2205.09541](https://arxiv.org/abs/2205.09541).
 - [BS62] W. Browder and E. Spanier, *H-spaces and duality*, Pacific J. Math. **12** (1962), 411–414.
 - [CE99] H. Cartan and S. Eilenberg, *Homological Algebra*, Princeton University Press, 1999. With an appendix by D. A. Buchsbaum; Reprint of the 1956 original.
 - [Jac79] N. Jacobson, *Lie Algebras*, Dover Publications, 1979. Republication of the 1962 original.
 - [KC02] V. Kac and P. Cheung, *Quantum Calculus*, Universitext, Springer-Verlag, 2002.
 - [Koc04] J. Kock, *Frobenius Algebras and 2D Topological Quantum Field Theories*, Lond. Math. Soc. Stud. Texts, vol. 59, Cambridge University Press, 2004.
 - [Lam99] T. Y. Lam, *Lectures on Modules and Rings*, Grad. Texts in Math., vol. 189, Springer-Verlag, 1999.
 - [LS69] R. G. Larson and M. E. Sweedler, *An associative orthogonal bilinear form for Hopf algebras*, Amer. J. Math. **91** (1969), 75–94.
 - [Lor11] M. Lorenz, *Some applications of Frobenius algebras to Hopf algebras*, Contemp. Math. **537** (2011), 269–289.

- [MR83] N. Metropolis and G-C. Rota, *Witt vectors and the algebra of necklaces*, Adv. in Math. **50** (1983), 95–125.