

Algebra III: Rings and Modules

Problem Sheet 4, Autumn Term 2022-23

John Nicholson

- Let R be a commutative ring and let $I \subseteq R$ be an ideal.
 - Prove that I is a free R -module if and only if I is principal and is generated by an element which is not a zero divisor. [Optional: Find a non-commutative ring where this is false.]
 - Deduce that a commutative ring R is a principal ideal domain if and only if every ideal $I \subseteq R$ is free as an R -module.
- Let R be a ring and let M be a free R -module. Give a proof or counterexample to each of the following statements:
 - Every spanning set for M over R contains a basis for M .
 - Every linearly independent subset of M over R can be extended to a basis for M .
- Let R be a non-trivial commutative ring. Prove that R is a field if and only if every finitely generated R -module is free. [Optional: Prove this is also equivalent to every R -module being free. You will need to use the axiom of choice.]
- Let R be a ring, let $S \subseteq R$ be a multiplicative submonoid and let $N \leq M$ be R -modules. Show that there is an isomorphism of $S^{-1}R$ -modules $S^{-1}(M/N) \cong S^{-1}M/S^{-1}N$.
- Let R be a ring, M a right R -module and N a left R -module. The tensor product $M \otimes_R N$ is defined to be the abelian group

$$M \otimes_R N = \mathbb{Z}[M \times N] / ((va, w) - (v, aw), (v, w) + (v', w) - (v + v', w), \\ (v, w) + (v, w') - (v, w + w') \mid a \in R, v, v' \in M, w, w' \in N).$$

For left R -modules M and N , let $\text{Hom}_R(M, N)$ denote the set of left R -module homomorphisms $f : M \rightarrow N$, which is an abelian group under pointwise addition.

From now on, let R be a commutative ring.

- Let M, N be left R -modules (which we can also view as right modules since R is commutative). Show that $M \otimes_R N$ is an R -module with action $a(v \otimes_R w) = av \otimes_R w$ for $a \in R, v \in M$ and $w \in N$.
- Let M, N be left R -modules. Show that $\text{Hom}_R(M, N)$ is an R -module, with action: for $a \in R$ and $\varphi : M \rightarrow N$, define $a \cdot \varphi : M \rightarrow N$ by $(a \cdot \varphi)(b) = a\varphi(b)$ for $b \in M$.
- Show that, if M, N , and T are all R -modules, then $\text{Hom}_R(M \otimes_R N, T)$ is identified with the set of R -bilinear maps $\varphi : M \times N \rightarrow T$, which means functions satisfying $\varphi(au, v) = a\varphi(u, v) = \varphi(u, av)$ and $\varphi(u + u', v) = \varphi(u, v) + \varphi(u', v)$ as well as $\varphi(u, v + v') = \varphi(u, v) + \varphi(u, v')$. Use this to give an alternative definition of tensor product.

6. Let R be a ring and let M be a left R -module. We say that R is a *ring with involution* (or a **-ring*) if R is equipped with a map $*$: $R \rightarrow R$ such that $(x + y)^* = x^* + y^*$, $(xy)^* = y^*x^*$, $1^* = 1$ and $(x^*)^* = x$ for all $x, y \in R$, i.e. $*$ is an anti-homomorphism and an involution.
- (i) Show that $M^* = \text{Hom}_R(M, R)$ is a right R -module with action: for $a \in R$ and $\varphi \in \text{Hom}_R(M, R)$, define $\varphi \cdot a : M \rightarrow R$ by $(\varphi \cdot a)(b) = \varphi(b) \cdot_R a$ for $b \in M$. This is known as the *dual module*.
 - (ii) Let R be a commutative ring. Show that R is a ring with involution. For a group G , show that $R[G]$ is a ring with involution.
 - (iii) Let R be a ring with involution. Show that any right R -module M can be viewed as a left R -module with action: for $a \in R$ and $m \in M$, define $x \cdot m = m \cdot_M x^*$. Use this to define a left R -module structure on $\text{Hom}_R(M, R)$. For left R -modules M and N , define a (sensible) left R -module structure on the tensor product of abelian groups $M \otimes_{\mathbb{Z}} N$. [Optional: How do these R -module structures compare to those defined in (5) in the commutative case?]
7. Let R be a ring and let M be an R -module and let $N \leq M$ be a submodule. Show that M is Noetherian if and only if N and M/N are Noetherian.
8. Let a, b be nonzero positive integers. Find the Smith normal form of the following matrices in their respective rings:

$$\begin{pmatrix} a & b \\ -b & a \end{pmatrix} \in M_2(\mathbb{Q}), \quad \begin{pmatrix} X^2 - 5X + 6 & X - 3 \\ (X - 2)^3 & X^2 - 5X + 6 \end{pmatrix} \in M_2(\mathbb{Q}[X]).$$

9. Let G be the abelian group given by generators a, b, c and the relations $6a + 10b = 0$, $6a + 15c = 0$, $10b + 15c = 0$ (i.e. G is the free abelian group generated by a, b, c quotiented by the subgroup $(6a + 10b, 6a + 15c, 10b + 15c)$). Determine the structure of G as a direct sum of cyclic groups.
10. A ring R has the *invariant basis number* property (IBN) if, for all positive integers m, n , $R^n \cong R^m$ as R -modules implies $m = n$.
- (i) For an ideal $I \subseteq R$ and an R -module M , we define an R -submodule $IM = \{am \in M : a \in I, m \in M\} \leq M$. Prove that M/IM is an R/I -module in a natural way.
 - (ii) Prove that non-zero commutative rings have IBN. You may assume that every non-zero commutative ring has a maximal ideal. [This is equivalent to the axiom of choice.]
 - (iii) Let S be a ring and M a free S -module with basis $\{x_i \mid i \geq 1\}$. Let $R = \text{End}_S(M)$. Prove that R does not have IBN. [Hint: Note that $M \cong M_{\text{even}} \oplus M_{\text{odd}}$ where M_{even} and M_{odd} are the submodules generated by x_i for i even and odd respectively. Use this to show that $R \cong R^2$ as R -modules.]
- +11. Let G be a finite group, let $N = \sum_{g \in G} g \in \mathbb{Z}[G]$ and let $r \in \mathbb{Z}$ be an integer with $(r, |G|) = 1$
- (i) Show that the ideal $(N, r) \subseteq \mathbb{Z}[G]$ is projective as a $\mathbb{Z}[G]$ -module.
 - (ii) Let $G = C_n$ be a finite cyclic group. Show that (N, r) is free as a $\mathbb{Z}[G]$ -module.
 - (iii) Let $G = Q_8$ be the quaternion group of order 8. Show that $(N, 3)$ is not free as a $\mathbb{Z}[G]$ -module. Is it stably free as a $\mathbb{Z}[G]$ -module?