

EMBEDDED SURFACES WITH INFINITE CYCLIC KNOT GROUP

ANTHONY CONWAY AND MARK POWELL

ABSTRACT. We study locally flat, compact, oriented surfaces in 4-manifolds whose exteriors have infinite cyclic fundamental group. We give algebraic topological criteria for two such surfaces, with the same genus g , to be related by an ambient homeomorphism, and further criteria that imply they are ambiently isotopic. Along the way, we provide a classification of a subset of the topological 4-manifolds with infinite cyclic fundamental group, and we apply our results to rim surgery.

1. INTRODUCTION

We study locally flat embeddings of compact, orientable surfaces in compact, oriented, simply-connected topological 4-manifolds, where the complement of the surface has infinite cyclic fundamental group. Extending the terminology for knotted spheres, we call this group the *knot group*, so we shall study knotted surfaces with knot group \mathbb{Z} , or \mathbb{Z} -surfaces.

We will present algebraic criteria for pairs of \mathbb{Z} -surfaces to be ambiently isotopic. As part of the proof we obtain an algebraic classification of a certain subset of the 4-manifolds with boundary and fundamental group \mathbb{Z} , simultaneously generalising work of Freedman-Quinn [FQ90] on the closed case and of Boyer [Boy86] on simply-connected 4-manifolds with nonempty boundary; see Section 1.7. We apply our results to show that, in simply-connected 4-manifolds, 1-twisted rim surgery on a surface with knot group \mathbb{Z} yields a topologically ambiently isotopic surface, extending results of Kim-Ruberman [KR08a] and Juhász-Miller-Zemke [JMZ20]; see Section 1.5.

1.1. Surfaces in S^4 and D^4 . We start with our main results on surfaces in S^4 and D^4 as the most important special cases, before going on to explain more general statements for surfaces in any simply-connected 4-manifold.

Theorem 1.1. *Any two locally flat, embedded, closed, orientable, genus $g \neq 1, 2$ surfaces $\Sigma_0, \Sigma_1 \subseteq S^4$ with $\pi_1(S^4 \setminus \Sigma_i) = \mathbb{Z}$ for $i = 0, 1$ are topologically ambiently isotopic.*

Theorem 1.2. *Let K be an Alexander polynomial one knot in S^3 . Any two locally flat, embedded, compact, orientable genus $g \neq 1, 2$ surfaces $\Sigma_0, \Sigma_1 \subseteq D^4$ with boundary K and $\pi_1(D^4 \setminus \Sigma_i) = \mathbb{Z}$ for $i = 0, 1$ are topologically ambiently isotopic rel. boundary.*

In a previous paper of ours, we proved Theorem 1.2 in the genus zero case [CP19, Theorem 1.2]. With regards to Theorem 1.1, a genus g surface $\Sigma \subseteq S^4$ is *unknotted* if it bounds a locally flat, embedded handlebody in S^4 . The *unknotting conjecture* for locally flat surfaces posits that a locally flat, embedded, closed, oriented surface $\Sigma \subseteq S^4$ is unknotted if and only if $\pi_1(S^4 \setminus \nu\Sigma) = \mathbb{Z}$. The forwards direction holds because any two embeddings of a handlebody in S^4 are ambiently isotopic, so an unknotted surface is ambiently isotopic to a standard embedding. For the reverse direction, Freedman and Quinn [FQ90, Theorem 11.7A] proved the $g = 0$ case of 2-knots, when $\Sigma \cong S^2$, while Hillman and Kawachi claimed it

for $g \geq 1$ [HK95]. Theorem 1.1 offers a new proof for surfaces of genus $g \geq 3$. We also give new proofs for the $g = 0$ cases, but it should be noted that this specialisation produces somewhat similar proofs to those of Freedman-Quinn and our previous work [CP19].

Our proof of Theorem 1.1 differs from that of Hillman-Kawauchi [HK95], who do not have any genus restrictions. In particular, in a key step in the proof, one considers a closed 4-manifold M built from the two surface exteriors, and shows that M is homeomorphic to $S^1 \times S^3 \#_{i=1}^{2g} S^2 \times S^2$. To see this, we control the $\mathbb{Z}[\mathbb{Z}]$ -valued intersection form of M , whereas [HK95, Proof of Lemma 3.1] just calculates the \mathbb{Z} -valued intersection form, and appeals to separate work of Kawauchi [Kaw94], in which it was claimed that every closed, spin 4-manifold with fundamental group \mathbb{Z} splits as $S^1 \times S^3 \# X$, where X is a closed, simply-connected 4-manifold. It would follow from this claim that computing the \mathbb{Z} -valued form suffices. A mistake in [Kaw94] was found by Hambleton-Teichner [HT97]. Although Kawauchi later updated his theorem [Kaw13] to include the hypothesis that the \mathbb{Z} -intersection form be indefinite, which is the case for the 4-manifolds arising in [HK95], the consensus in the community seems to be that an independent account is also desirable.

1.2. Surfaces in simply-connected 4-manifolds with boundary S^3 . To fix our terminology, throughout the article, by a *4-manifold* we shall always mean a compact, connected, oriented, topological 4-manifold with connected boundary. Surfaces will always be compact, connected, and orientable. We will consider locally flat, embedded, closed surfaces $\Sigma \subseteq X$ with X a closed 4-manifold, and locally flat properly embedded surfaces $\Sigma \subseteq N$, where N is a 4-manifold with boundary S^3 and $\partial\Sigma = K \subseteq S^3$ is a fixed knot. The exteriors of such surfaces will be denoted by X_Σ and N_Σ respectively.

Next we formulate our most general statement, for the nonempty boundary case, pairs of surfaces $\Sigma_i \subseteq N$, where $\partial\Sigma_i \subseteq \partial N = S^3$ is a fixed knot. Theorem 1.2 is an immediate consequence of Theorem 1.3 below, which we state after introducing the prerequisites. One quickly deduces consequences for closed surfaces in closed 4-manifolds X , in particular Theorem 1.1, by removing an unknotted (D^4, D^2) pair from (X, Σ) , as we will explain in Section 1.3.

Given a compact, oriented 4-manifold M with $\pi_1(M) = \mathbb{Z}$, set $\Lambda := \mathbb{Z}[t^{\pm 1}]$ and write $H_*(M; \Lambda)$ for the homology of the infinite cyclic cover \widetilde{M} of M , considered as a Λ -module. Taking signed intersections in \widetilde{M} endows the homology Λ -module $H_2(M; \Lambda)$ with a sesquilinear Hermitian intersection form

$$\lambda_M: H_2(M; \Lambda) \times H_2(M; \Lambda) \rightarrow \Lambda.$$

The adjoint of λ_M is the Λ -module homomorphism

$$\widehat{\lambda}_M: H_2(M; \Lambda) \rightarrow \text{Hom}_\Lambda(H_2(M; \Lambda), \Lambda) =: H_2(M; \Lambda)^*.$$

If two 4-manifolds M_0, M_1 with infinite cyclic fundamental group are orientation-preserving homeomorphic, then their Λ -intersection forms are isometric. That is, there exists an isomorphism $F: H_2(M_0; \Lambda) \xrightarrow{\cong} H_2(M_1; \Lambda)$ such that $\widehat{\lambda}_{M_0} = F^* \widehat{\lambda}_{M_1} F$. We write $F: \lambda_{M_0} \cong \lambda_{M_1}$ and call F an *isometry*.

Let Y be a compact oriented 3-manifold, and let $\varphi: \pi_1(Y) \rightarrow \mathbb{Z}$ be an epimorphism. Associated with this data, there is a Λ -module $H_1(Y; \Lambda)$, called the *Alexander module*, which is the first homology group of the infinite cyclic cover associated to $\ker(\varphi)$. If this module is torsion over $\Lambda = \mathbb{Z}[t^{\pm 1}]$, then it is endowed with a sesquilinear Hermitian *Blanchfield form*

$$\text{Bl}_Y: H_1(Y; \Lambda) \times H_1(Y; \Lambda) \rightarrow \mathbb{Q}(t)/\Lambda.$$

More details on this pairing appear in Subsection 3.1, but we note that Bl_Y should be thought of as the analogue of the linking pairing of a \mathbb{Q} -homology sphere on the level of infinite cyclic covers.

As we recall in Sections 2 and 3, if M_0, M_1 are 4-manifolds with $\pi_1(M_i) = \mathbb{Z}$ whose boundaries ∂M_i have torsion Alexander modules and if $\pi_1(\partial M_i) \rightarrow \pi_1(M_i)$ is surjective for $i = 0, 1$, then an isometry $F: \lambda_{M_0} \cong \lambda_{M_1}$ of the Λ -intersection forms induces an isometry of the Blanchfield forms of the boundary

$$\partial F: (H_1(\partial M_0; \Lambda), \text{Bl}_{\partial M_0}) \xrightarrow{\cong} (H_1(\partial M_1; \Lambda), \text{Bl}_{\partial M_1}).$$

Here is the construction: via universal coefficients, Poincaré-Lefschetz duality, and the long exact sequence of a pair, the composition

$$H_2(M_i; \Lambda)^* \cong H^2(M_i; \Lambda) \cong H_2(M_i, \partial M_i; \Lambda) \xrightarrow{\delta} H_1(\partial M_i; \Lambda)$$

induces an identification $\text{coker}(\widehat{\lambda}_{M_i}) \cong H_1(\partial M_i; \Lambda)$, and the map

$$F^{-*} := (F^*)^{-1}: H_2(M_0; \Lambda)^* \rightarrow H_2(M_1; \Lambda)^*$$

induces $\partial F: \text{coker}(\widehat{\lambda}_{M_0}) \rightarrow \text{coker}(\widehat{\lambda}_{M_1})$.

Now we focus on the case where $M_i = N_{\Sigma_i}$ are surface exteriors, with N a compact, simply-connected 4-manifold with boundary S^3 . The boundary ∂N_{Σ_i} is homeomorphic to

$$M_{K,g} := E_K \cup_{\partial} (\Sigma_{g,1} \times S^1),$$

where $E_K = S^3 \setminus \nu(K)$ is the knot exterior and $\Sigma_{g,1}$ denotes the (abstract) genus g surface with one boundary component. In Proposition 5.7, we show that every automorphism h of the Blanchfield pairing $\text{Bl}_{M_{K,g}}$ decomposes as $h_K \oplus h_{\Sigma}$, where h_K is an automorphism of $\text{Bl}_K := \text{Bl}_{E_K}$ and h_{Σ} is an automorphism of $\text{Bl}_{\Sigma_{g,1} \times S^1}$.

Let $f_K: E_K \rightarrow E_K$ be an orientation-preserving homeomorphism that is the identity on ∂E_K . Extend f_K via the identity on νK to an orientation-preserving self-homeomorphism of S^3 . The mapping class group of S^3 is trivial, so there is an isotopy $\Theta(f_K): S^3 \times [0, 1] \rightarrow S^3$ between the extension and the identity, such that $\Theta(f_K)|_{S^3 \times \{0\}} = \text{Id}$ and $\Theta(f_K)|_{E_K \times \{1\}} = f_K$.

Here is the central theorem of the paper on embedded surfaces with knot group \mathbb{Z} .

Theorem 1.3. *Let X be a closed, simply-connected 4-manifold, let $N = X \setminus \overset{\circ}{D}^4$ be a punctured X , and let $K \subseteq S^3 = \partial N$ be a knot. Let $\Sigma_0, \Sigma_1 \subseteq N$ be two locally flat, embedded, compact, oriented genus g surfaces with boundary K and $\pi_1(N_{\Sigma_i}) = \mathbb{Z}$ for $i = 0, 1$. Suppose there is an isometry $F: \lambda_{N_{\Sigma_0}} \cong \lambda_{N_{\Sigma_1}}$ and write $\partial F = h_K \oplus h_{\Sigma}$.*

- *If h_K is induced by an orientation-preserving homeomorphism $f_K: E_K \rightarrow E_K$ that is the identity on ∂E_K , then f_K extends to an orientation-preserving homeomorphism of pairs*

$$(N, \Sigma_0) \xrightarrow{\cong} (N, \Sigma_1)$$

inducing the given isometry $F: H_2(N_{\Sigma_0}; \Lambda) \cong H_2(N_{\Sigma_1}; \Lambda)$.

- *If in addition $N = D^4$, then for any choice of isotopy $\Theta(f_K)$, the surfaces Σ_0 and Σ_1 are topologically ambiently isotopic via an ambient isotopy of D^4 extending $\Theta(f_K)$.*

In particular note that if $h_K = \text{Id}$, then we can take $f_K = \text{Id}$ and Θ_{f_K} the constant isotopy, so that the homeomorphism of pairs in the first item can be assumed to fix the boundary pointwise, and the ambient isotopy in the second item can be assumed to be rel. boundary.

In general, $\lambda_{N_{\Sigma_0}}$ and $\lambda_{N_{\Sigma_1}}$ need not be isometric, even for $N = D^4$, as shown by examples due to Oba [Oba17].

The deduction of the last item uses Alexander's coning trick, which shows that every homeomorphism of D^4 that restricts to the identity on ∂D^4 is topologically isotopic to the identity. So in D^4 , a homeomorphism of pairs can be upgraded to a topological ambient isotopy. See Section 5.3 for details.

As explained in Section 1.5 below, we will apply Theorem 1.3 to 1-twisted rim surgery, a method which has been effective at producing exotic embeddings of surfaces. In Section 1.6, we also apply Theorem 1.3 to study Seifert surfaces that are pushed in to D^4 . Further applications can be obtained by finding classes of knots K for which every automorphism of the Blanchfield pairing is realised by a symmetry of the knot exterior E_K .

1.3. Ambient isotopy in closed 4-manifolds. For closed surfaces embedded in closed, simply-connected 4-manifolds, by applying the classification of self-homeomorphisms due to Kreck, Perron, and Quinn [Kre79, Per86, Qui86], we can potentially upgrade a homeomorphism of pairs obtained from Theorem 1.3 to an ambient isotopy. The theorem (see [Qui86] or [FQ90, Theorem 10.1]) is that two self-homeomorphisms of a closed, simply-connected 4-manifold are isotopic if and only if they induce the same self-isomorphism on second homology. An analogous classification of homeomorphisms for simply-connected 4-manifolds with boundary has not yet been proven.

Theorem 1.4. *Let X be a closed, simply-connected 4-manifold. Let $\Sigma_0, \Sigma_1 \subseteq X$ be two locally flat, embedded, closed, oriented genus g surfaces with $\pi_1(X_{\Sigma_i}) = \mathbb{Z}$ for $i = 0, 1$.*

- (1) *If the intersection forms $\lambda_{X_{\Sigma_0}}$ and $\lambda_{X_{\Sigma_1}}$ are isometric via an isometry F , then there is an orientation-preserving homeomorphism of pairs*

$$\Phi: (X, \Sigma_0) \xrightarrow{\cong} (X, \Sigma_1)$$

inducing the given isometry $\Phi_ = F: H_2(X_{\Sigma_0}; \Lambda) \cong H_2(X_{\Sigma_1}; \Lambda)$.*

- (2) *The isometry F induces an isometry $F_{\mathbb{Z}}: H_2(X) \rightarrow H_2(X)$ of the standard intersection form Q_X of X . The surfaces Σ_0 and Σ_1 are topologically ambiently isotopic if and only if $F_{\mathbb{Z}} = \text{Id}$.*

With regards to (2), we refer to Lemma 5.10 for details of how the isometry $F: \lambda_{\Sigma_0} \cong \lambda_{\Sigma_1}$ induces an isometry $F_{\mathbb{Z}}$ of the standard intersection form Q_X . Theorem 1.4 also has applications to rim surgery, as we explain in Section 1.5.

Theorem 1.4 (1) follows from Theorem 1.3. Here is a short outline; see Section 5.4 for details. After an isotopy, we assume Σ_0 and Σ_1 coincide on a disc D^2 . Remove an open neighbourhood of this common disc ($\mathring{D}^4, \mathring{D}^2$) from (X, Σ_i) to obtain a pair $(N, \tilde{\Sigma}_i)$, with $\partial N \cong S^3$ and $\partial \tilde{\Sigma}_i$ an unknot K . The exterior of Σ_i in X equals the exterior of $\tilde{\Sigma}_i$ in N , so the assumptions of Theorem 1.3 hold. Here, since the unknot has trivial Alexander module, $h_K = \text{Id}$ and so we take $f_K = \text{Id}$. Then Theorem 1.3 produces a homeomorphism of pairs $(N, \tilde{\Sigma}_0) \cong (N, \tilde{\Sigma}_1)$ rel. boundary, which we complete with the identity on the missing 4-ball to prove (1). Deducing (2) uses the classification of homeomorphisms from [Qui86], [FQ90, Theorem 10.1] mentioned above.

Remark 1.5. It is worth contrasting Theorem 1.4 with a result of Sunukjian. Indeed, [Sun15, Theorem 7.2] shows that closed surfaces $\Sigma_0, \Sigma_1 \subseteq X$ of the same genus with knot group \mathbb{Z} are topologically isotopic provided $b_2(X) \geq |\sigma(X)| + 2$. Thus, when X is "big enough", restrictions on the equivariant intersection form are not needed to establish isotopy.

1.4. Deducing Theorems 1.1 and 1.2 from Theorems 1.3 and 1.4. In the closed case, with $X = S^4$ if we have an isometry F of the intersection form of $X_{\Sigma_0} = S_{\Sigma_0}^4$ and $X_{\Sigma_1} = S_{\Sigma_1}^4$ then the map $F_{\mathbb{Z}}: H_2(S^4) \rightarrow H_2(S^4)$ is automatically the identity automorphism.

For the case with nonempty boundary, with $N = D^4$, Alexander polynomial one implies that $H_1(E_K; \Lambda) = 0$, so $h_K = \text{Id}$ automatically, and we may take $f_K: E_K \rightarrow E_K$ also to be the identity.

Therefore, in both cases, it suffices to find an isometry between the intersection forms of the surface exteriors. It is an open question whether this is true in general. For genus zero, i.e. for discs and spheres, this is automatically the case since the Λ -coefficient second homology of the surface exterior vanishes. Therefore the genus zero cases of Theorems 1.1 and 1.2, due to [FQ90, Theorem 11.7A] and [CP19, Theorem 1.2] respectively, follow from Theorems 1.3 and 1.4.

For genus at least three, our strategy to show that the intersection forms are isometric is as follows. As explained above, the exterior of a closed surface in S^4 can be considered as the exterior of a properly embedded surface in D^4 , so we discuss only the latter case.

By [BS16, Theorem 5] (see also Theorem A.1), Σ_0 and Σ_1 are stably equivalent, meaning that they become isotopic after adding some number of trivial tubes. Therefore the intersection forms of $\lambda_{D_{\Sigma_0}^4}$ and $\lambda_{D_{\Sigma_1}^4}$ satisfy $\lambda_{D_{\Sigma_0}^4} \oplus \mathcal{H}_2^{\oplus n} \cong \lambda_{D_{\Sigma_1}^4} \oplus \mathcal{H}_2^{\oplus n}$ for some $n \geq 0$, where

$$\mathcal{H}_2 := \begin{pmatrix} 0 & t-1 \\ t^{-1}-1 & 0 \end{pmatrix}.$$

Then provided g is at least three we are able to leverage algebraic cancellation results for hyperbolic forms from [Bas73] (see also [HT97, Kha17, MvdKV88, CS11]) to improve such a stable isometry to an isometry. While we refer to Subsection 7.1 for details, we record one of the aforementioned intermediate results as it might be of independent interest.

Proposition 1.6. *Let $\Sigma_0, \Sigma_1 \subseteq N = X \setminus \mathring{D}^4$ be two locally flat, properly embedded, compact, oriented genus g surfaces with boundary the same knot $K \subseteq S^3$ and $\pi_1(N_{\Sigma_i}) = \mathbb{Z}$ for $i = 0, 1$. There exists an integer $n \geq 0$ and an isometry*

$$\lambda_{N_{\Sigma_0}} \oplus \mathcal{H}_2^{\oplus n} \cong \lambda_{N_{\Sigma_1}} \oplus \mathcal{H}_2^{\oplus n}.$$

Whereas Theorems 1.1 and 1.2 hold for $g \neq 1, 2$, Proposition 1.6 leads to some results for arbitrary g . Indeed, Corollary 6.6 shows that for any genus g surface $\Sigma \subseteq N = X \setminus \mathring{D}^4$ with knot group \mathbb{Z} and $\partial\Sigma$ an Alexander polynomial one knot, $\lambda_{N_{\Sigma}} \oplus \mathcal{H}_2^{\oplus n} \cong Q_X \oplus \mathcal{H}_2^{\oplus (g+n)}$ for some $n \geq 0$, where Q_X denotes the standard intersection form of X . The same result holds for closed surfaces $\Sigma \subseteq X$ with knot group \mathbb{Z} .

1.5. Application to rim surgery. Rim surgery is an effective way to produce smoothly inequivalent embeddings of surfaces [FS97, Kim06, KR08a, Mar13, JMZ20]. Given a locally flat embedded closed oriented surface Σ in a closed 4-manifold X , a knot $J \subseteq S^3$ and simple closed curve $\alpha \subseteq \Sigma$, n -roll m -twist rim surgery outputs another locally flat embedded closed oriented surface $\Sigma_n^m(\alpha, J) \subseteq X$. The case where $m = 0, n = 0$ was first introduced by Fintushel-Stern [FS97], while m -twisted rim surgery was introduced by Kim [Kim06], and n -roll m -twist rim surgery first appeared in [KR08b].

Kim and Ruberman showed that if X is a simply-connected 4-manifold, $\pi_1(X_{\Sigma}) = \mathbb{Z}_d$ is a finite cyclic group and $(m, d) = 1$, then $\Sigma_n^m(\alpha, J)$ and Σ are topologically isotopic [KR08b, Theorem 1.3 and Proposition 3.1]. In order to extend this to infinite cyclic fundamental

groups, we consider n -roll 1-twist rim surgery, and set $\Sigma_n(\alpha, J) := \Sigma_n^1(\alpha, J)$. In this case if $\pi_1(X_\Sigma) = \mathbb{Z}$, then also $\pi_1(X_{\Sigma_n(\alpha, J)}) = \mathbb{Z}$. Our first result on rim surgery then reads as follows.

Theorem 1.7. *Let X be a closed, simply-connected 4-manifold, let $\Sigma \subseteq X$ be a locally flat, embedded, orientable surface with knot group \mathbb{Z} , let $\alpha \subseteq \Sigma$ be a simple closed curve, let J be a knot, and let $n \in \mathbb{Z}$. Then the surfaces Σ and $\Sigma_n(\alpha, J)$ are topologically ambiently isotopic.*

For good fundamental groups $\pi_1(X \setminus \Sigma)$ and J a slice knot, Kim and Ruberman showed in [KR08a, Theorem 4.5] that there is a homeomorphism of pairs $(X, \Sigma) \cong (X, \Sigma_0(\alpha, J))$, so for $\pi_1(X \setminus \Sigma) \cong \mathbb{Z}$ our result is an extension.

Rim surgery is also defined for a properly embedded surface in a 4-manifold N with boundary. Juhász, Miller, and Zemke showed that if $\Sigma \subseteq N$ is a properly embedded locally flat oriented surface, and if α bounds a locally flat disc in N_Σ , then $\Sigma_0(\alpha, J)$ is topologically isotopic to Σ [JMZ20, Corollary 2.7]. In the case where Σ has knot group \mathbb{Z} , we generalise this result as follows.

Theorem 1.8. *Let X be a closed, simply-connected 4-manifold, and let $N := X \setminus \mathring{D}^4$ be a punctured X . Let $\Sigma \subseteq N$ be a locally flat, properly embedded, orientable surface with knot group \mathbb{Z} , let $\alpha \subseteq \Sigma$ be a simple closed curve, let J be a knot, and let $n \in \mathbb{Z}$ be an integer. There is a rel. boundary orientation-preserving homeomorphism of pairs*

$$(N, \Sigma) \xrightarrow{\cong} (N, \Sigma_n(\alpha, J)).$$

If $N = D^4$ the surfaces Σ and $\Sigma_n(\alpha, J)$ are topologically ambiently isotopic rel. boundary.

Note that in both of the previous two results, we have no restrictions on the genera of the surfaces. This is because we show that the Λ -intersection forms of $N_{\Sigma_n(\alpha, J)}$ and N_Σ coincide, allowing us to apply the first item of Theorem 1.3, for which there is no genus restriction.

For the case $N = D^4$, Theorem 1.8 means that the construction of exotic surfaces in [JMZ20] applies more generally than realised in that article, because the condition that α bounds a locally flat embedded disc in N_Σ is not needed.

1.6. Application to pushed-in Seifert surfaces. Pushing a Seifert surface for a knot K into D^4 yields a surface with knot group \mathbb{Z} . It is then intriguing to wonder whether any two Seifert surfaces of the same genus for a knot K become isotopic once pushed into D^4 ; see for instance [Kir97, Problem 1.20 (C)] and [Liv82, Section 6]. In Theorem 7.11, relying on Theorem 1.3, we show that this is the case for Alexander polynomial one knots.

Theorem 1.9. *If $F_0, F_1 \subseteq D^4$ are genus g pushed-in Seifert surfaces for an Alexander polynomial one knot K , then they are topologically ambiently isotopic rel. boundary.*

This result provides another setting where we are able to obtain results for $g = 1, 2$. In particular, it rules out the most naive potential counter examples to the conjecture that Theorem 1.2 holds for \mathbb{Z} -surfaces of any genus.

1.7. A classification result for topological 4-manifolds. Several of the steps from the proof of Theorem 1.3 fit in a classification scheme that applies to a wider class of 4-manifolds than surface exteriors. We state the resulting theorem, which is analogous to Boyer's classification for the simply-connected case [Boy86].

We say that a compact, oriented, topological 4-manifold M with nonempty connected boundary has *ribbon boundary* if the inclusion induced map $\iota: \pi_1(\partial M) \rightarrow \pi_1(M)$ is surjective. For example, in simply-connected 4-manifolds, (connected) surface exteriors with infinite cyclic fundamental group have ribbon boundaries. Set $\Lambda := \mathbb{Z}[t^{\pm 1}]$. If we assume that $\pi_1(M) = \mathbb{Z}$, then we have infinite cyclic covers $\widetilde{M} \rightarrow M$ and $\partial \widetilde{M} \rightarrow \partial M$ and Λ -modules $H_*(M; \Lambda) = H_*(\widetilde{M})$ and $H_*(\partial M; \Lambda) = H_*(\partial \widetilde{M})$.

Let M_0, M_1 be two 4-manifolds with infinite cyclic fundamental group whose boundaries are ribbon and have Λ -torsion Alexander modules. Fix an orientation-preserving homeomorphism $f: \partial M_0 \rightarrow \partial M_1$ that intertwines the inclusion induced maps $\varphi_i: \pi_1(\partial M_i) \rightarrow \mathbb{Z}$, and an isometry $F: (H_2(M_0; \Lambda), \lambda_{M_0}) \rightarrow (H_2(M_1; \Lambda), \lambda_{M_1})$ of the Λ -valued intersection forms λ_{M_0} and λ_{M_1} . As described in Subsection 3.4 below, f and F respectively induce isometries \widetilde{f}_* and ∂F of the boundary Blanchfield forms:

$$\widetilde{f}_*, \partial F: (H_1(\partial M_0; \Lambda), \text{Bl}_{\partial M_0}) \rightarrow (H_1(\partial M_1; \Lambda), \text{Bl}_{\partial M_1}).$$

We call (f, F) a *compatible pair* if f and F induce the same isometry. For conciseness, we write

$$\text{Homeo}_{\varphi}^+(\partial M_0, \partial M_1) := \{f \in \text{Homeo}^+(\partial M_0, \partial M_1) \mid \varphi_1 \circ f_* = \varphi_0\}$$

for the set of orientation-preserving homeomorphisms of the boundaries that intertwine the maps to \mathbb{Z} .

Recall that the Kirby-Siebenmann invariant $\text{ks}(M_i) \in \mathbb{Z}/2$ is the unique obstruction for the stable tangent bundle $M_i \rightarrow \text{BTOP}$ to lift to $M_i \rightarrow \text{BPL}$, or equivalently for M_i to be smoothable after adding copies of $S^2 \times S^2$ [FQ90, Theorem 8.6], [FNOP19, Section 8]. This is relevant in the next theorem for the non-spin case; note that for 4-manifolds with fundamental group \mathbb{Z} , whether or not they are spin is determined by the intersection pairing.

Theorem 1.10. *Let M_0 and M_1 be two compact, oriented 4-manifolds with infinite cyclic fundamental group, whose boundaries are ribbon and have Λ -torsion Alexander modules. Let $f \in \text{Homeo}_{\varphi}^+(\partial M_0, \partial M_1)$ be a homeomorphism and let $F: \lambda_{M_0} \cong \lambda_{M_1}$ be an isometry. If M_0 and M_1 are not spin, then assume that the Kirby-Siebenmann invariants satisfy $\text{ks}(M_0) = \text{ks}(M_1) \in \mathbb{Z}/2$. Then the following assertions are equivalent:*

- (1) *the pair (f, F) is compatible;*
- (2) *the homeomorphism f extends to an orientation-preserving homeomorphism*

$$\Phi: M_0 \xrightarrow{\cong} M_1$$

inducing the given isometry $F: H_2(M_0; \Lambda) \cong H_2(M_1; \Lambda)$.

Remark 1.11. Theorem 1.10 should be compared with both Boyer's classification of simply-connected 4-manifolds with boundary [Boy86] and with Freedman-Quinn's classification of closed 4-manifolds with infinite cyclic fundamental group [FQ90]. Boyer uses a notion similar to our compatible pairs that he calls *morphisms*. In a nutshell, given simply-connected 4-manifolds M_0, M_1 with rational homology spheres as their boundaries, Boyer shows that a morphism (f, F) can be extended to a homeomorphism $M_0 \rightarrow M_1$ [Boy86, Theorem 0.7 and Proposition 0.8]. Note that Boyer's methods differ from ours, as he does not use the union of forms. On the other hand, Freedman-Quinn show that closed 4-manifolds with infinite cyclic fundamental groups are classified by their Λ -intersection form and their Kirby-Siebenmann invariant [FQ90, Theorem 10.7A], [SW00], [HKT09].

A major step in the proof of Theorem 1.10 is the following intermediate result that might be of independent interest; a more detailed statement and a proof can be found in Theorem 3.12.

Theorem 1.12. *Let M_0 and M_1 be two compact oriented 4-manifolds with infinite cyclic fundamental group, whose boundaries are ribbon and have Λ -torsion Alexander modules. Let (f, F) be a compatible pair. If M_0 and M_1 are not spin, then assume that the Kirby-Siebenmann invariants satisfy $\text{ks}(M_0) = \text{ks}(M_1) \in \mathbb{Z}/2$. Then there is an orientation-preserving homeomorphism*

$$M_0 \cup_f -M_1 \cong S^1 \times S^3 \#_{i=1}^a S^2 \times S^2 \#_{j=1}^b S^2 \tilde{\times} S^2$$

for some a, b with $a + b = b_2(M_0)$. If M_0 and M_1 are spin then $b = 0$.

Remark 1.13. We compare Theorem 1.12 with a particular case of a result due to Hambleton-Teichner [HT97]. Hambleton and Teichner show that if M is a closed, oriented, topological 4-manifold with infinite cyclic fundamental group and with $b_2(M) - |\sigma(M)| \geq 6$, then M is homeomorphic to the connected sum of $S^1 \times S^3$ with a unique closed, simply-connected 4-manifold [HT97, Corollary 3]. In most cases, this result is stronger than Theorem 1.12. However, there are some situations (such as the union of two genus one surfaces exteriors $X_{\Sigma_0}, X_{\Sigma_1} \subseteq S^4$ with $\pi_1(X_{\Sigma_i}) = \mathbb{Z}$ for $i = 0, 1$) where Theorem 1.12 applies while [HT97, Corollary 3] does not. Finally, note that Sunukjian has used these results to study closed surfaces $\Sigma \subseteq X$ with $\pi_1(X \setminus \nu\Sigma) = \mathbb{Z}$ in closed 4-manifolds that satisfy $b_2(X) \geq |\sigma(X)| + 6$ [Sun15, Theorem 7.2].

1.8. Isometries of the Blanchfield form. Theorem 1.10 provides a criterion for extending a homeomorphism $f: \partial M_0 \rightarrow \partial M_1$ to a homeomorphism $M_0 \rightarrow M_1$: one must fit f into a compatible pair. However, in practice, finding compatible pairs is difficult and so we provide some sufficient conditions for their existence.

Recall that if V is a 4-manifold with $\pi_1(V) = \mathbb{Z}$ whose boundary $Y = \partial V$ is ribbon and has torsion Alexander module, then an isometry F of the Λ -intersection form λ_V induces an isometry

$$\partial F: (H_1(Y; \Lambda), \text{Bl}_Y) \xrightarrow{\cong} (H_1(Y; \Lambda), \text{Bl}_Y).$$

We write $\text{Aut}(\lambda_V)$ and $\text{Aut}(\text{Bl}_Y)$ for the groups of isometries of λ_V and Bl_Y , and note that there is a left action of $\text{Aut}(\lambda_V)$ on $\text{Aut}(\text{Bl}_Y)$ given by $F \cdot h = h \circ \partial F^{-1}$. We will be interested in a quotient of the orbit set

$$\text{Aut}(\text{Bl}_Y) / \text{Aut}(\lambda_V).$$

In order to find compatible pairs, we need to know when an isometry of the Blanchfield form is induced by a homeomorphism. As we have already alluded to, we note in Proposition 3.7 that any homeomorphism $f \in \text{Homeo}_\varphi^+(Y)$ induces a Λ -isometry \tilde{f}_* of Bl_Y by lifting f to the infinite cyclic covers. Here, $\text{Homeo}_\varphi^+(Y) := \text{Homeo}_\varphi^+(Y, Y)$ denotes the set of orientation-preserving self-homeomorphisms of Y that intertwine the map $\pi_1(Y) \rightarrow \mathbb{Z}$. The assignment $(f, F) \cdot h = \tilde{f}_* \circ h \circ \partial F^{-1}$ then gives rise to a left action

$$\text{Homeo}_\varphi^+(Y) \times \text{Aut}(\lambda_V) \curvearrowright \text{Aut}(\text{Bl}_Y).$$

We return to our 4-manifolds M_0, M_1 whose boundaries are ribbon and have torsion Alexander module. If M_0 and M_1 are orientation-preserving homeomorphic, then a compatible pair exists; recall Theorem 1.10. Assuming that we are given a homeomorphism $f' \in \text{Homeo}_\varphi^+(\partial M_0, \partial M_1)$ and an isometry $F': \lambda_0 \cong \lambda_1$, Proposition 3.14 notes that (f', F') gives rise to a compatible

pair (f, F) if the composition $f'_* \circ \partial F'^{-1}$ is trivial in $\text{Aut}(\text{Bl}_{\partial M_1}) / \text{Homeo}_\varphi^+(\partial M_1) \times \text{Aut}(\lambda_{M_1})$. Using this result, we obtain the following consequence of Theorem 1.10.

Theorem 1.14. *Let M_0 and M_1 be two compact, oriented 4-manifolds with infinite cyclic fundamental group whose boundaries are ribbon and have Λ -torsion Alexander modules. Assume that M_0 and M_1 have orientation-preserving homeomorphic boundaries, via a homeomorphism that intertwines the maps to \mathbb{Z} , and isometric intersection forms. If the orbit set*

$$\text{Aut}(\text{Bl}_{\partial M_1}) / \text{Homeo}_\varphi^+(\partial M_1) \times \text{Aut}(\lambda_{M_1})$$

is trivial, then there is an orientation-preserving homeomorphism $M_0 \cong M_1$.

Theorem 1.14 is helpful to summarise a programme to prove that two 4-manifolds M_0, M_1 (with infinite cyclic fundamental group and identical ribbon boundaries with torsion Alexander modules) are homeomorphic:

- (1) decide whether the Λ -intersection forms λ_{M_0} and λ_{M_1} are isometric;
- (2) show that the orbit set $\text{Aut}(\text{Bl}_{\partial M_1}) / \text{Homeo}_\varphi^+(\partial M_1) \times \text{Aut}(\lambda_{M_1})$ is trivial.

To illustrate this paradigm, we explain how Theorem 1.3 can be applied to classify \mathbb{Z} -surfaces for some knots purely in terms of their intersection pairings. For any knot $K \subseteq S^3$, multiplication by a monomial $\pm t^k$ gives rise to an automorphism of its Blanchfield pairing Bl_{E_K} on the exterior of the knot. For some knots K , these are the only automorphisms, or at least they form a finite index subgroup, and in this case we obtain a simpler classification of \mathbb{Z} -surfaces, i.e. locally flat Σ in N with $\pi_1(N_\Sigma) \cong \mathbb{Z}$, with boundary K .

Corollary 1.15. *Let X be a closed, simply-connected 4-manifold, and let $N := X \setminus \mathring{D}^4$ be a punctured X . Let Σ_0 and Σ_1 be two \mathbb{Z} -surfaces in N with the same genus and boundary $\partial \Sigma_0 = \partial \Sigma_1 = K \subseteq S^3$. Suppose that $\text{Aut}(\text{Bl}_{E_K}) \subseteq \{\pm t^k \mid k \in \mathbb{Z}\}$.*

- (i) *There is a rel. boundary homeomorphism of pairs $(N, \Sigma_0) \cong (N, \Sigma_1)$ if and only if there is an isometry $\lambda_{N_{\Sigma_0}} \cong \lambda_{N_{\Sigma_1}}$.*
- (ii) *If $N = D^4$ then Σ_0 and Σ_1 are ambiently isotopic rel. boundary if and only if there is an isometry $\lambda_{D_{\Sigma_0}^4} \cong \lambda_{D_{\Sigma_1}^4}$.*

Proof. Let $F: \lambda_{N_{\Sigma_0}} \cong \lambda_{N_{\Sigma_1}}$ be an isometry. Consider the isomorphism $g := \partial F|_{H_1(E_K; \Lambda)}$ induced on the Alexander module of K . It is an automorphism of the Blanchfield form and therefore by the hypothesis on $\text{Aut}(\text{Bl}_{E_K})$, g is multiplication by $\pm t^k$ for some $k \in \mathbb{Z}$. Precompose F with multiplication by $\pm t^{-k}$, to obtain a new isometry inducing Id on $H_1(E_K; \Lambda)$. Then apply Theorem 1.3 with $f_K = \text{Id}$. \square

Remark 1.16. If the Alexander module $H_1(E_K; \Lambda)$ is cyclic with order Δ_K , then the automorphisms of the Blanchfield pairing can be computed directly as follows. Suppose that $\text{Bl}_K(1, 1) = b/\Delta_K \in \mathbb{Q}(t)/\Lambda$ for some $b \in \Lambda$. Then $\text{Aut}(\text{Bl}_{E_K}) = \{[p] \in \Lambda/\Delta\Lambda \mid [p \cdot \bar{p} \cdot b] = [b] \in \Lambda/\Delta\Lambda\}$. In particular for $K = \pm T_{2,3}$ a trefoil, it is not too hard to compute that $\text{Aut}(\text{Bl}_{E_K}) = \{\pm t^k \mid k = -1, 0, 1\}$, so that Corollary 1.15 applies to classify all \mathbb{Z} -surfaces with boundary a trefoil in terms of the isometry type of the intersection pairing.

Organisation. In Section 2, we review some notions on linking forms and define the union of Hermitian forms over Λ . In Section 3, we show how this algebraic union can be used to express the intersection form of a union of two 4-manifolds; we also prove Theorem 1.12. In Section 4, we prove Theorem 1.10, our partial classification result for 4-manifolds with infinite cyclic fundamental group. In Section 5, we prove Theorems 1.3 and 1.4, our main results

on properly embedded surfaces in simply-connected 4-manifolds. In Section 6, we discuss the equivariant intersection forms for surfaces with knot group \mathbb{Z} , before focusing on surfaces in S^4 and D^4 in Section 7, where we prove Theorems 1.1 and 1.2. In Section 8, we prove Theorems 1.7 and 1.8 on rim surgery. In Appendix A, we adapt the work of Baykur-Sunukjian to properly embedded surfaces in 4-manifolds with boundary.

Acknowledgments. We thank Peter Teichner for pointing out a mistake in a previous version of this work and for stressing the importance of $(-t)$ -Hermitian forms in the proof of Theorem 7.4. We are indebted to anonymous referees for several very useful suggestions which helped us to significantly improve the exposition and simplify some proofs. We thank Lisa Piccirillo for helpful discussions, and İnanç Baykur and Nathan Sunukjian for discussions on their stabilisation theorem, which helped us when writing the appendix. Part of this work was completed while the authors were visitors at the Max Planck Institute for Mathematics in Bonn.

Conventions.

- (1) From now on, all manifolds are assumed to be compact, connected, based and oriented; if a manifold has a nonempty boundary, then the basepoint is assumed to be in the boundary. We work in the topological category with locally flat embeddings unless otherwise stated.
- (2) X will always denote a simply-connected, closed 4-manifold and we will write $N := X \setminus \overset{\circ}{D}^4$ for the complement of an open ball $\overset{\circ}{D}^4 \subseteq X$.
- (3) A \mathbb{Z} -surface Σ will always refer to a compact, connected, orientable, locally flat, embedded surface in a 4-manifold whose knot group is \mathbb{Z} ; a surface $\Sigma \subseteq X$ is understood to be closed, while a surface $\Sigma \subseteq N$ is understood to be properly embedded.
- (4) If P is manifold and $Q \subseteq P$ is a submanifold with tubular neighborhood $\nu Q \subseteq P$, then $P_Q := P \setminus \nu Q$ will always denote the exterior of Q in P . The only exception to this is that the exterior of a knot K in S^3 will be denoted E_K instead of S_K^3 .
- (5) Throughout the article, we set $\Lambda := \mathbb{Z}[t^{\pm 1}]$ for the ring of Laurent polynomials and $Q := \mathbb{Q}(t)$ for its field of fractions.
- (6) We write $\bar{p} \mapsto p$ for the involution on Λ induced by $t \mapsto t^{-1}$. Given a Λ -module H , we write \bar{H} for the Λ -module whose underlying abelian group is H but with module structure given by $\bar{p} \cdot h = \overline{p \cdot h}$ for $h \in H$ and $p \in \Lambda$.
- (7) We write $H^* := \overline{\text{Hom}}_{\Lambda}(H, \Lambda)$.
- (8) For any ring R , elements of R^n are considered as column vectors.
- (9) Let $f: X \rightarrow Y$ be a morphism in a category \mathcal{C} , and let $F: \mathcal{C} \rightarrow \mathcal{D}$ be a contravariant functor. Assume that $F(f)$ is invertible. We shall often write the induced morphism in \mathcal{D} as $f^* = F(f)$, and denote its inverse by $f^{-*} := (f^*)^{-1} = F(f)^{-1}$.

2. THE UNION OF FORMS ALONG ISOMETRIES OF THEIR BOUNDARY LINKING FORMS

We develop the theory required for computing the intersection form of a closed 4-manifold obtained as the union of two compact 4-manifolds along their common boundary. Here is a summary of this section. In Subsection 2.1, we review the boundary linking form of a Hermitian form. In Subsection 2.2, we discuss isometries of Hermitian forms and linking forms. In Subsection 2.3, we study the union of two Hermitian forms over $\Lambda := \mathbb{Z}[t^{\pm 1}]$. In Subsection 2.4, we provide a condition for this union to admit a metaboliser.

2.1. The boundary of Hermitian forms and their isometries. We fix our terminology on Hermitian forms, linking forms and their boundaries. References include [Ran81, Section 3.4] and [CS11, Section 6].

A *Hermitian form* over Λ is a pair (H, λ) , where H is a free Λ -module and $\lambda: H \times H \rightarrow \Lambda$ is a sesquilinear Hermitian pairing. Here by sesquilinear, we mean that $\lambda(px, qy) = p\lambda(x, y)\bar{q}$ for all $x, y \in H$ and all $p, q \in \Lambda$. By Hermitian, we mean that $\lambda(y, x) = \lambda(x, y)$ for all $x, y \in H$. The *adjoint* of λ is the Λ -linear map $\hat{\lambda}: H \rightarrow \overline{\text{Hom}}_{\Lambda}(H, \Lambda) =: H^*$ such that $\hat{\lambda}(y)(x) = \lambda(x, y)$. A Hermitian form is *nondegenerate* if its adjoint is injective and *nonsingular* if its adjoint is an isomorphism. The *standard hyperbolic form* is

$$H^+(\Lambda) := \left(\Lambda \oplus \Lambda, \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \right).$$

Remark 2.1. If (H, λ) is a nondegenerate Hermitian form, then $\text{coker}(\hat{\lambda})$ is a torsion Λ -module. This can be seen by tensoring the following exact sequence with the field of fractions $Q := \mathbb{Q}(t)$ of Λ :

$$0 \rightarrow H \xrightarrow{\hat{\lambda}} H^* \rightarrow \text{coker}(\hat{\lambda}) \rightarrow 0.$$

A *linking form* (T, β) over Λ consists of a torsion Λ -module T together with a sesquilinear Hermitian form $\beta: T \times T \rightarrow Q/\Lambda$. The *adjoint* of β is the Λ -linear map $\hat{\beta}: T \rightarrow \overline{\text{Hom}}_{\Lambda}(T, Q/\Lambda)$ such that $\hat{\beta}(y)(x) = \beta(x, y)$. A linking form is *nondegenerate* if its adjoint is injective and *nonsingular* if its adjoint is an isomorphism.

Definition 2.2. The *boundary linking form* of a nondegenerate Hermitian form (H, λ) over Λ is the linking form $(\text{coker}(\hat{\lambda}), \partial\lambda)$, where $\partial\lambda$ is defined as

$$\begin{aligned} \partial\lambda: \text{coker}(\hat{\lambda}) \times \text{coker}(\hat{\lambda}) &\rightarrow Q/\Lambda \\ ([x], [y]) &\mapsto \frac{1}{s}(y(z)), \end{aligned}$$

where, since $\text{coker}(\hat{\lambda})$ is Λ -torsion, there exists an $s \in \Lambda$ and an $z \in H$ such that $sx = \hat{\lambda}(z)$.

It is not difficult to show that $\partial\lambda$ is independent of the choices involved, is sesquilinear and Hermitian. We conclude with two remarks that we will use throughout this section.

Remark 2.3. Let (H, λ) be a Hermitian form, and set $H_Q := H \otimes_{\Lambda} Q$. Since H is free, we can identify $\overline{\text{Hom}}_{\Lambda}(H, \Lambda) \otimes_{\Lambda} Q$ with $H_Q^* := \overline{\text{Hom}}_Q(H_Q, Q)$. As stated in Remark 2.1, if (H, λ) is nondegenerate, then $\text{coker}(\hat{\lambda})$ is a torsion Λ -module and therefore $\hat{\lambda}_Q := \hat{\lambda} \otimes_{\Lambda} \text{Id}_Q: H_Q \rightarrow H_Q^*$ is a nonsingular Hermitian form over Q .

Using this remark, we describe an equivalent definition of the boundary linking form.

Remark 2.4. The boundary linking form of a nondegenerate Hermitian form (H, λ) can be described as $\partial\lambda([x], [y]) = y(\hat{\lambda}_Q^{-1}(x))$ for $[x], [y] \in \text{coker}(\hat{\lambda})$. Choose a basis $\mathbf{b} = (e_i)_{i=1}^n$ for H and endow $\overline{\text{Hom}}_{\Lambda}(H, \Lambda)$ with the dual basis \mathbf{b}^* . If $A_{ij} = \lambda(e_i, e_j)$ is a Hermitian matrix representing λ , then \bar{A} is a matrix for $\hat{\lambda}$ and $\partial\lambda([x], [y]) = y(\hat{\lambda}_Q^{-1}(x)) = (\bar{A}^{-1}x)^T \bar{y} = x^T \bar{A}^{-1} \bar{y}$. For future reference, note that since A is Hermitian, we have $\bar{A} = A^T$.

2.2. Isometries of forms. We discuss isometries of Hermitian forms and linking forms. References include [Ran81, Section 3.4] and [CS11, Section 6].

Let (H_0, λ_0) and (H_1, λ_1) be Hermitian forms over Λ . A Λ -linear isomorphism $F: H_0 \rightarrow H_1$ is an *isometry* if $\lambda_1(F(x), F(y)) = \lambda_0(x, y)$ for all $x, y \in H_0$. Let $\text{Iso}(\lambda_0, \lambda_1)$ denote the set of isometries between the Hermitian forms (H_0, λ_0) and (H_1, λ_1) , and let $\text{Aut}(\lambda) := \text{Iso}(\lambda, \lambda)$ denote the group of self-isometries of a Hermitian form (H, λ) .

We make the analogous definitions for linking forms. Let (T_0, β_0) and (T_1, β_1) be two linking forms over Λ . A Λ -linear isomorphism $h: T_0 \rightarrow T_1$ is an *isometry* if $\beta_1(h(x), h(y)) = \beta_0(x, y)$ for all $x, y \in T_0$. We let $\text{Iso}(\beta_0, \beta_1)$ denote the set of isometries between the linking forms (T_0, β_0) and (T_1, β_1) , and let $\text{Aut}(\beta) := \text{Iso}(\beta, \beta)$ denote the group of self-isometries of a linking form (T, β) .

An isomorphism $F: H_0 \rightarrow H_1$ induces an isomorphism $F^{-*} := (F^*)^{-1}: H_0^* \rightarrow H_1^*$. If additionally, the isomorphism $F \in \text{Iso}(\lambda_0, \lambda_1)$ is an isometry, then F^{-*} descends to an isomorphism

$$\partial F := F^{-*}: \text{coker}(\widehat{\lambda}_0) \rightarrow \text{coker}(\widehat{\lambda}_1).$$

For later use, note that $F: (H_0, \lambda_0) \rightarrow (H_1, \lambda_1)$ is an isometry if and only if $\widehat{\lambda}_0 = F^* \widehat{\lambda}_1 F$. Next, we verify that ∂F is an isometry of the boundary linking forms.

Lemma 2.5. *If $F: (H_0, \lambda_0) \rightarrow (H_1, \lambda_1)$ is an isometry of nondegenerate Hermitian forms, then ∂F is an isometry of linking forms:*

$$\partial F: (\text{coker}(\widehat{\lambda}_0), \partial \lambda_0) \rightarrow (\text{coker}(\widehat{\lambda}_1), \partial \lambda_1).$$

This construction provides a map

$$\partial: \text{Iso}(\lambda_0, \lambda_1) \rightarrow \text{Iso}(\partial \lambda_0, \partial \lambda_1)$$

which is a homomorphism on automorphism groups.

Proof. Given $[x], [y] \in \text{coker}(\widehat{\lambda}_0)$, there exists $s \in \Lambda$ and $z \in H_0$ such that $sx = \widehat{\lambda}_0(z)$. Since F is an isometry, it follows that $sF^{-*}(x) = F^{-*}\widehat{\lambda}_0(z) = \widehat{\lambda}_1 F(z)$. We can now conclude that ∂F is an isometry because

$$\partial \lambda_1(\partial F([x]), \partial F([y])) = \partial \lambda_1(F^{-*}([x]), F^{-*}([y])) = \frac{1}{s} F^{-*}(y) F(z) = \frac{1}{s} y(z) = \partial \lambda_0([x], [y]).$$

The last assertion follows from the equality $(G \circ F)^{-*} = G^{-*} \circ F^{-*}$. \square

Using Lemma 2.5, we are led to the following definition.

Definition 2.6. The *boundary of an isometry* $F: (H_0, \lambda_0) \rightarrow (H_1, \lambda_1)$ of nondegenerate Hermitian forms is the isometry of linking forms

$$\partial F: \partial(H_0, \lambda_0) \rightarrow \partial(H_1, \lambda_1).$$

2.3. The union of nondegenerate Hermitian forms. We define the union of two Hermitian forms (H_0, λ_0) and (H_1, λ_1) over Λ along an isometry of their boundary linking forms. The definition is inspired by [Cro02, Chapter 3], which was concerned with forms over the integers.

We describe the main construction of this subsection. In what follows, for $i = 0, 1$, we use $\pi_i: H_i^* \rightarrow \text{coker}(\widehat{\lambda}_i)$ to denote the canonical projections.

Construction 2.7. Let (H_0, λ_0) and (H_1, λ_1) be two nondegenerate Hermitian forms over Λ , and let $h: (\text{coker}(\widehat{\lambda}_0), \partial\lambda_0) \rightarrow (\text{coker}(\widehat{\lambda}_1), \partial\lambda_1)$ be an isometry of their boundary linking forms. Consider the pair (H, λ) with

$$H := \ker (h\pi_0 - \pi_1: H_0^* \oplus H_1^* \rightarrow \text{coker}(\widehat{\lambda}_1))$$

$$\lambda \left(\begin{pmatrix} x_0 \\ x_1 \end{pmatrix}, \begin{pmatrix} y_0 \\ y_1 \end{pmatrix} \right) = \frac{1}{s_0} y_0(z_0) - \frac{1}{s_1} y_1(z_1) \in Q,$$

where since $\text{coker}(\widehat{\lambda}_i)$ is torsion, there exists $s_i \in \Lambda$ and $z_i \in H_i$ such that $s_i x_i = \widehat{\lambda}_i(z_i)$. Since the Hermitian forms λ_0 and λ_1 are nondegenerate, it is not difficult to prove that the pairing λ does not depend on the choice of s_0, s_1, z_0, z_1 .

The next proposition establishes some facts about the pairing (H, λ) .

Proposition 2.8. *The pair (H, λ) from Construction 2.7 satisfies the following properties.*

(1) *For $(x_0, x_1), (y_0, y_1) \in H$, the pairing λ can equivalently be defined as*

$$\lambda \left(\begin{pmatrix} x_0 \\ x_1 \end{pmatrix}, \begin{pmatrix} y_0 \\ y_1 \end{pmatrix} \right) = y_0(\widehat{\lambda}_{0,Q}^{-1}(x_0)) - y_1(\widehat{\lambda}_{1,Q}^{-1}(x_1)).$$

(2) *If we choose bases for H_0, H_1 and dual bases for H_0^*, H_1^* and let A_0, A_1 be Hermitian matrices representing λ_0, λ_1 , then*

$$\lambda \left(\begin{pmatrix} x_0 \\ x_1 \end{pmatrix}, \begin{pmatrix} y_0 \\ y_1 \end{pmatrix} \right) = x_0^T A_0^{-1} \bar{y}_0 - x_1^T A_1^{-1} \bar{y}_1.$$

(3) *The pairing λ is sesquilinear, Hermitian and takes values in Λ .*

(4) *The following two maps, which we abridge by $\widehat{\lambda}_0, \widehat{\lambda}_1$, are injective:*

$$(H_0, \lambda_0) \begin{pmatrix} \widehat{\lambda}_0 \\ 0 \end{pmatrix} \hookrightarrow (H, \lambda) \begin{pmatrix} 0 \\ \widehat{\lambda}_1 \end{pmatrix} \hookrightarrow (H_1, -\lambda_1).$$

Furthermore, these maps satisfy $\widehat{\lambda}_0(H_0)^\perp = \widehat{\lambda}_1(H_1)$.

Proof. To prove the first assertion, notice that since $\lambda_{i,Q}$ is a nonsingular pairing for $i = 0, 1$, we can write $\widehat{\lambda}_{i,Q}^{-1}(x_i) = z_i/s_i$ for some $z_i \in H$ and some $s_i \in \Lambda$. Consequently, $\widehat{\lambda}_i(z_i) = s_i x_i$ and we have $\frac{1}{s_i} y_i(z_i) = y_i(\widehat{\lambda}_{i,Q}^{-1}(x_i))$, as desired. This proves the first assertion. The second assertion now follows as in Remark 2.4.

We prove the third assertion. The fact that λ is sesquilinear and Hermitian follows from the second assertion and the fact that A_0^{-1} and A_1^{-1} are Hermitian. Next, using the definition of the boundary linking forms $\partial\lambda_0$ and $\partial\lambda_1$ from Definition 2.2, we have

$$\lambda \left(\begin{pmatrix} x_0 \\ x_1 \end{pmatrix}, \begin{pmatrix} y_0 \\ y_1 \end{pmatrix} \right) = \partial\lambda_0([x_0], [y_0]) - \partial\lambda_1([x_1], [y_1]) \pmod{\Lambda}.$$

By definition of H , we have $h[x_0] = x_1$ and $h[y_0] = y_1$. Then since h is an isometry, this latter expression vanishes:

$$\begin{aligned} \partial\lambda_0([x_0], [y_0]) - \partial\lambda_1([x_1], [y_1]) &\equiv \partial\lambda_0([x_0], [y_0]) - \partial\lambda_1(h[x_0], h[y_0]) \\ &\equiv \partial\lambda_0([x_0], [y_0]) - \partial\lambda_0([x_0], [y_0]) \equiv 0 \in Q/\Lambda. \end{aligned}$$

It follows that λ takes values in Λ , concluding the proof of the third assertion. We now prove the fourth assertion. First, we check that $\widehat{\lambda}_0$ is an isometric embedding; the proof for $\widehat{\lambda}_1$ is

identical. Since the pairings are nondegenerate, the maps are injective and given $z_0, z'_0 \in H_0$, we have

$$\lambda \left(\begin{pmatrix} \widehat{\lambda}_0(z_0) \\ 0 \end{pmatrix}, \begin{pmatrix} \widehat{\lambda}_0(z'_0) \\ 0 \end{pmatrix} \right) = \widehat{\lambda}(z'_0)(z_0) = \lambda_0(z_0, z'_0).$$

It remains to check that $\widehat{\lambda}_0(H_0)^\perp = \widehat{\lambda}_1(H_1)$. The inclusion $\widehat{\lambda}_0(H_0)^\perp \supseteq \widehat{\lambda}_1(H_1)$ is clear and so we prove the inclusion $\widehat{\lambda}_0(H_0)^\perp \subseteq \widehat{\lambda}_1(H_1)$. Assume that $(x_0, x_1) \in H$ (with $sx_0 = \widehat{\lambda}_0(z_0)$) satisfies for all $a_0 \in H_0$

$$0 = \lambda \left(\begin{pmatrix} x_0 \\ x_1 \end{pmatrix}, \begin{pmatrix} \widehat{\lambda}_0(a_0) \\ 0 \end{pmatrix} \right) = \frac{1}{s} \widehat{\lambda}_0(a_0)(z_0) = \frac{1}{s} \lambda_0(z_0, a_0) \in \Lambda.$$

As λ_0 is nondegenerate, this implies that $z_0 = 0$ and therefore $x_0 = 0$. But since $(x_0, x_1) \in H$, we also have $[x_1] = h[x_0] = 0$ and so $x_1 = \widehat{\lambda}_1(z_1)$ for some $z_1 \in H_1$. We therefore conclude that $\widehat{\lambda}_0(H_0)^\perp = \widehat{\lambda}_1(H_1)$, establishing the proposition. \square

Note that Proposition 2.8 does not contain a statement about the Λ -module H being free. Since we defined Hermitian forms over free Λ -modules, a *Hermitian module* (H, λ) will refer to a pair consisting of a finitely generated Λ -module H and a sesquilinear Hermitian pairing $\lambda: H \times H \rightarrow \Lambda$; we drop the requirement that H be free.

Definition 2.9. Let (H_0, λ_0) and (H_1, λ_1) be two nondegenerate Hermitian forms over Λ , and let $h: (\text{coker}(\widehat{\lambda}_0), \partial\lambda_0) \rightarrow (\text{coker}(\widehat{\lambda}_1), \partial\lambda_1)$ be an isometry of their boundary linking forms. The *union* of (H_0, λ_0) and (H_1, λ_1) along h is the Hermitian module $(H_0 \cup_h H_1, \lambda_0 \cup_h -\lambda_1)$ described in Construction 2.7 and Proposition 2.8:

$$\begin{aligned} H_0 \cup_h H_1 &:= \ker (h\pi_0 - \pi_1: H_0^* \oplus H_1^* \rightarrow \text{coker}(\widehat{\lambda}_1)) \\ \lambda_0 \cup_h -\lambda_1 \left(\begin{pmatrix} x_0 \\ x_1 \end{pmatrix}, \begin{pmatrix} y_0 \\ y_1 \end{pmatrix} \right) &= \frac{1}{s_0} y_0(z_0) - \frac{1}{s_1} y_1(z_1) \in \Lambda, \end{aligned}$$

where since $\text{coker}(\widehat{\lambda}_i)$ is torsion, there exists $s_i \in \Lambda$ and $z_i \in H_i$ such that $s_i x_i = \widehat{\lambda}_i(z_i)$.

Remark 2.10. We make a brief remark about the nonsingularity of $(H_0 \cup_h H_1, \lambda_0 \cup_h -\lambda_1)$. Proposition 2.8 contains no statement about $(H, \lambda) := (H_0 \cup_h H_1, \lambda_0 \cup_h -\lambda_1)$ being nonsingular. While this holds over the integers [Cro02, Lemma 3.6] and in the topological setting of Proposition 3.9, we will not prove it in the algebraic generality of this section. Note that if $\widehat{\lambda}: H \xrightarrow{\cong} \overline{\text{Hom}}_\Lambda(H, \Lambda) =: H^*$ is nonsingular, then H must be free, since for any finitely generated Λ -module H , the dual H^* is free [BF14, Lemma 2.1].

Given a Λ -module H , we set $H^* := \overline{\text{Hom}}_\Lambda(H, \Lambda)$. If H is free, then there is a Λ -linear evaluation isomorphism $\text{ev}: H \rightarrow H^{**}$ given by $\text{ev}_x(\varphi) = \overline{\varphi(x)}$ where $x \in H$ and $\varphi \in H^*$. The next remark uses ev to describe the adjoint of $\lambda_0 \cup_h -\lambda_1$.

Remark 2.11. Let (H_0, λ_0) and (H_1, λ_1) be two nondegenerate Hermitian forms over Λ , and let $h \in \text{Iso}(\partial\lambda_0, \partial\lambda_1)$ be an isometry. The adjoint of $\lambda := \lambda_0 \cup_h -\lambda_1$ is given by

$$\begin{aligned} \widehat{\lambda}: H_0 \cup_h H_1 &\rightarrow (H_0 \cup_h H_1)^* \\ \begin{pmatrix} y_0 \\ y_1 \end{pmatrix} &\mapsto (\text{ev} \quad \text{ev}) \begin{pmatrix} \widehat{\lambda}_{0,Q}^{-1} & 0 \\ 0 & -\widehat{\lambda}_{1,Q}^{-1} \end{pmatrix} \begin{pmatrix} y_0 \\ y_1 \end{pmatrix}. \end{aligned}$$

This follows by combining the definition of ev with the second and third items of Proposition 2.8. Indeed for $(x_0, y_0) \in H_0 \cup_h H_1$, we have

$$\begin{aligned} \lambda \left(\begin{pmatrix} x_0 \\ x_1 \end{pmatrix}, \begin{pmatrix} y_0 \\ y_1 \end{pmatrix} \right) &= \overline{\lambda \left(\begin{pmatrix} y_0 \\ y_1 \end{pmatrix}, \begin{pmatrix} x_0 \\ x_1 \end{pmatrix} \right)} = \overline{x_0(\widehat{\lambda}_{0,Q}^{-1}(y_0)) - x_1(\widehat{\lambda}_{1,Q}^{-1}(y_1))} \\ &= \begin{pmatrix} \text{ev}_{\widehat{\lambda}_{0,Q}^{-1}(y_0)} & \text{ev}_{-\widehat{\lambda}_{1,Q}^{-1}(y_1)} \end{pmatrix} \begin{pmatrix} x_0 \\ x_1 \end{pmatrix} = \widehat{\lambda} \begin{pmatrix} y_0 \\ y_1 \end{pmatrix} \begin{pmatrix} x_0 \\ x_1 \end{pmatrix}. \end{aligned}$$

We conclude this section by describing how the union interacts with boundary isometries. The proof follows immediately from the definitions and is left to the reader.

Proposition 2.12. *Let (H_0, λ_0) and (H_1, λ_1) be Hermitian forms, and let $h \in \text{Iso}(\partial\lambda_0, \partial\lambda_1)$ be an isometry. If $F \in \text{Iso}(\lambda_0, \lambda_1)$ is an isometry, then $F^{-*} \oplus \text{Id}$ induces an isometry*

$$(H_0, \lambda_0) \cup_h (H_1, -\lambda_1) \rightarrow (H_1, \lambda_1) \cup_{h \circ \partial F^{-1}} (H_1, -\lambda_1).$$

2.4. Lagrangian complements. This subsection provides a criterion for a union of Hermitian forms to be metabolic. It motivates the notion of a compatible pair that will be introduced in the next subsection. The idea for this criterion stems from work of Kreck [Kre99, Proposition 8] and Crowley-Sixt [CS11, Theorem 5.11] and their work on the monoid $\ell_{2q+1}(\mathbb{Z}[\pi])$.

A *Lagrangian* for a nonsingular Hermitian form (H, λ) is a direct summand $L \subseteq H$ such that $L^\perp = L$. A nonsingular Hermitian form that admits a Lagrangian is called *metabolic*. A *Lagrangian complement* for a half rank direct summand $V \subseteq (H, \lambda)$ is a Lagrangian of (H, λ) such that $L \oplus V = H$.

In the next proposition, for an isomorphism $F: H_0 \rightarrow H_1$ we consider the graph

$$\Gamma_{F^{-*}} = \{(x, F^{-*}(x)) \mid x \in H_0^*\} \subseteq H_0^* \oplus H_1^*.$$

Since, by definition of ∂F , we have $\partial F(x) - F^{-*}(x) = F^{-*}(x) - F^{-*}(x) = 0$, we deduce the inclusions $\Gamma_{F^{-*}} \subseteq \ker(\partial F \circ \pi_0 - \pi_1) = H_0 \cup_{\partial F} H_1 \subseteq H_0^* \oplus H_1^*$.

Proposition 2.13. *Let $F: (H_0, \lambda_0) \rightarrow (H_1, \lambda_1)$ be a Λ -isometry such that $\lambda_0 \cup_{\partial F} -\lambda_1$ is nonsingular. The graph $\Gamma_{F^{-*}} \subseteq H_0 \cup_{\partial F} H_1$ is a Lagrangian complement for $\widehat{\lambda}_0(H_0)$:*

$$H_0 \cup_{\partial F} H_1 = \widehat{\lambda}_0(H_0) \oplus \Gamma_{F^{-*}}.$$

Proof. Set $\lambda := \lambda_0 \cup_{\partial F} -\lambda_1$. We show that $\Gamma_{F^{-*}}$ is a Lagrangian. To see that $\Gamma_{F^{-*}} \subseteq \Gamma_{F^{-*}}^\perp$, we must show that for all $x_0, y_0 \in H_0^*$,

$$\lambda \left(\begin{pmatrix} x_0 \\ F^{-*}(x_0) \end{pmatrix}, \begin{pmatrix} y_0 \\ F^{-*}(y_0) \end{pmatrix} \right) = 0.$$

Pick $s_0 \in \Lambda$ and $z_0 \in H_0$ such that $s_0 x_0 = \widehat{\lambda}_0(z_0)$. We apply F^{-*} to both sides of this equation. Since F is an isometry, we have $F^* \widehat{\lambda}_1 F = \widehat{\lambda}_0$, and we therefore obtain $s_0 F^{-*}(x_0) = \widehat{\lambda}_1(F(z_0))$. Using the definition of λ , we now obtain the desired conclusion:

$$\lambda \left(\begin{pmatrix} x_0 \\ F^{-*}(x_0) \end{pmatrix}, \begin{pmatrix} y_0 \\ F^{-*}(y_0) \end{pmatrix} \right) = \frac{1}{s_0} y_0(z_0) - \frac{1}{s_0} F^{-*}(y_0)(F(z_0)) = 0.$$

Next, we show that $\Gamma_{F^{-*}}^\perp \subseteq \Gamma_{F^{-*}}$. We therefore assume that $(x_0, x_1) \in H_0 \cup_{\partial F} H_1$ satisfies, for all $y_0 \in H_0^*$, the equation

$$\lambda \left(\begin{pmatrix} x_0 \\ x_1 \end{pmatrix}, \begin{pmatrix} y_0 \\ F^{-*}(y_0) \end{pmatrix} \right) = 0$$

and we must show that $x_1 = F^{-*}(x_0)$. Pick $s_0, s_1 \in \Lambda, z_0 \in H_0$ and $z_1 \in H_1$ so that $s_0 x_0 = \widehat{\lambda}_0(z_0)$ and $s_1 x_1 = \widehat{\lambda}_1(z_1)$. We have $s_0 F^{-*}(x_0) = \widehat{\lambda}_1(F(z_0))$. Use consecutively the definition of λ and the definition of pullback to obtain

$$0 = \lambda \left(\begin{pmatrix} x_0 \\ x_1 \end{pmatrix}, \begin{pmatrix} y_0 \\ F^{-*}(y_0) \end{pmatrix} \right) = y_0 \begin{pmatrix} z_0 \\ s_0 \end{pmatrix} - F^{-*}(y_0) \begin{pmatrix} z_1 \\ s_1 \end{pmatrix} = y_0 \begin{pmatrix} z_0 \\ s_0 \end{pmatrix} - F^{-1} \begin{pmatrix} z_1 \\ s_1 \end{pmatrix}.$$

Since this equation holds for all y_0 , we deduce that

$$\frac{z_0}{s_0} = F^{-1} \begin{pmatrix} z_1 \\ s_1 \end{pmatrix}.$$

Using that F is an isometry, as well as the definitions of s_0, s_1, z_0 and z_1 , we obtain the desired equation:

$$x_0 = \widehat{\lambda}_0 \begin{pmatrix} z_0 \\ s_0 \end{pmatrix} = \widehat{\lambda}_0 \left(F^{-1} \begin{pmatrix} z_1 \\ s_1 \end{pmatrix} \right) = F^* \left(\widehat{\lambda}_1 \begin{pmatrix} z_1 \\ s_1 \end{pmatrix} \right) = F^*(x_1).$$

To conclude the proof that $\Gamma_{F^{-*}}$ is a Lagrangian, it remains to show that it is a direct summand of $H := \ker(\partial F \circ \pi_0 - \pi_1)$. In fact, we will show that

$$\widehat{\lambda}_0(H_0) \oplus \Gamma_{F^{-*}} = H.$$

First we establish the inclusion $\widehat{\lambda}_0(H_0) + \Gamma_{F^{-*}} \subseteq H$. To do so, we must prove that the sum of arbitrary elements $(\widehat{\lambda}_0(z_0), 0) \in \widehat{\lambda}_0(H_0)$ and $(y, F^{-*}(y)) \in \Gamma_{F^{-*}}$ belongs to H . In other words, we must show that $\partial F[\widehat{\lambda}_0(z_0) + y] = [F^{-*}(y)]$. This uses the fact that F^{-*} induces ∂F (by Definition 2.6) and the fact that, since F is an isometry, $F^{-*} \circ \widehat{\lambda}_0(z_0) = \widehat{\lambda}_1(F(z_0))$ belongs to $\text{im}(\widehat{\lambda}_1)$ and so the class $[F^{-*} \circ \widehat{\lambda}_0(z_0)]$ must vanish in the quotient $\text{coker}(\widehat{\lambda}_1)$:

$$\partial F[\widehat{\lambda}_0(z_0) + y] = [F^{-*} \widehat{\lambda}_0(z_0)] + [F^{-*}(y)] = [F^{-*}(y)].$$

Next, we prove the reverse inclusion, namely that $H \subseteq \widehat{\lambda}_0(H_0) + \Gamma_{F^{-*}}$. To show this, we must write $(x, y) \in H$ as a sum of an element in $\widehat{\lambda}_0(H_0)$ with an element in $\Gamma_{F^{-*}}$. Consider the decomposition

$$(x, y) = (x - F^*(y), 0) + (F^*(y), y).$$

The second term certainly lies in $\Gamma_{F^{-*}}$, so we need to argue that the first term lies in $\widehat{\lambda}_0(H_0)$. Since $(x, y) \in H$, we know that $\partial F[x] = [y]$ in $\text{coker}(\widehat{\lambda}_1)$. It follows that $F^{-*}(x) - y \in H_1^*$ belongs to $\text{im}(\widehat{\lambda}_1)$. Since F is an isometry, this is equivalent to saying that $x - F^*(y) \in \text{im}(\widehat{\lambda}_0)$, as desired.

Having proved that $\widehat{\lambda}_0(H_0) + \Gamma_{F^{-*}} = H$, it remains to check that $\widehat{\lambda}_0(H_0) \cap \Gamma_{F^{-*}} = 0$ (so that the sum is direct): if $(\widehat{\lambda}_0(x_0), 0) = (x, F^{-*}(x))$ belongs to this intersection, then $F^{-*}(x) = 0$ and therefore $0 = x = \widehat{\lambda}_0(x_0)$. This concludes the proof of Proposition 2.13. \square

3. THE INTERSECTION FORM OF THE UNION OF TWO 4-MANIFOLDS

We start our study of 4-manifolds with infinite cyclic fundamental group and of their Λ -intersection forms. First, recall the following definition from the introduction.

Definition 3.1. A 4-manifold M has *ribbon boundary* if ∂M is nonempty and path-connected, and the map $\pi_1(\partial M) \rightarrow \pi_1(M)$ induced by the inclusion is surjective.

Here is a summary of Section 3. In Subsection 3.1, we show that if M is a 4-manifold with infinite cyclic fundamental group whose boundary is ribbon and has torsion Alexander module, then the boundary linking form $\partial\lambda_M$ of λ_M is isometric to minus the Blanchfield form of ∂M . In Subsection 3.2, we describe when homeomorphisms of 3-manifolds induce isometries of the Blanchfield pairing. In Subsection 3.3 we show that the Λ -intersection form of a union of two manifolds can be expressed using the algebraic union from Subsection 2.3. In Subsection 3.4, we introduce the notion of a compatible pair and prove Theorem 1.12 from the introduction.

3.1. The boundary of the intersection form. If a 4-manifold M with $\pi_1(M) = \mathbb{Z}$ has ribbon boundary and Λ -torsion Alexander module, then ∂M is endowed with a sesquilinear Hermitian nonsingular linking form over Λ , namely the Blanchfield form

$$\text{Bl}_{\partial M}: H_1(\partial M; \Lambda) \times H_1(\partial M; \Lambda) \rightarrow Q/\Lambda.$$

This subsection establishes that the boundary linking form of the Λ -intersection form of M is isometric to minus the Blanchfield form of ∂M . Since results of this form are known (see e.g. [BF15, Theorem 2.6]), we only outline the argument so as to fix some notations for later use.

We collect some homological facts about 4-manifolds with infinite cyclic fundamental group.

Lemma 3.2. *Let M be a 4-manifold with $\pi_1(M) = \mathbb{Z}$. When ∂M is nonempty, we assume it is ribbon and has torsion Alexander module $H_1(\partial M; \Lambda)$. The following assertions hold:*

- (1) $H_0(M; \Lambda) = \mathbb{Z}$ and, if $\partial M \neq \emptyset$, then $H_2(\partial M; \Lambda) = \mathbb{Z}$;
- (2) the Λ -modules $H_1(M; \Lambda)$, $H_1(M, \partial M; \Lambda)$ and $H_3(M; \Lambda)$ all vanish;
- (3) the Λ -modules $H_2(M; \Lambda)$ and $H_2(M, \partial M; \Lambda)$ are free;
- (4) when $\partial M = \emptyset$, the Λ -intersection form λ_M is nonsingular, whereas for $\partial M \neq \emptyset$, λ_M is nondegenerate, and any matrix representing it presents the Alexander module of ∂M .

Proof. Throughout this proof we use that since the manifold has fundamental group \mathbb{Z} , homology with Λ -coefficients can be computed as the homology of the universal cover. Since universal covers are 1-connected, we immediately deduce that $H_0(M; \Lambda) = \mathbb{Z}$ and $H_1(M; \Lambda) = 0$. If the boundary is nonempty, the long exact sequence of the pair shows that $H_1(M, \partial M; \Lambda) = 0$; here we used that ∂M is connected, as assumed in Definition 3.1. For later use, we also note that $H_0(M, \partial M; \Lambda) = 0$, also owing to the fact that ∂M is connected. As $H_1(\partial M; \Lambda)$ is torsion, we have $\text{Hom}_\Lambda(H_1(\partial M; \Lambda), \Lambda) = 0$, and thus $H_2(\partial M; \Lambda) \cong \text{Ext}_\Lambda^1(H_0(\partial M; \Lambda), \Lambda) \cong \mathbb{Z}$ by duality and the universal coefficient spectral sequence [Lev77, Theorem 2.3]. Since $H_i(M, \partial M; \Lambda) = 0$ for $i = 0, 1$, duality and the universal coefficient spectral sequence (UCSS for short) also show that $H_3(M; \Lambda) = H^1(M, \partial M; \Lambda) = 0$. Thus we have proved the first two assertions; we now prove the third. Poincaré duality and the UCSS imply that

$$\begin{aligned} H_2(M; \Lambda) &\cong H^2(M, \partial M; \Lambda) \cong \overline{\text{Hom}_\Lambda(H_2(M, \partial M; \Lambda), \Lambda)}, \\ H_2(M, \partial M; \Lambda) &\cong H^2(M; \Lambda) \cong \overline{\text{Hom}_\Lambda(H_2(M; \Lambda), \Lambda)}. \end{aligned}$$

Since the dual of a finitely generated Λ -module is free [BF14, Lemma 2.1], we deduce that these second homology modules are free over Λ . This proves the third assertion and also establishes the fourth, namely that in the closed case, the Λ -intersection form is nonsingular. Finally, when the boundary is nonempty, the map $\mathbb{Z} = H_2(\partial M; \Lambda) \rightarrow H_2(M; \Lambda)$ is the zero map (since $H_2(M; \Lambda)$ is free) and therefore the intersection form is nondegenerate; it presents $H_1(\partial M; \Lambda)$ because we established that $H_1(M; \Lambda) = 0$. \square

In the case with nonempty connected boundary, we fix bases for the free Λ -modules $H_2(M; \Lambda)$ and $H_2(M, \partial M; \Lambda)$.

Remark 3.3. Fix a basis \mathbf{b} for $H := H_2(M; \Lambda)$, endow the dual $H^* := \overline{\text{Hom}_\Lambda(H, \Lambda)}$ with the dual basis \mathbf{b}^* , and equip $H_2(M, \partial M; \Lambda)$ with the basis $\text{PD ev}^{-1}(\mathbf{b}^*)$ coming from the isomorphisms $\text{ev}: H^2(M; \Lambda) \rightarrow H^*$ and $\text{PD}: H^2(M; \Lambda) \rightarrow H_2(M, \partial M; \Lambda)$. A short computation shows that if the matrix A is defined as $A_{ij} := \lambda_M(b_i, b_j)$, then $A^T = \bar{A}$ is a matrix for $\widehat{\lambda}_M: H \rightarrow H^*$; recall Remark 2.1. The same conclusion holds for the map $H \rightarrow H_2(M, \partial M; \Lambda)$ induced by the inclusion [CFT18, Section 5.2]. As mentioned in the fourth item of Lemma 3.2, the connecting homomorphism δ in the long exact sequence of the pair $(M, \partial M)$ induces an isomorphism

$$\text{coker}(\widehat{\lambda}_M) = \text{coker}(A^T) \xrightarrow{\cong, \delta} H_1(\partial M; \Lambda).$$

Next, we briefly recall the definition of the Blanchfield pairing, referring to [CFT18, FP17] as references in which the conventions are identical to ours.

Definition 3.4. Let N be a closed 3-manifold with an epimorphism $\varphi: \pi_1(N) \rightarrow \mathbb{Z}$ such that the resulting Alexander module is Λ -torsion. The adjoint of the Blanchfield pairing is defined by the composition

$$H_1(N; \Lambda) \xrightarrow{\text{PD}^{-1}, \cong} H^2(N; \Lambda) \xrightarrow{BS^{-1}, \cong} H^1(N; Q/\Lambda) \xrightarrow{\text{ev}, \cong} \overline{\text{Hom}_\Lambda(H_1(N; \Lambda), Q/\Lambda)}.$$

of the inverse of Poincaré duality, the inverse of a Bockstein homomorphism, and the evaluation map. The Blanchfield pairing is a nonsingular linking form, so in particular it is sesquilinear and Hermitian.

Now we can prove the main result of this subsection.

Proposition 3.5. *If M is a 4-manifold with $\pi_1(M) = \mathbb{Z}$ whose boundary is ribbon and has torsion Alexander module, then there is an isometry*

$$\partial(H_2(M; \Lambda), \lambda_M) \cong (H_1(\partial M; \Lambda), -\text{Bl}_{\partial M}).$$

Proof. Choose bases for $H_2(M; \Lambda)$ and $H_2(M, \partial M; \Lambda)$ as in Remark 3.3, set $A_{ij} := \lambda_M(b_i, b_j)$, and use δ to denote the connecting homomorphism from the long exact sequence of the pair $(M, \partial M)$. The same argument as in [CFT18, Section 5] shows that the following diagram (with exact columns) commutes:

$$\begin{array}{ccc} 0 & & 0 \\ \downarrow & & \downarrow \\ H_2(M; \Lambda) \times H_2(M; \Lambda) & \xrightarrow{(x, y) \mapsto -x^T A \bar{y}} & \Lambda \\ \downarrow A^T \times A^T & & \downarrow \\ H_2(M, \partial M; \Lambda) \times H_2(M, \partial M; \Lambda) & \xrightarrow{(x, y) \mapsto -x^T A^{-1} \bar{y}} & Q \\ \downarrow \delta \times \delta & & \downarrow \\ H_1(\partial M; \Lambda) \times H_1(\partial M; \Lambda) & \xrightarrow{\text{Bl}_{\partial M}} & Q/\Lambda \\ \downarrow & & \downarrow \\ 0 & & 0. \end{array}$$

In particular, δ induces an isometry between $(\text{coker}(\widehat{\lambda}_M), \partial \lambda_M)$ and $(H_1(\partial M; \Lambda), -\text{Bl}_{\partial M})$; here we are using the characterisation of $\partial \lambda_M$ from Remark 2.4. \square

The next remark records that the presence of a minus sign in Proposition 3.5 is immaterial once one passes to isometry groups.

Remark 3.6. Given linking forms (H_0, β_0) and (H_1, β_1) , observe that canonically we have $\text{Iso}(\beta_0, \beta_1) = \text{Iso}(-\beta_0, -\beta_1)$; the presence of the minus sign has no effect. In particular if M_0, M_1 are 4-manifolds with infinite cyclic fundamental group whose boundaries are ribbon and have torsion Alexander modules, then we can use Proposition 3.5 to obtain an identification

$$\text{Iso}(\partial\lambda_{M_0}, \partial\lambda_{M_1}) = \text{Iso}(\text{Bl}_{\partial M_0}, \text{Bl}_{\partial M_1}).$$

3.2. Homeomorphisms and isometries. In this short subsection, we provide a condition for a homeomorphism between 3-manifolds to induce an isometry of Blanchfield forms.

Proposition 3.7. *Let Y_0^3, Y_1^3 be 3-manifolds equipped with epimorphisms $\varphi_i: \pi_1(Y_i) \rightarrow \mathbb{Z}$ and assume that the resulting Alexander modules $H_1(Y_i; \Lambda)$ are Λ -torsion for $i = 0, 1$. If an orientation-preserving homotopy equivalence $f: Y_0^3 \rightarrow Y_1^3$ satisfies $\varphi_1 \circ f_* = \varphi_0$ on $\pi_1(Y_0)$, then it induces an isometry between the Blanchfield forms:*

$$\tilde{f}_*: (H_1(Y_0; \Lambda), \text{Bl}_{Y_0}) \rightarrow (H_1(Y_1; \Lambda), \text{Bl}_{Y_1}).$$

Proof. The homotopy equivalence f induces Λ -isomorphisms $f: H_*(Y_0; \Lambda) \rightarrow H_*(Y_1; \Lambda)$ (and similarly on cohomology) because we assumed that $\varphi_1 \circ f_* = \varphi_0$. The fact that f has degree one, together with the naturality of the UCSS and the Bockstein homomorphism ensure that \tilde{f}_* intertwines the Blanchfield pairings, concluding the proof of the proposition. \square

For pairs (Y_0, φ_0) and (Y_1, φ_1) as in the statement of Proposition 3.7, we consider those orientation-preserving homeomorphisms that intertwine the φ_i :

$$\text{Homeo}_\varphi^+(Y_0, Y_1) = \{f \in \text{Homeo}^+(Y_0, Y_1) \mid \varphi_1 \circ f_* = \varphi_0\}.$$

In the case where $(Y_0, \varphi_0) = (Y_1, \varphi_1)$, we simply write $\text{Homeo}_\varphi^+(Y_0)$ for the resulting group.

3.3. Algebraic unions and topological unions. In this subsection, we show that under favourable conditions, the Λ -intersection form of a union of two 4-manifolds can be expressed using the algebraic union of Hermitian forms from Subsection 2.3. For that however, given two manifolds M_0, M_1 with $\pi_1(M_i) \cong \mathbb{Z}$, we need to verify that $\pi_1(M_0 \cup_f -M_1) \cong \mathbb{Z}$.

Lemma 3.8. *If $f \in \text{Homeo}_\varphi^+(\partial M_0, \partial M_1)$ is a homeomorphism where M_0, M_1 are 4-manifolds with ribbon boundaries and $\pi_1(M_i) \cong \mathbb{Z}$, then $\pi_1(M_0 \cup_f -M_1) \cong \mathbb{Z}$.*

Proof. For the sake of applying van Kampen's theorem, for $i = 0, 1$ we write $\pi_1(M_i) = \mathbb{Z}\langle t_i \rangle$, so that $\pi_1(M_0 \cup_f -M_1)$ is generated by t_0, t_1 . Consider the following commutative diagram:

$$\begin{array}{ccc} \pi_1(\partial M_0) & \xrightarrow{f_*, \cong} & \pi_1(\partial M_1) \\ \downarrow \iota_0 & \searrow \varphi_0 & \swarrow \varphi_1 & \downarrow \iota_1 \\ \pi_1(M_0) = \mathbb{Z}\langle t_0 \rangle & \xrightarrow{\psi_0, \cong} & \mathbb{Z}\langle t \rangle & \xleftarrow{\psi_1, \cong} & \mathbb{Z}\langle t_1 \rangle = \pi_1(M_1). \end{array}$$

Here, for $j = 0, 1$, $\iota_j: \pi_1(\partial M_j) \rightarrow \pi_1(M_j)$ is the inclusion induced map and $\psi_j: t_j \mapsto t$. In the group $\pi_1(M_0 \cup_f -M_1)$, the relations identify $\iota_1 f_*(g)$ with $\iota_0(g)$ for every $g \in \pi_1(\partial M_0)$. Since ι_0 is surjective, there is a $\mu \in \pi_1(\partial M_0)$ with $\iota_0(\mu) = t_0$. We deduce that in $\pi_1(M_0 \cup_f -M_1)$, the relations identify $t_0 = \varphi_0(\mu)$ with $\iota_1(f_*(\mu)) = t_1^n$ for some n . Under the identification given by the ψ_i and using the fact that $\varphi_0(\mu) = \varphi_1(f_*(\mu))$, we deduce that $n = 1$. Thus $\pi_1(M_0 \cup_f -M_1)$

is cyclic. For every $g \in \pi_1(\partial M_0)$, we have $\iota_0(g) = t_0^m$ for some m , and the same reasoning as above shows that $\iota_0(g) = t_0^m$ is identified with $\iota_1(f_*(g)) = t_1^m$ in $\pi_1(M_0 \cup_f -M_1)$. It follows that $\pi_1(M_0 \cup_f -M_1)$ is isomorphic to $\langle t_0, t_1 \mid t_0 = t_1 \rangle \cong \mathbb{Z}$. \square

Next, we prove the main result of this section.

Proposition 3.9. *Let M_0 and M_1 be two 4-manifolds with infinite cyclic fundamental group whose boundaries are ribbon and have torsion Alexander modules. Let $f \in \text{Homeo}_\varphi^+(\partial M_0, \partial M_1)$ be a homeomorphism, set $M := M_0 \cup_f -M_1$, and let $\iota_j : M_j \rightarrow M$ be the inclusion. The map $\begin{pmatrix} \iota_0^* \\ \iota_1^* \end{pmatrix} : H_2(M; \Lambda)^* \rightarrow H_2(M_0; \Lambda)^* \oplus H_2(M_1; \Lambda)^*$ induces an isometry*

$$(1) \quad (H_2(M; \Lambda)^*, \lambda_M^{-1}) \xrightarrow{\cong} (H_2(M_0; \Lambda) \cup_{\tilde{f}_*} H_2(M_1; \Lambda), \lambda_{M_0} \cup_{\tilde{f}_*} -\lambda_{M_1}).$$

The map $\widehat{\lambda}_M$ induces an isometry $(H_2(M; \Lambda), \lambda_M) \rightarrow (H_2(M; \Lambda)^*, \lambda_M^{-1})$ which, using the identification from (1), makes the following diagram of isometries commutes:

$$(2) \quad \begin{array}{ccc} (H_2(M_0; \Lambda), \lambda_{M_0}) & \begin{array}{c} \xrightarrow{\iota_0} \\ \searrow \widehat{\lambda}_{M_0} \end{array} & (H_2(M; \Lambda), \lambda_M) & \begin{array}{c} \xleftarrow{\iota_1} \\ \swarrow \widehat{\lambda}_{M_1} \end{array} & (H_2(M_1; \Lambda), -\lambda_{M_1}) \\ & & \cong \downarrow \widehat{\lambda}_M & & \\ & & (H_2(M_0; \Lambda) \cup_{\tilde{f}_*} H_2(M_1; \Lambda), \lambda_{M_0} \cup_{\tilde{f}_*} -\lambda_{M_1}). & & \end{array}$$

In particular, $H_2(M_0; \Lambda) \cup_{\tilde{f}_*} H_2(M_1; \Lambda)$ is a free Λ -module and $\lambda_{M_0} \cup_{\tilde{f}_*} -\lambda_{M_1}$ is nonsingular.

Here, note that $\widehat{\lambda}_M : H_2(M; \Lambda) \rightarrow H_2(M; \Lambda)^*$ is an isomorphism because $\partial M = \emptyset$; recall the fourth item of Lemma 3.2. This endows $H_2(M; \Lambda)^*$ with a Hermitian form which we denoted λ_M^{-1} in (1).

Proof. Using Proposition 3.5, we think of the Blanchfield form on ∂M_i as being defined on the torsion module $\text{coker}(\widehat{\lambda}_i)$. We also recall from Proposition 3.7 that the homeomorphism $f : \partial M_0 \rightarrow \partial M_1$ induces an isometry \tilde{f}_* between these linking forms. Lemma 3.8 implies that $\pi_1(M) = \mathbb{Z}$. Consider the following commutative diagram in which the first and third rows are exact and Λ coefficients are understood:

$$(3) \quad \begin{array}{ccccccc} 0 & \longrightarrow & H_2(M_0) \oplus H_2(M_1) & \xrightarrow{(\iota_0 \ \iota_1)} & H_2(M) & \longrightarrow & H_1(\partial M_1) \longrightarrow 0 \\ & & \downarrow j_0 \oplus j_1 & & \downarrow \text{PD}_M^{-1} & & \\ & & H_2(M_0, \partial M_0) \oplus H_2(M_1, \partial M_1) & & & & \\ & & \downarrow \text{PD}_{M_0}^{-1} \oplus -\text{PD}_{M_1}^{-1} & & \downarrow & & \\ H^2(\partial M_1) & \xleftarrow{\tilde{f}_*^{-1} \text{incl}_0^* - \text{incl}_1^*} & H^2(M_0) \oplus H^2(M_1) & \xleftarrow{\begin{pmatrix} \iota_0^* \\ \iota_1^* \end{pmatrix}} & H^2(M) & \xleftarrow{\quad} & 0 \\ & & \downarrow \cong \text{ev} \oplus \text{ev} & & \downarrow \cong \text{ev} & & \\ & & H_2(M_0)^* \oplus H_2(M_1)^* & \xleftarrow{\begin{pmatrix} \iota_0^* \\ \iota_1^* \end{pmatrix}} & H_2(M)^* & & \end{array}$$

We justify the three zeros that appear. In the first row, the rightmost zero comes from the fact that $H_1(M_i; \Lambda) = 0$ for $i = 0, 1$ by the second item of Lemma 3.2. For the leftmost zero of the first row, the first and third items of Lemma 3.2 respectively imply that $H_2(\partial M_i; \Lambda) = \mathbb{Z}$

is Λ -torsion and that $H_2(M_i; \Lambda)$ is free. It follows that the inclusion induced homomorphism $H_2(\partial M_i; \Lambda) \rightarrow H_2(M_i; \Lambda)$ is zero. The same argument explains the appearance of the zero in the third row: $H^2(M; \Lambda)$ is free and $H^1(\partial M_1; \Lambda)$ is Λ -torsion. The commutativity of the middle square follows by applying [Bre93, Lemma 8.2] with $K = M_0$ and $L = M_1$, as well as applying excision to the pairs $(M, M \setminus M_i)$.

We choose bases for these various free Λ -modules as in Remark 3.3. First, fix bases $\mathbf{b}^0, \mathbf{b}^1$ for

$$H_0 := H_2(M_0; \Lambda) \text{ and } H_1 := H_2(M_1; \Lambda).$$

Then, for $i = 0, 1$ we endow H_i^* with the dual bases \mathbf{b}_i^* , and $H_2(M_i, \partial M_i; \Lambda)$ with the basis $\text{PD}_{M_i} \text{ev}^{-1}(\mathbf{b}_i^*)$. Finally, endow $H^2(M_i; \Lambda)$ with the basis $\text{ev}^{-1}(\mathbf{b}_i^*)$. We set $\lambda_i := \lambda_{M_i}$ as well as $(A_i)_{kl} := \lambda_i(b_k^i, b_l^i)$ for $i = 0, 1$. Using these bases, we recalled in Remark 3.3 that $\widehat{\lambda}_i$ and the map $H_2(M_i; \Lambda) \rightarrow H_2(M_i, \partial M_i; \Lambda)$ induced by the inclusion are represented by A_i^T , and that $H_1(M_i; \Lambda)$ can be identified with $\text{coker}(A_i^T) = \text{coker}(\widehat{\lambda}_i)$. Using the commutative diagram

$$\begin{array}{ccccc} H_2(M_i; \Lambda) & \xrightarrow{j_i} & H_2(M_i, \partial M_i; \Lambda) & \longrightarrow & H_1(\partial M_i; \Lambda) \\ \downarrow \text{PD}^{-1} & & \downarrow \text{PD}^{-1} & & \downarrow \text{PD}^{-1} \\ H^2(M_i, \partial M_i; \Lambda) & \longrightarrow & H^2(M_i; \Lambda) & \xrightarrow{\text{incl}_i^*} & H^2(\partial M_i; \Lambda), \end{array}$$

we deduce that $H^2(\partial M_i; \Lambda)$ can also be identified with $\text{coker}(A_i^T) = \text{coker}(\widehat{\lambda}_i)$. Similarly, the map $\text{PD}^{-1} \circ j_i: H_i \rightarrow H^2(M_i; \Lambda)$ is also represented by A_i^T . We therefore allow ourselves to write H_i^* both for $H_2(M_i; \Lambda)^*$ and for $H^2(M_i; \Lambda)$. We set $V := H_2(M; \Lambda)^*$ and identify $H_2(M; \Lambda)$ with the double dual $H_2(M; \Lambda)^{**} = V^*$ via the evaluation isomorphism ev . Considering the projection $\pi_i: H^2(M_i; \Lambda) \rightarrow \text{coker}(A_i^T)$ and using λ_M to denote the Λ -intersection form on M , and looking at (3), we obtain the following commutative diagram in which both rows are exact:

$$(4) \quad \begin{array}{ccccccc} 0 & \longrightarrow & H_0 \oplus H_1 & \xrightarrow{(\text{ev} \ \text{ev})} & V^* & \longrightarrow & T \\ & & \downarrow \widehat{\lambda}_0 \oplus -\widehat{\lambda}_1 & & \cong \downarrow \widehat{\lambda}_M & & \\ \text{coker}(\widehat{\lambda}_1) & \xleftarrow{\widetilde{f}_* \pi_0 - \pi_1} & H_0^* \oplus H_1^* & \longleftarrow & V & \longleftarrow & 0. \end{array}$$

Identify $V = H_2(M; \Lambda)^*$ with $\ker(\widetilde{f}_* \pi_0 - \pi_1)$ and view it as a subspace of $H_0^* \oplus H_1^*$. The diagram in (4) therefore yields the equation

$$(5) \quad \begin{pmatrix} \widehat{\lambda}_0 & 0 \\ 0 & -\widehat{\lambda}_1 \end{pmatrix} = \widehat{\lambda}_M \circ (\text{ev} \ \text{ev}).$$

Since Lemma 3.2 implies that $H_2(M; \Lambda)$ is a free Λ -module and $H_1(M; \Lambda) = 0$, we may identify $H^2(M; Q)$ with $H^2(M; \Lambda) \otimes_{\Lambda} Q$, and similarly for $H^2(M_i, \partial M_i; Q) \cong H^2(M_i, \partial M_i; \Lambda) \otimes_{\Lambda} Q$ and $H^2(M_i; Q) \cong H^2(M_i; \Lambda) \otimes_{\Lambda} Q$ for $i = 0, 1$. As a consequence, we may identify the tensored up intersection pairings $\lambda_M \otimes_{\Lambda} \text{Id}_Q$ and $\lambda_i \otimes_{\Lambda} \text{Id}_Q$ with the nonsingular Q -valued intersections pairings on $H_2(M; Q)$ and $H_2(M_i; Q)$. Since $\widehat{\lambda}_M$ is invertible by Lemma 3.2, we deduce from (5) that

$$\widehat{\lambda}_M^{-1} = (\text{ev} \ \text{ev}) \circ \begin{pmatrix} \widehat{\lambda}_{0,Q}^{-1} & 0 \\ 0 & -\widehat{\lambda}_{1,Q}^{-1} \end{pmatrix}.$$

As noted in Remark 2.11, this is precisely the adjoint of $\lambda_0 \cup_{\widetilde{f}_*} -\lambda_1$.

We have thus proved that the inclusions ι_0 and ι_1 induce an isometry $(H_2(M; \Lambda)^*, \lambda_M^{-1}) \cong (H_0 \cup_{\tilde{f}_*} H_1, \lambda_0 \cup_{\tilde{f}_*} -\lambda_1)$. Now, since $\lambda_M^* = \lambda_M$, a quick verification shows that the isomorphism $\widehat{\lambda}_M: H_2(M; \Lambda) \rightarrow H_2(M; \Lambda)^*$ induces an isometry between the forms $(H_2(M; \Lambda), \lambda_M)$ and $(H_2(M; \Lambda)^*, \lambda_M^{-1})$. Indeed, writing $H := H_2(M; \Lambda)$, this follows from the following commutative diagram:

$$\begin{array}{ccc} H & \xrightarrow{\widehat{\lambda}_M} & H^* \\ \downarrow \widehat{\lambda}_M & \cong & \downarrow \widehat{\lambda}_M^{-1} \\ H^* & \xleftarrow{\widehat{\lambda}_M^* = \widehat{\lambda}_M} & H. \end{array}$$

The commutativity of the diagram in (2) follows from the commutativity of (4): we have shown that $(H_0 \cup_{\tilde{f}_*} H_1, \lambda_0 \cup_{\tilde{f}_*} -\lambda_1) \cong (H_2(M; \Lambda), \lambda_M)$ and, recalling that $(\text{ev}, \text{ev}) = (\iota_0, \iota_1)$, the diagram in (4) gives $\widehat{\lambda}_M \circ \iota_0(x_0, 0) = \widehat{\lambda}_{M_0}(x_0, 0)$ leading to the commutativity of the left hand triangle of (2); the reasoning for the right hand triangle is identical. \square

3.4. Compatible pairs. Throughout this section, M_0 and M_1 are 4-manifolds with infinite cyclic fundamental group whose boundaries are ribbon and have torsion Alexander modules. This section introduces a notion of compatibility for an isometry $F: H_2(M_0; \Lambda) \rightarrow H_2(M_1; \Lambda)$ and a homeomorphism $f: \partial M_0 \rightarrow \partial M_1$. We then prove Theorem 1.12 from the introduction.

As the boundaries are ribbon, the inclusions $\partial M_i \hookrightarrow M_i$ induce surjections $\varphi_i: \pi_1(M_i) \twoheadrightarrow \mathbb{Z}$ for $i = 0, 1$. Recall from Subsection 3.2 that we set

$$\text{Homeo}_\varphi^+(\partial M_0, \partial M_1) = \{f \in \text{Homeo}^+(\partial M_0, \partial M_1) \mid \varphi_1 \circ f_* = \varphi_0\}.$$

Proposition 3.7 ensures that a homeomorphism $f \in \text{Homeo}_\varphi^+(\partial M_0, \partial M_1)$ induces an isometry $\tilde{f}_* \in \text{Iso}(\text{Bl}_{\partial M_0}, \text{Bl}_{\partial M_1})$ of the Blanchfield forms. On the other hand, recall from Lemma 2.5 that an isometry $F \in \text{Iso}(\lambda_{M_0}, \lambda_{M_1})$ of the Λ -intersection forms of M_0 and M_1 induces an isometry $\partial F = F^{-*}$ of the boundary linking forms. Using Proposition 3.5, we identify these boundary linking forms with minus the corresponding Blanchfield forms so that we may consider $\partial F \in \text{Iso}(\text{Bl}_{\partial M_0}, \text{Bl}_{\partial M_1})$; recall also Remark 3.6.

Definition 3.10. An orientation-preserving homeomorphism $f \in \text{Homeo}_\varphi^+(\partial M_0, \partial M_1)$ is *compatible* with an isometry $F \in \text{Iso}(\lambda_{M_0}, \lambda_{M_1})$ if

$$\partial F = \tilde{f}_*: (H_1(\partial M_0; \Lambda), \text{Bl}_{\partial M_0}) \rightarrow (H_1(\partial M_1; \Lambda), \text{Bl}_{\partial M_1}).$$

In this case, the pair (f, F) is called a *compatible pair*.

The next proposition shows that the existence of a compatible pair is necessary for a homeomorphism $f \in \text{Homeo}_\varphi^+(\partial M_0, \partial M_1)$ to extend to a homeomorphism $M_0 \rightarrow M_1$.

Proposition 3.11. *Let $f \in \text{Homeo}_\varphi^+(\partial M_0, \partial M_1)$ be a homeomorphism, and let $F \in \text{Iso}(\lambda_{M_0}, \lambda_{M_1})$ be an isometry. If f extends to an orientation-preserving homeomorphism $\Phi: M_0 \xrightarrow{\cong} M_1$ that induces F , then (f, F) is a compatible pair.*

Proof. The homeomorphism Φ induces an isomorphism $\Phi_*: \pi_1(M_0) \rightarrow \pi_1(M_1)$. Arrange the isomorphisms $\pi_1(M_i) \cong \mathbb{Z}$ for $i = 0, 1$ so that Φ_* intertwines them. It follows that the

homeomorphism Φ induces maps on twisted homology. In particular, we obtain the isometry $\tilde{\Phi}_*: H_2(M_0; \Lambda) \xrightarrow{\cong} H_2(M_1; \Lambda)$ that we assumed to coincide with F . We argue that (f, F) is a compatible pair. Naturality of homology shows that the following diagram commutes:

$$\begin{array}{ccccc} H_2(M_0; \Lambda) & \longrightarrow & H_2(M_0, \partial M_0; \Lambda) & \longrightarrow & H_1(\partial M_0; \Lambda) & \longrightarrow & 0 \\ \downarrow F & & \downarrow & & \downarrow \tilde{f}_* & & \\ H_2(M_1; \Lambda) & \longrightarrow & H_2(M_1, \partial M_1; \Lambda) & \longrightarrow & H_1(\partial M_1; \Lambda) & \longrightarrow & 0. \end{array}$$

Using the isomorphisms $H_2(M_i, \partial M_i; \Lambda) \cong H^2(M_i; \Lambda) \cong H_2(M_i; \Lambda)^*$ and choosing bases as in Remark 3.3, we see that the map induced by Φ on $H_2(M_0, \partial M_0; \Lambda)$ is given by F^{-*} . The commutativity of the diagram shows that (f, F) is a compatible pair, as desired. \square

The next result shows that the existence of a compatible pair (f, F) imposes strong restrictions on the topology of $M := M_0 \cup_f -M_1$. This is Theorem 1.12 from the introduction.

Theorem 3.12. *Let (f, F) be a compatible pair. If M_0 and M_1 are not spin, assume that the Kirby-Siebenmann invariants satisfy $\text{ks}(M_0) = \text{ks}(M_1) \in \mathbb{Z}/2$. Then there is an orientation-preserving homeomorphism*

$$M := M_0 \cup_f -M_1 \cong S^1 \times S^3 \#_{i=1}^a S^2 \times S^2 \#_{j=1}^b S^2 \tilde{\times} S^2$$

for some a, b with $a + b = b_2(M_0)$. If M_0 and M_1 are spin then $b = 0$.

Moreover it can be assumed that the Λ -isometry induced by this homeomorphism takes the Lagrangian $\widehat{\lambda}_M^{-1}(\Gamma_{F^{-*}}) \subseteq H_2(M; \Lambda)$ to the Lagrangian generated by $\{[\{\text{pt}\} \times S^2]\}_{i=1}^{b_2(M_0)}$.

Proof. Lemma 3.8 establishes that $\pi_1(M) = \mathbb{Z}$. Proposition 3.9 proves that λ_M is isometric to $\lambda_{M_0} \cup_{\tilde{f}_*} -\lambda_{M_1}$. We have $\lambda_{M_0} \cup_{\tilde{f}_*} -\lambda_{M_1} = \lambda_{M_0} \cup_{\partial F} -\lambda_{M_1}$ because the pair (f, F) is compatible. It follows from Proposition 2.13 that $\lambda_{M_0} \cup_{\partial F} -\lambda_{M_1}$ is metabolic with $\Gamma_{F^{-*}}$ as a Lagrangian. We deduce that M is a closed 4-manifold with $\pi_1(M) = \mathbb{Z}$ and metabolic Λ -intersection form. The (nonsingular) intersection form of M (which can be obtained from λ_M by setting $t = 1$) is therefore isometric to

$$\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}^{\oplus a} \oplus \begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix}^{\oplus b}$$

for some a, b . As $\text{ks}(M_0) = \text{ks}(M_1)$, by additivity of the Kirby-Siebenmann invariant [FNOP19, Theorem 8.2] under gluing it follows that $\text{ks}(M) = 0$. Apart from the statement that $b = 0$ for M_i spin, the theorem now follows from the classification of closed 4-manifolds with infinite cyclic fundamental group due to [FQ90, Theorem 10.7A (2)] and [SW00]: every isometry between the Λ -intersection forms of two closed, oriented 4-manifolds with fundamental group \mathbb{Z} can be realised by a homeomorphism.

Now assume that M_0 and M_1 are spin. To show that $b = 0$, it is sufficient to show that λ_M is hyperbolic, as the result will then once again follow from [FQ90, Theorem 10.7A (2)] and [SW00]. We already argued that λ_M is isometric to $\lambda_{M_0} \cup_{\partial F} -\lambda_{M_1}$ because the pair (f, F) is compatible. Applying Proposition 2.12 to the isometry $h = \partial F$, this latter form is isometric to $\lambda_{M_1} \cup_{\text{Id}} -\lambda_{M_1}$. The identity certainly belongs to $\text{Homeo}_{\varphi}^+(\partial M_1, \partial M_1)$ and so a second application of Proposition 3.9 ensures that $\lambda_{M_1} \cup_{\text{Id}} -\lambda_{M_1}$ is isometric to $\lambda_{M_1 \cup_{\text{Id}} -M_1}$. Summarising, we have the following sequence of isometries:

$$(6) \quad \lambda_M \cong \lambda_{M_0 \cup_{\tilde{f}_*} -M_1} \cong \lambda_{M_0} \cup_{\partial F} -\lambda_{M_1} \cong \lambda_{M_1} \cup_{\text{Id}} -\lambda_{M_1} \cong \lambda_{M_1 \cup_{\text{Id}} -M_1}.$$

Since M_1 is spin, the double $DM_1 := M_1 \cup_{\text{Id}} -M_1$ is also spin. As $\pi_2(DM_1) = H_2(DM_1; \Lambda)$ is free (by the third item of Lemma 3.2 and Lemma 3.8) and DM_1 is spin, it follows from [KPT20,

Remark 1.7] that λ_{DM_1} is even, meaning that $\lambda_{DM_1} = q + \bar{q}$ for some sesquilinear form q . Since λ_{DM_1} and λ_M are isometric by (6), it follows that λ_M is even. Now since λ_M is an even metabolic nonsingular Hermitian form on a finitely generated free Λ -module, it is hyperbolic by [Knu91, Corollary 3.7.3] with Γ_{F-*} as a Lagrangian. \square

We make the Lagrangian $\widehat{\lambda}_M^{-1}(\Gamma_{F-*}) \subseteq H_2(M; \Lambda)$ somewhat more concrete. In the setting of Theorem 3.12, combine the inclusion induced maps as

$$\iota = (\iota_0 \quad \iota_1) : H_2(M_0; \Lambda) \oplus H_2(M_1; \Lambda) \rightarrow H_2(M; \Lambda).$$

We record the following fact; compare with [Boy86, Proposition 4.2] in the simply connected case.

Lemma 3.13. *The Lagrangian $\widehat{\lambda}_M^{-1}(\Gamma_{F-*}) \subseteq H_2(M; \Lambda)$ contains the graph Γ_{-F} of the isometry $-F : H_2(M_0; \Lambda) \rightarrow H_2(M_1; \Lambda)$.*

Proof. Given $x_0 \in H_2(M_0; \Lambda)$, we have

$$\iota \begin{pmatrix} x_0 \\ -F(x_0) \end{pmatrix} = \widehat{\lambda}_M^{-1} \begin{pmatrix} \widehat{\lambda}_{M_0}(x_0) \\ \widehat{\lambda}_{M_1}(F(x_0)) \end{pmatrix} = \widehat{\lambda}_M^{-1} \begin{pmatrix} \widehat{\lambda}_0(x_0) \\ F^{-*}(\widehat{\lambda}_{M_0}(x_0)) \end{pmatrix}.$$

Here, the first equality holds by the commutativity of the diagram in (4), while the second holds because F is an isometry. It follows that $\Gamma_{-F} \subseteq \widehat{\lambda}_M^{-1}(\Gamma_{F-*})$ as desired. \square

We introduce some notation needed to provide a sufficient criterion to produce compatible pairs. Under the identification $\text{Aut}(\partial\lambda_{M_1}) = \text{Aut}(\text{Bl}_{\partial M_1})$ as in Remark 3.6, there is a left action

$$\begin{aligned} (\text{Homeo}_\varphi^+(\partial M_1) \times \text{Aut}(\lambda_{M_1})) \times \text{Aut}(\text{Bl}_{\partial M_1}) &\rightarrow \text{Aut}(\text{Bl}_{\partial M_1}) \\ (f, F) \cdot h &:= \widetilde{f}_* \circ h \circ \partial F^{-1}. \end{aligned}$$

The next proposition gives a criterion to find a compatible pair.

Proposition 3.14. *For M_0 and M_1 as above, if there is $f \in \text{Homeo}_\varphi^+(\partial M_0, \partial M_1)$ and $F \in \text{Iso}(\lambda_{M_0}, \lambda_{M_1})$, and if the orbit set*

$$\text{Aut}(\text{Bl}_{\partial M_1}) / \text{Homeo}_\varphi^+(\partial M_1) \times \text{Aut}(\lambda_{M_1})$$

is trivial, then a compatible pair (f', F') exists.

Proof. The composition $\widetilde{f}_* \circ \partial F^{-1}$ is an isometry of $\partial\lambda_{M_1} \cong \text{Bl}_{\partial M_1}$. Using the assumption, there exists a homeomorphism $g \in \text{Homeo}_\varphi^+(\partial M_1)$ and an isometry $G \in \text{Aut}(\lambda_{M_1})$ such that we have $\widetilde{f}_* \circ \partial F^{-1} = \widetilde{g}_*^{-1} \circ \partial G$. Equivalently, we can write $(\widetilde{g} \circ f)_* = \partial(G \circ F)$ and consequently the pair $(f', F') := (g \circ f, G \circ F)$ is compatible. \square

4. PARTIAL CLASSIFICATION OF COMPACT 4-MANIFOLDS WITH FUNDAMENTAL GROUP \mathbb{Z}

In this section, we prove Theorem 1.10 from the introduction. The proof of this result was inspired by [Boy86] and [Kre99].

Theorem 4.1. *Let M_0 and M_1 be two 4-manifolds with infinite cyclic fundamental group whose boundaries are ribbon and have torsion Alexander modules. Let $f \in \text{Homeo}_\varphi^+(\partial M_0, \partial M_1)$ be an orientation-preserving homeomorphism and let $F \in \text{Iso}(\lambda_{M_0}, \lambda_{M_1})$ be an isometry. If M_0 and M_1 are not spin, assume that the Kirby-Siebenmann invariants satisfy $\text{ks}(M_0) = \text{ks}(M_1) \in \mathbb{Z}/2$. Then the following assertions are equivalent:*

- (1) the pair (f, F) is compatible;
(2) the homeomorphism f extends to an orientation-preserving homeomorphism

$$\Phi: M_0 \xrightarrow{\cong} M_1$$

inducing the given isometry $F: H_2(M_0; \Lambda) \cong H_2(M_1; \Lambda)$.

Proof. Proposition 3.11 shows that (2) implies (1). To prove the converse, given a compatible pair (f, F) , we must therefore establish the existence of a homeomorphism Φ . Under this assumption, and in the non-spin case assuming that the Kirby-Siebenmann invariants coincide, we saw in Theorem 3.12 that for some a, b such that $a+b = b_2(M_0)$, there is a homeomorphism

$$M_0 \cup_f -M_1 \cong S^1 \times S^3 \#_{i=1}^a S^2 \times S^2 \#_{i=1}^b S^2 \widetilde{\times} S^2.$$

Define $M := M_0 \cup_f -M_1$. With this notation, Theorem 3.12 additionally states that this homeomorphism takes the Lagrangian $\widehat{\lambda}_M^{-1}(\Gamma_{F-*}) \subseteq H_2(M; \Lambda)$ to the Lagrangian generated by the spheres $\{[\text{pt}] \times S^2\}_{i=1}^{b_2(M_0)}$. Consider the following 5-dimensional null-bordism for M

$$W := S^1 \times D^4 \natural_{i=1}^a S^2 \times D^3 \natural_{i=1}^b S^2 \widetilde{\times} D^3,$$

where $S^2 \widetilde{\times} D^3$ denotes the twisted linear D^3 bundle over S^2 . We prove that W is an h -cobordism; it will then automatically be an s -cobordism as $\text{Wh}(\mathbb{Z}) = 0$. As $H_i(W, M_0; \Lambda) = 0$ for $i \neq 2, 3$, we must show that $H_2(W, M_0; \Lambda) = 0$ for $i = 2, 3$.

Use $D^3_{\frac{1}{2}} \subseteq D^3$ to denote a 3-ball of smaller radius inside D^3 . Choosing the obvious set of generators for $H_2(W; \Lambda)$ and representing them by embedded spheres, we obtain

$$U := \natural_{i=1}^a S^2 \times D^3_{\frac{1}{2}} \natural_{i=1}^b S^2 \widetilde{\times} D^3_{\frac{1}{2}} \subseteq W.$$

Note that $H_*(U; \Lambda) = H_*(W; \Lambda)$ and $H_*(\partial U; \Lambda) = H_*(M; \Lambda)$. In particular, under this identification, we have

$$(7) \quad \ker(H_2(M; \Lambda) \xrightarrow{j} H_2(W; \Lambda)) = \text{im}(H_3(U, \partial U; \Lambda) \rightarrow H_2(\partial U; \Lambda)) = \widehat{\lambda}_M^{-1}(\Gamma_{F-*}).$$

In what follows, we identify $H_3(U, \partial U; \Lambda)$ with its image in $H_2(\partial U; \Lambda)$ allowing ourselves for instance to write $H_3(U, \partial U; \Lambda) \subseteq H_2(\partial U; \Lambda)$. We also think of the adjoint $\widehat{\lambda}_M$ of the intersection form on M as having $H_2(\partial U; \Lambda)$ as its domain.

Claim 1. The connecting homomorphism

$$\partial_0: H_3(W \setminus \overset{\circ}{U}, M_0 \sqcup \partial U; \Lambda) \rightarrow H_2(M_0; \Lambda),$$

arising from the long exact sequence of the triple $(W \setminus \overset{\circ}{U}, M_0 \sqcup \partial U, \partial U)$, is an isomorphism.

Proof of Claim 1. It suffices to note that by excision and by our choice of U and W , we have $H_i(W \setminus \overset{\circ}{U}, \partial U; \Lambda) \cong H_i(W, U; \Lambda) = 0$ for $i = 2, 3$. \square

Claim 2. The following diagram commutes:

$$\begin{array}{ccccc} H_2(M_0; \Lambda) & \xleftarrow{\widehat{\lambda}_0} & ((H_2(M_0; \Lambda) \cup_{\partial F} H_2(M_1; \Lambda), \lambda_0 \cup_{\partial F} -\lambda_1) & \xleftarrow{\quad} & \Gamma_{F-*} \\ \cong \uparrow \partial_0 & & \cong \uparrow \widehat{\lambda}_M & & \cong \uparrow \widehat{\lambda}_M \\ H_3(W \setminus \overset{\circ}{U}, M_0 \sqcup \partial U; \Lambda) & \xleftarrow{\quad} & (H_2(\partial U; \Lambda), \lambda_{\partial U}) & \xleftarrow{\quad} & H_3(U, \partial U; \Lambda). \end{array}$$

Proof of Claim 2. The commutativity of the right square follows from (7), and so we focus on the left square. Consider the long exact sequence of the triples $(W \setminus \mathring{U}, M_0 \sqcup \partial U, \partial U)$ and $(W \setminus \mathring{U}, M_0 \sqcup \partial U, M_0)$. The portions of interest can be seen in the two horizontal rows of the following diagram, where Λ coefficients are understood:

(8)

$$\begin{array}{ccccccc}
H_3(W \setminus \mathring{U}, \partial U) = 0 & \rightarrow & H_3(W \setminus \mathring{U}, M_0 \sqcup \partial U) & \xrightarrow{\partial_0} & H_2(M_0) & \xrightarrow{\quad} & 0 = H_2(W \setminus \mathring{U}, \partial U) \\
& & \uparrow = & & \downarrow \iota_0 & \nearrow \hat{\lambda}_0 & \\
& & & & H_2(M) & \xrightarrow{\hat{\lambda}_M, \cong} & H_2(M_0) \cup_{\partial F} H_2(M_1) \\
& & & & \uparrow = & \nwarrow \hat{\lambda}_M, \cong & \\
H_3(W \setminus \mathring{U}, M_0) = 0 & \rightarrow & H_3(W \setminus \mathring{U}, M_0 \sqcup \partial U) & \rightarrow & H_2(\partial U) & \xrightarrow{\quad} & H_2(W \setminus \mathring{U}, M_0).
\end{array}$$

The claim will follow from the central portion of the diagram, once we explain all its features and its commutativity. The zeros in the first row are a consequence of Claim 1. The bottom leftmost zero is stated in Kreck's work [Kre99, page 734] but we outline a proof in this setting. The exact sequence of the triple $(W \setminus \mathring{U}, M_1 \sqcup \partial U, M_1)$, together with a Mayer-Vietoris argument, give rise to an isomorphism $H_1(W \setminus \mathring{U}, M_1 \sqcup \partial U; \Lambda) \cong H_0(\partial U; \Lambda) = \Lambda$. Similarly, using the long exact sequence of the triple $(W \setminus \mathring{U}, M_1 \sqcup \partial U, \partial U)$, one can deduce that $H_2(W \setminus \mathring{U}, M_1 \sqcup \partial U; \Lambda) = 0$. Since we also have $H_0(W \setminus \mathring{U}, M_1 \sqcup \partial U; \Lambda) = 0$, Poincaré duality and the UCSS imply that

$$H_3(W \setminus \mathring{U}, M_0; \Lambda) = H^2(W \setminus \mathring{U}, M_1 \sqcup \partial U; \Lambda) = 0.$$

To establish the claim, it only remains to show that the diagram in (8) is commutative. The middle square clearly commutes. Proposition 3.9 establishes the commutativity of the triangles but with $H_2(M_0) \cup_{\tilde{f}_*} H_2(M_1)$ in place of $H_2(M_0) \cup_{\partial F} H_2(M_1)$. However, since (f, F) is a compatible pair, these two modules are isomorphic. This concludes our explanation of the diagram in (8) and concludes the proof of Claim 2. \square

Write A and B for the images of $H_3(U, \partial U; \Lambda)$ and $H_3(W \setminus \mathring{U}; M_0 \sqcup \partial U; \Lambda)$ in $H_2(\partial U; \Lambda)$. As Proposition 2.13 implies that $H_2(M_0; \Lambda) \cup_{\partial F} H_2(M_1; \Lambda) = \Gamma_{F-*} \oplus H_2(M_0; \Lambda)$, we deduce from Claim 2 that

$$H_2(\partial U; \Lambda) = A \oplus B.$$

Assemble the long exact sequence of the triple $(W \setminus \mathring{U}, M_0 \sqcup \partial U, M_0)$ and the long exact sequence of the triple $(W, W \setminus \mathring{U}, M_0)$ into the following diagram, based on that from [Kre99, p. 738], in which Λ -coefficients are understood:

$$\begin{array}{ccccccc}
& & & & 0 & & \\
& & & & \downarrow & & \\
& & & & B \cong H_3(W \setminus \mathring{U}, M_0 \sqcup \partial U) & & \\
& & & & \downarrow & & \\
A \cong H_3(U, \partial U) & \longrightarrow & & \longrightarrow & H_2(\partial U) & & \\
& & \downarrow \cong & & \downarrow & & \\
0 \longrightarrow & H_3(W, M_0) \longrightarrow & H_3(W, W \setminus \mathring{U}) & \longrightarrow & H_2(W \setminus \mathring{U}, M_0) & \longrightarrow & H_2(W, M_0) \longrightarrow 0. \\
& & & & \downarrow & & \\
& & & & 0 & &
\end{array}$$

Here, the left vertical isomorphism comes from excision. Since $H_2(\partial U; \Lambda) = A \oplus B$, the right-down composition $A \rightarrow H_2(W \setminus \hat{U}, M_0; \Lambda)$ is an isomorphism. It follows that the central map in the long row is an isomorphism, and therefore $H_i(W, M_0; \Lambda) = 0$ for $i = 2, 3$. Thus W is a relative h -cobordism, and therefore an s -cobordism, as desired.

Since \mathbb{Z} is a good group, by the topological s -cobordism theorem [FQ90, Theorem 7.1A], M_0 and M_1 are homeomorphic, via a homeomorphism Φ that extends f . It remains to show that Φ induces the isometry F on $H_2(-; \Lambda)$. The inclusions of M_0, M_1 into $M = M_0 \cup_f -M_1 \subseteq W$ give rise to the following homomorphism:

$$H_2(M_0) \oplus H_2(M_1) \xrightarrow{\iota=(\iota_0 \ \iota_1)} H_2(M) \xrightarrow{j} H_2(W).$$

Set $j_i = j \circ \iota_i$; these are isomorphisms because W is an s -cobordism. By definition $j_1^{-1}j_0$ is the isometry induced by the homeomorphism Φ . We noted that $H_2(U; \Lambda) = H_2(W; \Lambda)$ as well as $H_2(\partial U; \Lambda) = H_2(M; \Lambda)$. We also noted in (7) that $\ker(j) = \widehat{\lambda}_M^{-1}(\Gamma_{F^*})$. Using Lemma 3.13, we deduce that $\iota(\Gamma_{-F}) \subseteq \ker(j)$. This inclusion implies that for all $x_0 \in H_2(M_0; \Lambda)$,

$$0 = j\iota \begin{pmatrix} x_0 \\ -F(x_0) \end{pmatrix} = j\iota_0(x_0) - j\iota_1(F(x_0)) = j_0(x_0) - j_1(F(x_0)).$$

Since the j_i are isomorphisms, we obtain the desired conclusion: $j_1^{-1}j_0(x_0) = F(x_0)$. This concludes the proof of Theorem 4.1. \square

5. KNOTTED SURFACES IN SIMPLY-CONNECTED 4-MANIFOLDS.

The goal of this section is to use Theorem 4.1 to prove Theorems 1.3 and 1.4 from the introduction. Recall from our conventions that X refers to a closed simply-connected 4-manifold, while $N = X \setminus \mathring{D}^4$, and Σ always denotes a \mathbb{Z} -surface of genus g either embedded in X or properly embedded in N (recall that a \mathbb{Z} -surface refers to a surface whose knot group is \mathbb{Z} .)

This section is organised as follows. In Subsection 5.1, we collect some initial facts concerning \mathbb{Z} -surfaces. Subsection 5.2 then shows that any isometry of $\text{Bl}_{\Sigma_{g,1} \times S^1}$ can be realised by a homeomorphism of $\Sigma_{g,1} \times S^1$ (allowing us to construct a compatible pair between \mathbb{Z} -surfaces exteriors N_{Σ_0} and N_{Σ_1}). In Subsections 5.3 and 5.4, we prove Theorems 1.3 and 1.4 from the introduction respectively.

5.1. Facts about \mathbb{Z} -surfaces.

Lemma 5.1. *If $\Sigma \subseteq N$ is a \mathbb{Z} -surface, then it is null-homologous.*

Proof. We must show that $[\Sigma, \partial\Sigma] = 0 \in H_2(N, \partial N)$. The intersection form Q_N pairs $H_2(N, \partial N)$ with $H_2(N)$ nonsingularly because N is simply-connected. Thus, it is equivalent to show that $Q_N([\Sigma, \partial\Sigma], x) = 0$ for every class $x \in H_2(N)$. Represent such an $x \in H_2(N)$ by a closed surface $S \subseteq N$ that intersects Σ transversely in points p_1, \dots, p_n , so that $Q_N([\Sigma, \partial\Sigma], x) = \sum_{i=1}^n \varepsilon(p_i)$, where $\varepsilon(p_i) = \pm 1$. Now the intersection $S \cap N_\Sigma$ is a properly embedded surface in N_Σ with oriented boundary (homologous to) $\sum_{i=1}^n \varepsilon(p_i)\mu_\Sigma$. This implies that $\sum_{i=1}^n \varepsilon(p_i)\mu_\Sigma = 0 \in H_1(N_\Sigma)$. But now, since the homology group $H_1(N_\Sigma) = \pi_1(N_\Sigma) = \mathbb{Z}\mu_\Sigma$ is torsion-free, we therefore deduce that $Q_N([\Sigma, \partial\Sigma], x) = \sum_{i=1}^n \varepsilon(p_i) = 0$, establishing that Σ is null-homologous. \square

It follows easily that a closed \mathbb{Z} -surface $\Sigma \subseteq X$ is also null-homologous in $H_2(X)$. For the next lemma, recall that orientable 2-plane bundles over compact, orientable surfaces are classified up to isomorphism by their (relative) Euler number.

Lemma 5.2. *Every \mathbb{Z} -surface has trivial normal bundle. In the case of nonempty boundary this holds moreover relative to the Seifert framing on the boundary.*

Proof. In the case that the boundary is nonempty, the surface is oriented and therefore the normal bundle is trivial. We need to show that this holds relative to the Seifert framing, and that the normal bundle is likewise trivial in the closed case. Cap off the surface with a Seifert surface in S^3 . Form a push-off by pushing the Seifert surface off itself in S^3 and extending this to a normal push-off in N . Now the argument is the same as in the closed case: by the proof of Lemma 5.1, the push-off intersects Σ algebraically trivially. Therefore the normal bundle of Σ has vanishing Euler number, relative to the Seifert framing in the case of nonempty boundary. In particular the normal bundle is trivial in the closed case. \square

The boundary of the exterior of N_Σ is homeomorphic to $M_{K,g} := E_K \cup_{\partial} (\Sigma_{g,1} \times S^1)$, where $\Sigma_{g,1}$ is the orientable genus g surface with one boundary component and $E_K := S^3 \setminus \nu(K)$. We give more details on this identification as we will then make it implicitly throughout the remainder of the paper.

Remark 5.3. If $\Sigma \subseteq X$ is a closed orientable surface with $\pi_1(X_\Sigma) = \mathbb{Z}$, then the homotopy class of the meridian of $\Sigma \subseteq X$ is the unique nontrivial primitive class in $\pi_1(\partial X_\Sigma)$ that bounds a disc in $\Sigma \times D^2$. Fix a framing of the normal bundle of Σ , i.e. an identification $\nu\Sigma \cong \Sigma \times \mathbb{R}^2$, with the property that for each simple closed curve $\gamma_k \subseteq \Sigma$, we have that $\gamma_k \times \{e_1\} \subseteq X \setminus \Sigma$ is null-homologous in $H_1(X \setminus \nu\Sigma) \cong \mathbb{Z}$. Use a choice of an identification $\Sigma \cong \Sigma_g$ and this condition to fix an identification of the boundary of the exterior with $\Sigma_g \times S^1$. Any two choices now differ by an element of the mapping class group of Σ_g . Similarly, if $\Sigma \subseteq N$ is a properly embedded orientable surface, then we can identify ∂N_Σ with $E_K \cup_{\partial} (\Sigma_{g,1} \times S^1)$.

Lemma 5.4. *The exterior of a \mathbb{Z} -surface Σ has ribbon boundary.*

Proof. Since N is simply-connected, we have

$$\{1\} = \pi_1(N_\Sigma) *_{\pi_1(\Sigma \times S^1)} \pi_1(\Sigma \times D^2),$$

which implies that $\pi_1(\Sigma \times S^1) \rightarrow \pi_1(N_\Sigma)$ is onto. In fact, by the parametrisation of $\Sigma \times S^1$ described in Remark 5.3, we have that $\pi_1(\Sigma)$ maps trivially to $\pi_1(N_\Sigma)$, so the fundamental group of N_Σ is generated by a meridian of Σ . The closed case follows, since the exterior of a closed surface in X can be thought of as the exterior of a surface in $N = X \setminus \mathring{D}^4$. \square

Lemma 5.5. *The boundary of the exterior of a \mathbb{Z} -surface Σ has Λ -torsion Alexander module. More precisely, the following hold.*

(1) *If $\Sigma \subseteq X$ is closed, then*

$$H_1(\partial X_\Sigma; \Lambda) = H_1(\Sigma_g \times S^1; \Lambda) = (\Lambda/(t-1))^{\oplus 2g}.$$

(2) *If $\Sigma \subseteq N$ has boundary a knot K , then the inclusion induces an isomorphism*

$$H_1(E_K; \Lambda) \oplus H_1(\Sigma_{g,1} \times S^1; \Lambda) \cong H_1(\partial N_\Sigma; \Lambda).$$

In particular, if K has Alexander polynomial 1, then $H_1(\partial N_\Sigma; \Lambda) = (\Lambda/(t-1))^{\oplus 2g}$.

Proof. Using infinite cyclic covers, the assertion in the closed case is immediate:

$$H_1(\partial X_\Sigma; \Lambda) = H_1(\Sigma_g \times S^1; \Lambda) = H_1(\widetilde{\Sigma_g \times S^1}; \mathbb{Z}) = H_1(\Sigma_g \times \mathbb{R}) = \mathbb{Z}^{2g} = (\Lambda/(t-1))^{\oplus 2g}.$$

We prove the second assertion. Consider the Mayer-Vietoris sequence for the decomposition $M_{K,g} = E_K \cup_{\partial} (\Sigma_{g,1} \times S^1)$ with Λ coefficients:

$$\cdots \rightarrow H_1(S^1 \times S^1; \Lambda) \rightarrow H_1(E_K; \Lambda) \oplus H_1(\Sigma_{g,1} \times S^1; \Lambda) \rightarrow H_1(M_{K,g}; \Lambda) \rightarrow 0.$$

We have $H_1(S^1 \times S^1; \Lambda) = H_1(S^1 \times \mathbb{R}) = \mathbb{Z}$ and $H_1(\Sigma_{g,1} \times S^1; \Lambda) = H_1(\Sigma_{g,1} \times \mathbb{R}) = \mathbb{Z}^{2g}$. The map $H_1(S^1 \times S^1; \Lambda) \rightarrow H_1(E_K; \Lambda)$ is the zero map: it sends the generator to the lift of the longitude of E_K ; this is null-homologous in the infinite cyclic cover E_K^∞ of E_K . Next, the homomorphism $H_1(S^1 \times S^1; \Lambda) \rightarrow H_1(\Sigma_{g,1} \times S^1; \Lambda)$ is also the zero map: it coincides with the map $H_1(S^1) \rightarrow H_1(\Sigma_{g,1})$ sending the generator to $[\partial \Sigma_{g,1}]$ which vanishes in $H_1(\Sigma_{g,1})$. This establishes the second assertion. \square

5.2. Isometries of the Blanchfield pairing. Given an orientable genus g surface $\Sigma_{g,1}$ with one boundary component, we determine the isometries of the Blanchfield pairing on $\Sigma_{g,1} \times S^1$ and show that they can be realised by orientation-preserving homeomorphisms. We then describe the Blanchfield pairing of

$$M_{K,g} := E_K \cup (\Sigma_{g,1} \times S^1),$$

where $E_K := S^3 \setminus \nu(K)$ is the knot exterior.

To describe the Blanchfield pairing $\text{Bl}_{\Sigma_{g,1} \times S^1}$, we fix some notation on the symplectic group:

$$J_g := \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}^{\oplus g},$$

$$\text{Sp}_{2g}(\mathbb{Z}) := \{A \in M_{2g \times 2g}(\mathbb{Z}) \mid A^T J_g A = J_g\}.$$

Given a genus g surface $\Sigma_{g,1}$ with one boundary component, we call a basis of $H_1(\Sigma_{g,1})$ *symplectic* if the intersection form on $\Sigma_{g,1}$ with respect to this basis is represented by J_g . We also consider the map $\varphi: \pi_1(\Sigma_{g,1} \times S^1) \rightarrow \mathbb{Z}$ induced by the projection on the second coordinate and recall that

$$\text{Homeo}_\varphi^+(\Sigma_{g,1} \times S^1) = \{f \in \text{Homeo}^+(\Sigma_{g,1} \times S^1) \mid f_* \circ \varphi = \varphi\}.$$

The next proposition identifies the isometries of $\text{Bl}_{\Sigma_{g,1} \times S^1}$ with $\text{Sp}_{2g}(\mathbb{Z})$ and uses this fact to show that the orbit set $\text{Aut}(\text{Bl}_{\Sigma_{g,1} \times S^1}) / \text{Homeo}_\varphi^+(\Sigma_{g,1} \times S^1)$ is trivial.

Proposition 5.6. *A choice of a symplectic basis for $H_1(\Sigma_{g,1})$ gives rise to an identification*

$$\text{Aut}(\text{Bl}_{\Sigma_{g,1} \times S^1}) = \text{Sp}_{2g}(\mathbb{Z}).$$

Every element of $\text{Aut}(\text{Bl}_{\Sigma_{g,1} \times S^1})$ can be realised by a homeomorphism in $\text{Homeo}_\varphi^+(\Sigma_{g,1} \times S^1)$:

$$\text{Aut}(\text{Bl}_{\Sigma_{g,1} \times S^1}) / \text{Homeo}_\varphi^+(\Sigma_{g,1} \times S^1) = \{\text{Id}\}.$$

Any such homeomorphism can be assumed to be of the form $j \times \text{Id}_{S^1}$, for some $j \in \text{Homeo}^+(\Sigma_{g,1})$ that fixes the boundary of $\Sigma_{g,1}$ pointwise.

Proof. Let h be an isometry of the Blanchfield pairing $\text{Bl}_{\Sigma_{g,1} \times S^1}$. Since we saw in Lemma 5.5 that $H_1(\Sigma_{g,1} \times S^1; \Lambda) = H_1(\Sigma_{g,1})$, the isometry h can be thought of as a \mathbb{Z} -linear map. We assert that it preserves the intersection form on the surface $\Sigma_{g,1}$. Pick a symplectic basis for $H_1(\Sigma_{g,1})$ so that its intersection form is represented by J_g . By [FP17, Corollary 1.2], the Blanchfield form on the fibred manifold $\Sigma_{g,1} \times S^1$ is given by

$$(v, w) \mapsto v^T (t^{-1} - 1)^{-1} J_g w \in \Lambda_S / \Lambda.$$

Here, $\Lambda_S = \mathbb{Z}[t^{\pm 1}, (t-1)^{-1}]$ denotes the ring obtained from $\Lambda = \mathbb{Z}[t^{\pm 1}]$ by inverting $(t-1)$. Since h can be thought of as a \mathbb{Z} -linear map, we represent it by a matrix H with coefficients in \mathbb{Z} . As h is an isometry, we have the following equalities of matrices with coefficients in Λ_S / Λ :

$$(t^{-1} - 1)^{-1} H^T J_g H = J_g (t^{-1} - 1)^{-1}.$$

Since H and J_g takes values in \mathbb{Z} , the coefficients of the matrix $H^T J_g H - J_g$ take values in $(t-1)\Lambda$. On the other hand, $H^T J_g H - J_g$ has coefficients in \mathbb{Z} , so $H^T J_g H - J_g = 0$, and we deduce that $H^T J_g H = J_g$. This shows that H is a symplectomorphism, proving the first assertion.

We prove the second assertion. We may realise any $h \in \text{Sp}_{2g}(\mathbb{Z})$ as the map on $H_1(\Sigma_{g,1})$ induced by an orientation-preserving homeomorphism $j: \Sigma_{g,1} \rightarrow \Sigma_{g,1}$ that fixes the boundary pointwise [FM12, Section 2.1 and the discussion following Theorem 6.4]. Cross j with the identity on S^1 to obtain an orientation-preserving homeomorphism $j \times \text{Id}: \Sigma_{g,1} \times S^1 \rightarrow \Sigma_{g,1} \times S^1$ that belongs to $\text{Homeo}_\varphi^+(\Sigma_{g,1} \times S^1)$. \square

The next proposition describes the Blanchfield pairing of $M_{K,g} = E_K \cup (\Sigma_{g,1} \times S^1)$ and its automorphism group; here recall that $E_K = S^3 \setminus \nu(K)$ is the knot exterior.

Proposition 5.7. *Given a knot $K \subseteq S^3$, there is an isometry $\text{Bl}_{M_{K,g}} \cong \text{Bl}_{E_K} \oplus \text{Bl}_{\Sigma_{g,1} \times S^1}$ and*

$$\text{Aut}(\text{Bl}_{M_{K,g}}) \cong \text{Aut}(\text{Bl}_{E_K}) \oplus \text{Aut}(\text{Bl}_{\Sigma_{g,1} \times S^1}).$$

Proof. We proved in Lemma 5.5 that the inclusion induces a Λ -isomorphism $H_1(E_K; \Lambda) \oplus H_1(\Sigma_{g,1} \times S^1; \Lambda) \cong H_1(M_{K,g}; \Lambda)$. The isometry $\text{Bl}_{M_{K,g}} \cong \text{Bl}_{E_K} \oplus \text{Bl}_{\Sigma_{g,1} \times S^1}$ now follows from [FLNP17, Theorem 1.1].

We must now show that $\text{Aut}(\text{Bl}_{E_K} \oplus \text{Bl}_{\Sigma_{g,1} \times S^1}) = \text{Aut}(\text{Bl}_{E_K}) \oplus \text{Aut}(\text{Bl}_{\Sigma_{g,1} \times S^1})$. The order of $H_1(M_K; \Lambda)$ is the Alexander polynomial Δ_K (which is not divisible by $(t-1)$) and the order of $H_1(\Sigma_{g,1} \times S^1; \Lambda)$ is $(t-1)^{2g}$. An automorphism $h \in \text{Aut}(\text{Bl}_{E_K}) \oplus \text{Aut}(\text{Bl}_{\Sigma_{g,1} \times S^1})$ can be written as $h = \begin{pmatrix} h_{11} & h_{12} \\ h_{21} & h_{22} \end{pmatrix}$. Since the orders of $H_1(M_K; \Lambda)$ and $H_1(\Sigma_{g,1} \times S^1; \Lambda)$ are coprime, we deduce that $h_{12} = h_{21} = 0$. \square

5.3. Surfaces in manifolds with boundary. The aim of this subsection is to prove Theorem 1.3 from the introduction. Given a knot $K \subseteq S^3$, thanks to Proposition 5.7, we can write automorphisms of $\text{Bl}_{M_{K,g}}$ as $h = h_K \oplus h_\Sigma$ with $h_K \in \text{Aut}(\text{Bl}_{E_K})$ and $h_\Sigma \in \text{Aut}(\text{Bl}_{\Sigma_{g,1} \times S^1})$.

Let us recall the construction of an isotopy, as introduced in the introduction. Let $f_K: E_K \rightarrow E_K$ be an orientation-preserving homeomorphism that is the identity on ∂E_K . Extend f_K via the identity on νK to an orientation-preserving self-homeomorphism \tilde{f}_K of S^3 . The mapping class group of S^3 is trivial, so there is an isotopy $\Theta(\tilde{f}_K)_t: S^3 \rightarrow S^3$ between the extension and the identity, such that $\Theta(\tilde{f}_K)_0 = \text{Id}$ and $\Theta(\tilde{f}_K)_1 = \tilde{f}_K$. We can now prove Theorem 1.3.

Theorem 5.8. *Let $\Sigma_0, \Sigma_1 \subseteq N$ be two \mathbb{Z} -surfaces of genus g with boundary K . Suppose there is an isometry $F: \lambda_{N_{\Sigma_0}} \cong \lambda_{N_{\Sigma_1}}$, and write $\partial F = h_K \oplus h_\Sigma$.*

- If h_K is induced by an orientation-preserving homeomorphism $f_K: E_K \rightarrow E_K$ that is the identity on ∂E_K , then f_K extends to an orientation-preserving homeomorphism of pairs

$$(N, \Sigma_0) \xrightarrow{\cong} (N, \Sigma_1)$$

inducing the given isometry $F: H_2(N_{\Sigma_0}; \Lambda) \cong H_2(N_{\Sigma_1}; \Lambda)$.

- If in addition $N = D^4$, then for any choice of isotopy $\Theta(f_K)$, the surfaces Σ_0 and Σ_1 are topologically ambiently isotopic via an ambient isotopy of D^4 extending $\Theta(f_K)$.

Proof. By assumption, the boundary of the isometry F is $\partial F = h_K \oplus h_\Sigma$ for some $h_K \in \text{Aut}(\text{Bl}_{E_K})$ and some $h_\Sigma \in \text{Aut}(\text{Bl}_{\Sigma_{g,1} \times S^1})$. By Proposition 5.6, we can realise h_Σ by an orientation-preserving homeomorphism of the form $j \times \text{Id}_{S^1}$ with $j: \Sigma_{g,1} \rightarrow \Sigma_{g,1}$ an orientation-preserving homeomorphism that fixes the boundary pointwise. Moreover, h_K is realised by an orientation-preserving homeomorphism $f_K: E_K \rightarrow E_K$ that fixes the boundary ∂E_K . Gluing these two homeomorphisms together, we obtain a homeomorphism

$$f := f_K \cup (j \times \text{Id}_{S^1}): M_{K,g} \rightarrow M_{K,g}.$$

Since f is the identity on $\partial E_K \subseteq M_{K,g}$, we deduce that $f \in \text{Homeo}_\varphi^+(M_{K,g})$. It follows that (f, F) is a compatible pair. The N_{Σ_i} are 4-manifolds whose boundaries are ribbon (recall Lemma 5.4) and have torsion Alexander modules for $i = 0, 1$ (by Lemma 5.5). Since $\Sigma_i \times D^2$ is a smooth manifold, $\text{ks}(\Sigma_i \times D^2) = 0$. Thus for $i = 0, 1$, by additivity of ks under gluing we have $\text{ks}(N_{\Sigma_i}) = \text{ks}(N_{\Sigma_i}) + \text{ks}(\Sigma_i \times D^2) = \text{ks}(N)$. In particular $\text{ks}(N_{\Sigma_0}) = \text{ks}(N_{\Sigma_1})$. We may therefore apply Theorem 4.1 to extend f to an orientation-preserving homeomorphism $\Phi: N_{\Sigma_0} \rightarrow N_{\Sigma_1}$ that induces F . As a consequence, we obtain the required homeomorphism of pairs:

$$\Phi' := \Phi \cup (j \times \text{Id}_{D^2}): (N, \Sigma_0) \rightarrow (N, \Sigma_1).$$

This establishes the first assertion.

For the second assertion, let $N = D^4$ and let $\Theta(f_K)_t: S^3 \rightarrow S^3$ be an isotopy of self-homeomorphisms of S^3 with $\Theta(f_K)_0 = \text{Id}_{S^3}$ and $\Theta(f_K)_1$ equal to the extension \tilde{f}_K of f_K by the identity to all of S^3 . Cone $\Theta(f_K)$ to obtain an isotopy $C(\Theta(f_K)_t): D^4 \rightarrow D^4$ with $C(\Theta(f_K)_0) = \text{Id}_{D^4}$ and $C(\Theta(f_K)_1)$ equal to the cone $C(\tilde{f}_K)$ of \tilde{f}_K .

Next, note that $C(\tilde{f}_K)$ and $\Phi': D^4 \rightarrow D^4$ are two homeomorphisms of D^4 that restrict to the same homeomorphism of S^3 on the boundary. Therefore $C(\tilde{f}_K)^{-1} \circ \Phi': D^4 \rightarrow D^4$ is a homeomorphism restricting to Id_{S^3} on the boundary. By the Alexander trick, $C(\tilde{f}_K)^{-1} \circ \Phi'$ is isotopic rel. boundary to the identity, via an isotopy $G_t: D^4 \rightarrow D^4$ with $G_0 = \text{Id}_{D^4}$ and $G_1 = C(\tilde{f}_K)^{-1} \circ \Phi'$. Note that $G_t|_{S^3} = \text{Id}_{S^3}$ for all $t \in [0, 1]$. Then let

$$H_t := C(\Theta(f_K)_t) \circ G_t: D^4 \rightarrow D^4.$$

This is an isotopy with

$$\begin{aligned} H_0 &= \text{Id}_{D^4} \circ \text{Id}_{D^4} = \text{Id}_{D^4} \\ H_1 &= C(\tilde{f}_K) \circ C(\tilde{f}_K)^{-1} \circ \Phi' = \Phi'. \end{aligned}$$

In addition, for every $t \in [0, 1]$ we have that $H_t|_{S^3} = \Theta(f_K)_t \circ G_t|_{S^3} = \Theta(f_K)_t$. Since $\text{Id}_{D^4}(\Sigma_0) = \Sigma_0$ and $\Phi'(\Sigma_0) = \Sigma_1$, the isotopy H_t is a topological ambient isotopy between Σ_0 and Σ_1 , and it extends $\Theta(f_K)$, as required. \square

We also note the following corollary, in which we give a slightly relaxed criterion, but without precise control on the isotopy of K . It is sometimes easier to construct homeomorphisms of E_K that do not fix the boundary ∂E_K , but rather only fix the basepoint, so this could be a useful variation.

Corollary 5.9. *Let $K \subseteq S^3$ be a knot and fix a basepoint in ∂E_K . Let $f: S^3 \rightarrow S^3$ be an orientation-preserving, basepoint-preserving homeomorphism with $f(K) = K$ as oriented knots, $f(E_K) = E_K$. Then f induces an isomorphism $h: H_1(E_K; \Lambda) \rightarrow H_1(E_K; \Lambda)$. Suppose that Σ_0 and Σ_1 are two \mathbb{Z} -surfaces in $N = X \setminus \mathring{D}^4$ with boundary $K \subseteq S^3 = \partial N$, and let $F: \lambda_{N_{\Sigma_0}} \cong \lambda_{N_{\Sigma_1}}$ be an isometry with $\partial F = h$. Then Σ_0 and Σ_1 are related by a homeomorphism of pairs, and if $N = D^4$ then they are ambiently isotopic.*

Proof. Since f is orientation-preserving as a map from S^3 to S^3 , and as a map $K \rightarrow f(K)$, it restricts to an orientation-preserving map on ∂E_K that preserves the homotopy class of the zero-framed longitude and therefore preserves the orientation of the meridian. Thus f commutes with the map $\pi_1(E_K) \rightarrow \mathbb{Z}$ inducing the Λ -coefficients, and therefore induces a map $h: H_1(E_K; \Lambda) \rightarrow H_1(E_K; \Lambda)$ as claimed; recall Proposition 3.7. Let $\theta_t: S^3 \rightarrow S^3$ be an isotopy from the identity of S^3 to f . Let $S^3 \times [0, 1] \hookrightarrow N$ be a collar on the boundary of N with $S^3 \times \{0\}$ mapping to ∂N . Let $\Theta_t: N \rightarrow N$ be the isotopy that is the identity outside $S^3 \times [0, 1]$ and defined by $\Theta_t(x, s) = \theta_{t(1-s)}(x)$. This performs θ_t on the boundary, and tapers it in the collar. Let $\Sigma'_0 := \Theta_1(\Sigma_0)$, and note that Σ'_0 is ambiently isotopic (not necessarily rel. boundary) to Σ_0 . In addition, $\partial(\Theta_1)_* = \partial F = h$ as isomorphisms of $H_1(E_K; \Lambda)$. Therefore $G := F \circ (\Theta_1)_*^{-1}: \lambda_{N_{\Sigma'_0}} \cong \lambda_{N_{\Sigma_1}}$ is an isometry with $\partial G = \text{Id}$. By Theorem 1.3, Σ'_0 and Σ_1 are related by a homeomorphism of pairs restricting to the identity on the boundary, and if $N = D^4$ then they are ambiently isotopic rel. boundary. The corollary now follows immediately. \square

5.4. Surfaces in closed manifolds. Now we recover the statements in the closed setting, proving Theorem 1.4 from the introduction. As discussed in the introduction, for a closed, simply-connected 4-manifold X there is a classification of self-homeomorphisms of X in terms of their action on $H_2(X)$. The hypothesis of Theorem 1.4 is in terms of an isometry between the Λ -intersection forms of the surface exteriors X_{Σ_0} and X_{Σ_1} . In order to carefully state Theorem 1.4, first we need to understand the relationship between such an isometry and isomorphisms of the second homology $H_2(X)$ of the ambient 4-manifold.

Lemma 5.10. *Let $\Sigma_0, \Sigma_1 \subseteq X$ be closed \mathbb{Z} -surfaces of the same genus.*

- (1) *An isometry $F: \lambda_{X_{\Sigma_0}} \cong \lambda_{X_{\Sigma_1}}$ induces an isomorphism $F_{\mathbb{Z}}: H_2(X) \xrightarrow{\cong} H_2(X)$.*
- (2) *If the isometry F is induced by a homeomorphism of pairs $\Phi': (X, \Sigma_0) \rightarrow (X, \Sigma_1)$, then $F_{\mathbb{Z}} = \Phi'_*$; in particular, $F_{\mathbb{Z}}$ is an isometry of the standard intersection form Q_X .*

Proof. We claim that $H_2(X_{\Sigma_i}; \Lambda) \otimes_{\Lambda} \mathbb{Z} \cong H_2(X_{\Sigma_i})$. To prove this, we use the universal coefficient spectral sequence with second page $E_{p,q}^2 = \text{Tor}_p^{\Lambda}(H_q(X_{\Sigma_i}; \Lambda), \mathbb{Z})$ and which converges to $H_*(X_{\Sigma_i})$. The claim now follows promptly from the fact that $H_1(X_{\Sigma_i}; \Lambda) = 0$ and

$$\text{Tor}_2^{\Lambda}(H_0(X_{\Sigma_i}; \Lambda), \mathbb{Z}) = \text{Tor}_2^{\mathbb{Z}[\mathbb{Z}]}(\mathbb{Z}, \mathbb{Z}) = H_2(\mathbb{Z}; \mathbb{Z}) = H_2(S^1) = 0.$$

The claim implies that the intersection form on $H_2(X_{\Sigma_i})$ is obtained from the intersection form on $H_2(X_{\Sigma_i}; \Lambda)$ by tensoring down. Therefore F induces an isometry $F \otimes_{\Lambda} \text{Id}_{\mathbb{Z}}: H_2(X_{\Sigma_0}) \rightarrow H_2(X_{\Sigma_1})$. A Mayer-Vietoris argument then yields the following commutative diagram, where

the dotted map labelled $F_{\mathbb{Z}}$ is defined as the unique homomorphism that makes the right hand square commutes (both existence and uniqueness follow from a short diagram chase):

$$(9) \quad \begin{array}{ccccccc} 0 & \longrightarrow & \mathbb{Z}^{2g} & \longrightarrow & H_2(X_{\Sigma_0}) & \xrightarrow{p_0} & H_2(X) \longrightarrow 0 \\ & & \downarrow (F \otimes_{\Lambda} \text{Id}_{\mathbb{Z}}) & & \downarrow F \otimes_{\Lambda} \text{Id}_{\mathbb{Z}} & & \downarrow F_{\mathbb{Z}} \\ 0 & \longrightarrow & \mathbb{Z}^{2g} & \longrightarrow & H_2(X_{\Sigma_1}) & \xrightarrow{p_1} & H_2(X) \longrightarrow 0. \end{array}$$

The second assertion is now immediate: the homeomorphism Φ' induces an isometry of $(H_2(X), Q_X)$ that satisfies $\Phi'_* \circ p_0 = p_1 \circ (F \otimes_{\Lambda} \text{Id}_{\mathbb{Z}})$, so $F_{\mathbb{Z}} = \Phi'_*$ is also an isometry. \square

We move on to the proof of Theorem 1.4, whose statement we recall here for the benefit of the reader.

Theorem 5.11. *Let $\Sigma_0, \Sigma_1 \subseteq X$ be two closed \mathbb{Z} -surfaces of the same genus.*

- (1) *If the intersection forms $\lambda_{X_{\Sigma_0}}$ and $\lambda_{X_{\Sigma_1}}$ are isometric via an isometry F , then there is an orientation-preserving homeomorphism of pairs*

$$\Phi: (X, \Sigma_0) \xrightarrow{\cong} (X, \Sigma_1)$$

inducing the given isometry $\Phi_ = F: H_2(X_{\Sigma_0}; \Lambda) \cong H_2(X_{\Sigma_1}; \Lambda)$.*

- (2) *The isometry F induces an isometry $F_{\mathbb{Z}}: H_2(X) \rightarrow H_2(X)$ of the standard intersection form Q_X of X by Lemma 5.10. The surfaces Σ_0 and Σ_1 are topologically ambiently isotopic if and only if $F_{\mathbb{Z}} = \text{Id}$.*

Proof. After an ambient isotopy, assume that Σ_0 and Σ_1 coincide on a disc $D^2 \subseteq \Sigma_0 \cap \Sigma_1$. Assume that the normal bundles also coincide over this D^2 . Consider the preimage $\dot{D}^2 \times \mathbb{R}^2 \subseteq \nu\Sigma_i$. This is homeomorphic to an open 4-ball \dot{D}^4 . Remove this (\dot{D}^4, \dot{D}^2) from (X, Σ_i) to obtain $(N, \tilde{\Sigma}_i)$, with $\partial\tilde{\Sigma}_i = \Sigma_i \cap \partial N$ the unknot K in S^3 . Then the exterior of Σ_i in X equals the exterior of $\tilde{\Sigma}_i$ in N .

Since $X_{\Sigma_i} = N_{\tilde{\Sigma}_i}$, the Λ -intersection forms are unchanged and the isometry F also induces an isometry $F: \lambda_{N_{\tilde{\Sigma}_0}} \cong \lambda_{N_{\tilde{\Sigma}_1}}$. Write $\partial F = h_K \oplus h_{\Sigma}$ as in Theorem 5.8. The unknot $K \subseteq S^3$ has trivial Alexander module, so $h_K = \text{Id}$ and so h_K is realised by the homeomorphism $f_K = \text{Id}: E_K \rightarrow E_K$. Theorem 5.8 provides a rel. boundary homeomorphism of pairs $\Phi': (N, \tilde{\Sigma}_0) \rightarrow (N, \tilde{\Sigma}_1)$ that induces the isometry F on the Λ -homology of the surface exteriors. We recover the required homeomorphism of pairs $\Phi: (X, \Sigma_0) \rightarrow (X, \Sigma_1)$ by gluing Φ' with the identity homeomorphism $(D^4, D^2) \rightarrow (D^4, D^2)$.

Lemma 5.10 implies that the isometry F induces an isometry $F_{\mathbb{Z}}$ of the standard intersection form Q_X . In particular, Lemma 5.10 (2) specifies that $F_{\mathbb{Z}}$ is induced by the homeomorphism Φ . If $F_{\mathbb{Z}} = \text{Id}$, then Φ is a self-homeomorphism of X inducing the identity on $H_2(X)$, so by [Qui86, Theorem 1.1], [FQ90, Theorem 10.1], Φ is isotopic to the identity. It follows that Σ_0 and Σ_1 are topologically ambiently isotopic.

On the other hand if Σ_0 and Σ_1 are topologically ambiently isotopic, then the induced homeomorphism between their exteriors extends to a homeomorphism from X to itself that is isotopic to the identity, and so certainly induces the identity map on $H_2(X)$. \square

6. EQUIVARIANT INTERSECTION FORMS OF SURFACE EXTERIORS

The goal of this section is to collect some results about the intersection forms of \mathbb{Z} -surface exteriors. The main result of Subsection 6.1 shows that up to direct summands with the

Hermitian form

$$\mathcal{H}_2 := \left(\Lambda^2, \begin{pmatrix} 0 & t-1 \\ t^{-1}-1 & 0 \end{pmatrix} \right),$$

the equivariant intersection forms of any two \mathbb{Z} -surface exteriors are isometric (Proposition 6.3). Subsection 6.2 then focuses on the case of \mathbb{Z} -surfaces $\Sigma \subseteq N = X \setminus \mathring{D}^4$ with boundary an Alexander polynomial one knot $K \subseteq S^3$. The main result is Corollary 6.6 which shows that λ_{N_Σ} becomes isometric to $Q_X \oplus \mathcal{H}_2^{\oplus g}$ after adding sufficiently many copies of \mathcal{H}_2 , where Q_X is the \mathbb{Z} -valued intersection form of X (note that $Q_X = Q_N$).

6.1. Equivariant intersection forms of \mathbb{Z} -surface exteriors. The goal of this section is to study the Λ -intersection form of \mathbb{Z} -surface exteriors up to stabilisations by \mathcal{H}_2 .

We start by describing the Λ -intersection form of unknotted surfaces in S^4 .

Lemma 6.1. *If $\Sigma \subseteq S^4$ is an unknotted oriented closed genus g surface, then the Λ -intersection form of S^4_Σ is isometric to $\mathcal{H}_2^{\oplus g}$.*

Proof. We carry out the computation for the standardly embedded genus g surface $\Sigma \subseteq S^4$. Use $U \subseteq S^3$ to denote the unknot. Slice (S^4, Σ) along an equatorial (S^3, U) to obtain a decomposition $(S^4, \Sigma) = (D^4, \mathring{\Sigma}) \cup (D^4, D)$, with $D \subseteq D^4$ a disc bounding U and $\mathring{\Sigma} \subseteq D^4$ a punctured unknotted surface in D^4 .

Note that S^4_Σ and $D^4_{\mathring{\Sigma}} := D^4 \setminus \nu\mathring{\Sigma}$ are homeomorphic, since we can assume that the removed \mathring{D}^4 lies in the regular neighbourhood $\nu\Sigma$ removed from S^4 to form S^4_Σ . It follows that $H_2(S^4_\Sigma; \Lambda) = H_2(D^4_{\mathring{\Sigma}}; \Lambda)$ and that the Λ -intersection forms agree.




FIGURE 1. Two handle diagrams for the exterior of a standardly embedded genus g surface $F \subseteq D^4$ with boundary the unknot in S^3 .

Thus it remains to compute the Λ -intersection form of the exterior D^4_F of a properly embedded unknotted surface $F \subseteq D^4$. A handle diagram with a single one handle and $2g$ two handles for D^4_F appears in the left hand side of Figure 1, produced using the formalism of [GS99, Section 6.2]. It can then be isotoped as in the right hand side of Figure 1, leading to a handle diagram for the infinite cyclic cover of D^4_F depicted in Figure 2. From this diagram, by taking the union of the cores of the 2-handles with null-homotopies of their attaching curves in the 4-ball, we obtain generators of $\pi_2(D^4_F) = H_2(D^4_F; \Lambda) = \Lambda^{2g}$. The Λ -intersection form can be computed via (equivariant) linking numbers, yielding the required result. \square

We recall the concept of a 1-handle stabilisation for a surface in a 4-manifold. The following definition was motivated by [JZ18]. Let $\Sigma \subseteq V$ be a locally flat (connected) surface embedded in a 4-manifold V . Let B be an embedding of D^4 into V such that ∂B intersects Σ transversely in a 2-component unlink L and B intersects Σ transversely in two discs D_0 and D_1 , which can




FIGURE 2. A handle diagram for the infinite cyclic cover of D_F^4 , where $F \subseteq D^4$ is an unknotted punctured surface.

be simultaneously isotoped within B to lie in ∂B . Suppose that, for $i = 0, 1$, a 3-dimensional 1-handle $D^2 \times [0, 1]$ is embedded in the interior of B such that $D^2 \times \{i\} = D_i$. The resulting 1-handle stabilisation of Σ is defined as

$$\Sigma' = (\Sigma \cap (S^4 \setminus B)) \cup_L (S^1 \times [0, 1]).$$

If $S^1 \times [0, 1]$ can be isotoped into ∂B relative to L , and if there exists a path γ in Σ between $(1, 0)$ and $(1, 1)$ in $S^1 \times [0, 1]$ such that $\gamma \cup (\{1\} \times [0, 1])$ is a null-homotopic loop in V , then we call the stabilisation *trivial*.

A detailed discussion of this construction in the locally flat setting can be found in [FNO19, Proposition 9.1]. The next result describes the effect of trivial 1-handle stabilisation on the Λ -intersection form of surface exteriors.

Lemma 6.2. *If $\Sigma \subseteq N$ is a \mathbb{Z} -surface and $\Sigma' \subseteq N$ is obtained from Σ by a trivial 1-handle stabilisation, then*

$$\lambda_{N_{\Sigma'}} \cong \lambda_{N_{\Sigma}} \oplus \mathcal{H}_2.$$

Proof. Since the 1-handle stabilisation is trivial, one can write $(N, \Sigma') = (N, \Sigma) \# (S^4, T^2)$, where $T^2 \subseteq S^4$ denotes a standardly embedded torus, and $\#$ denotes the interior connect sum. It follows that $N_{\Sigma'} = N_{\Sigma} \cup S_{T^2}^4$, where the identification takes place along thickened meridians: $\mu_{\Sigma} \times D^2$ is identified with $\mu_{T^2} \times D^2$. One thus deduces the isomorphism $\pi_1(N_{\Sigma'}) \cong \pi_1(N_{\Sigma}) = \mathbb{Z}$. As the coefficient system maps these meridians to $1 \in \mathbb{Z}$, a straightforward Mayer-Vietoris argument shows that $H_2(N_{\Sigma'}; \Lambda) = H_2(N_{\Sigma}; \Lambda) \oplus H_2(S_{T^2}^4; \Lambda)$, noting that $H_1(\mu_{\Sigma}; \Lambda) = 0$. It then follows that $\lambda_{N_{\Sigma'}} = \lambda_{N_{\Sigma}} \oplus \lambda_{S_{T^2}^4}$. The result is now a consequence of Lemma 6.1. \square

The next proposition shows that the Λ -intersection forms of any two \mathbb{Z} -surfaces exteriors become isometric after adding sufficiently many \mathcal{H}_2 summands.

Proposition 6.3. *For \mathbb{Z} -surfaces $\Sigma_0, \Sigma_1 \subseteq N$ of the same genus with boundary K , there exists an integer $n \geq 0$ and an isometry*

$$\lambda_{N_{\Sigma_0}} \oplus \mathcal{H}_2^{\oplus n} \cong \lambda_{N_{\Sigma_1}} \oplus \mathcal{H}_2^{\oplus n}.$$

Proof. Any two null-homologous surfaces in a 4-manifold can be made isotopic by enough 1-handles stabilisations to each surface [BS16, Theorem 5]. While Baykur and Sunukjian prove this result in the smooth category for surfaces in closed manifolds, it also applies in the topological category for properly embedded surfaces in 4-manifolds with boundary; see Theorem A.1 for a detailed proof in the case at hand.

We apply this to Σ_0, Σ_1 . This is possible because \mathbb{Z} -surfaces are nullhomologous by Lemma 5.1. Since $\pi_1(\partial(\nu\Sigma_i)) \rightarrow \pi_1(N_{\Sigma_i}) = \mathbb{Z}$ is surjective, all 1-handles stabilisations can be taken to be trivial [BS16, Lemma 3]. We deduce that after sufficiently many (say n) trivial 1-handles stabilisations, Σ_0 becomes isotopic to Σ_1 , also stabilised n times. Using Lemma 6.2, each such stabilisation adds an \mathcal{H}_2 -summand to the Λ -intersection form of the corresponding surface exterior, from which the result follows. \square

6.2. \mathbb{Z} -surfaces for Alexander polynomial one knots. We now restrict our attention to \mathbb{Z} -surfaces with boundary Alexander polynomial one knots.

An Alexander polynomial one knot $K \subseteq S^3$ bounds a disc $D \subseteq D^4$ with $\pi_1(D^4 \setminus D) = \mathbb{Z}$ [FQ90]. We call this disc a \mathbb{Z} -slice disc of K . When $S^3 = \partial N$ with $N = X \setminus \mathring{D}^4$, we arrange that this disc D belong to a collar neighborhood $S^3 \times [0, 1] \subseteq N$ of $S^3 = \partial N$. We use this disc to build a genus g surface with boundary K .

Definition 6.4. A genus g target surface Σ_g^t for an Alexander polynomial one knot K is an embedded surface obtained from a \mathbb{Z} -disc D of K by g trivial 1-handle stabilisations.

Although we do not need this fact in the sequel, an Alexander polynomial one knot $K \subseteq S^3$ bounds a *unique* \mathbb{Z} -disc: this follows either from [CP19, Theorem 1.2] or from Theorem 5.8 above, because $H_2(D_D^4; \Lambda) = 0$. This is why we do not keep track of the \mathbb{Z} -disc D in the notation for target surfaces.

The next lemma describes the Λ -intersection form of a target surface exterior.

Lemma 6.5. *If $D \subseteq S^3 \times [0, 1] \subseteq N$ is a \mathbb{Z} -disc, then $\lambda_{N_D} = Q_X$ and if Σ_g^t is a target surface, then $\lambda_{N_{\Sigma_g^t}} \cong Q_X \oplus \mathcal{H}_2^{\oplus g}$.*

Proof. Since D is properly embedded in a collar neighbourhood of ∂N , N_D is homeomorphic to the interior connected sum $N_D \cong D_D^4 \# X$. The exterior D_D^4 of a \mathbb{Z} -slice disc is aspherical, and therefore we have $H_i(D_D^4; \Lambda) = 0$ for $i > 0$ [CP19, Lemma 2.1]. The first assertion now follows from a straightforward Mayer-Vietoris argument. Since a target surface is obtained from a \mathbb{Z} -slice disc by g handle additions, the second assertion follows from the first and Lemma 6.2. \square

Finally, we describe the Λ -intersection form of a \mathbb{Z} -surface exterior for an Alexander polynomial one knot K .

Corollary 6.6. *If $\Sigma \subseteq N = X \setminus \mathring{D}^4$ is a genus g \mathbb{Z} -surface for an Alexander polynomial one knot K , then $\lambda_{N_\Sigma} \oplus \mathcal{H}_2^{\oplus n} \cong Q_X \oplus \mathcal{H}_2^{\oplus(n+g)}$ for some integer $n \geq 0$.*

Proof. Since K has Alexander polynomial one, it bounds a genus g target surface Σ_g^t . Proposition 6.3 ensures that $\lambda_{N_\Sigma} \oplus \mathcal{H}_2^{\oplus n}$ is isometric to $\lambda_{N_{\Sigma_g^t}} \oplus \mathcal{H}_2^{\oplus n}$ for some $n \geq 0$. The result now follows from Lemma 6.5, thanks to which $\lambda_{N_{\Sigma_g^t}} = Q_X \oplus \mathcal{H}_2^{\oplus g}$. \square

7. KNOTTED SURFACES IN S^4 AND D^4

As we outlined in Subsection 1.4, our results about surfaces in D^4 and S^4 can be deduced from Theorem 5.8 once we prove that the surface exteriors have isometric equivariant intersection forms. In Subsection 7.1, we show that if $\Sigma \subseteq D^4$ is a genus $g \geq 3$ \mathbb{Z} -surface with boundary an Alexander polynomial one knot K , then $\lambda_{D_\Sigma^4} \cong \mathcal{H}_2^{\oplus g}$. In Subsection 7.2 we then

combine this result with Theorem 5.8 to deduce our results about surfaces in D^4 and S^4 , in particular proving Theorems 1.1 and 1.2. In Subsection 7.3, we apply these results to study Seifert surfaces pushed into D^4 .

7.1. Intersection forms in D^4 . The aim of this subsection is to show that if $\Sigma \subseteq D^4$ is a \mathbb{Z} -surface of genus $g \geq 3$ with $\Delta_{\partial\Sigma} \doteq 1$, then $\lambda_{D^4_\Sigma} \cong \mathcal{H}_2^{\oplus g}$.

Before we continue, we need to recall some general theory on ε -Hermitian forms, and on their ε -quadratic counterparts; further details can be found in [Ran81, Section 1.1] and [CS11, Section 2], where the terminology “ ε -symmetric” is used, as is customary in L -theory. Let R be a ring with involution and let $\varepsilon \in R$ be a central unit with $\varepsilon\bar{\varepsilon} = 1$. Let M be a finitely generated free left R -module. Let $\text{Sesq}(M)$ denote the abelian group of R -sesquilinear forms on M , meaning that $b(rx, sy) = rb(x, y)\bar{s}$ for $b \in \text{Sesq}(M)$, $x, y \in M$, and $r, s \in R$. Define an involution:

$$\begin{aligned} T_\varepsilon: \text{Sesq}(M) &\rightarrow \text{Sesq}(M) \\ (T_\varepsilon b)(y, x) &= \varepsilon \overline{b(x, y)}. \end{aligned}$$

A quick check shows that $(T_\varepsilon^2 b)(x, y) = \varepsilon b(x, y)\bar{\varepsilon} = \varepsilon\bar{\varepsilon}b(x, y) = b(x, y)$, so the conditions that ε is central and $\varepsilon\bar{\varepsilon} = 1$ are crucial for T_ε to be an involution. This enables us to define the *symmetric Q -group* $Q^\varepsilon(M)$ of M and the *quadratic Q -group* $Q_\varepsilon(M)$ of M via the exact sequence

$$0 \rightarrow Q^\varepsilon(M) \rightarrow \text{Sesq}(M) \xrightarrow{1-T_\varepsilon} \text{Sesq}(M) \rightarrow Q_\varepsilon(M) \rightarrow 0.$$

In other words,

$$Q^\varepsilon(M) := \ker(1 - T_\varepsilon) \text{ and } Q_\varepsilon(M) := \text{coker}(1 - T_\varepsilon).$$

A sesquilinear form $b: M \times M \rightarrow R$ is called *ε -Hermitian* if it belongs to $Q^\varepsilon(M)$, that is if $b(y, x) = \varepsilon \overline{b(x, y)}$ for every $x, y \in M$. A $(+1)$ -Hermitian form is a Hermitian form in the usual sense, while a (-1) -Hermitian form is a skew-Hermitian form in the usual sense. A ε -Hermitian form is called *hyperbolic* if it is isometric to $H^\varepsilon(R)^{\oplus g} = \left(R^{2g}, \begin{pmatrix} 0 & 1 \\ \varepsilon & 0 \end{pmatrix}^{\oplus g}\right)$ for some $g \geq 0$. The reason for introducing this terminology is that we will shortly be concerned with $(-t)$ -Hermitian forms over Λ , and their quadratic analogues.

An element $\psi \in Q_\varepsilon(M)$ is called an *ε -quadratic form*. To an ε -quadratic form in $Q_\varepsilon(M)$ is associated its *symmetrisation* $\varphi := (1 + T_\varepsilon)(\psi) \in Q^\varepsilon(M)$. Given $\varphi \in Q^\varepsilon(M)$, a quadratic form ψ with $(1 + T_\varepsilon)(\psi) = \varphi$ is called a *quadratic refinement of φ* . The symmetrisation is well-defined on equivalence classes in $Q_\varepsilon(M)$ because $(1 + T_\varepsilon)(1 - T_\varepsilon) = 1 - T_\varepsilon^2 = 0$. Note that quadratic forms are considered up to addition of forms in the image of $1 - T_\varepsilon$. A quadratic form over R is *hyperbolic* if it is isometric to $H_\varepsilon(R)^{\oplus g} = \left(R^{2g}, \begin{pmatrix} 0 & 1 \\ \varepsilon & 0 \end{pmatrix}^{\oplus g}\right)$ for some $g \geq 0$. Let us emphasise that subscripts denote quadratic forms, while superscripts denote ε -Hermitian forms.

Here is the key relationship between $(-t)$ -Hermitian and $(-t)$ -quadratic forms that we shall exploit in the proof of Theorem 7.4 below.

Lemma 7.1. *Let M be a finitely generated free Λ -module. Then the map*

$$(1 + T_{-t}): Q_{-t}(M) \rightarrow Q^{-t}(M)$$

is injective.

Proof. To prove the lemma we will show the following facts.

(1) For $\varphi \in Q^{-t}(M)$, the assignment $\varphi \mapsto (t^{-1} - 1)\varphi$ determines a map

$$\Omega: Q^{-t}(M) \rightarrow Q^{+1}(M).$$

(2) For $\psi \in \text{Sesq}(M)$, the assignment $\psi \mapsto (t^{-1} - 1)\psi$ determines an injective map

$$\Xi: Q_{-t}(M) \rightarrow Q_{+1}(M).$$

(3) The map $(1 + T_{+1}): Q_{+1}(M) \rightarrow Q^{+1}(M)$ is injective.

(4) The diagram

$$\begin{array}{ccc} Q_{-t}(M) & \xrightarrow{\Xi} & Q_{+1}(M) \\ \downarrow 1+T_{-t} & & \downarrow 1+T_{+1} \\ Q^{-t}(M) & \xrightarrow{\Omega} & Q^{+1}(M) \end{array}$$

commutes.

Then, since the right-down composition is injective, it follows that the left vertical map is injective, as desired. So it suffices to prove these four assertions.

For (1), suppose that $\varphi \in Q^{-t}(M) = \ker(1 - T_{-t})$, that is $\varphi = -t\bar{\varphi}^T$. Then

$$\overline{(t^{-1} - 1)\varphi}^T = (t - 1)\bar{\varphi}^T = (t^{-1} - 1)(-t)\bar{\varphi}^T = (t^{-1} - 1)\varphi.$$

Therefore $\Omega(\varphi) = (t^{-1} - 1)\varphi \in \ker(1 - T_{+1}) = Q^{+1}(M)$ as desired.

To prove (2), first we show that $\Xi: Q_{-t}(M) \rightarrow Q_{+1}(M)$ is well-defined. We need to see that for every sesquilinear form $\theta \in \text{Sesq}(M)$, the element $(1 - T_{-t})(\theta) = \theta - (-t)\bar{\theta}^T = \theta + t\bar{\theta}^T$ maps to the trivial element in $Q_{+1}(M)$. This is indeed the case since

$$(t^{-1} - 1)(\theta + t\bar{\theta}^T) = (t^{-1} - 1)\theta - (t - 1)\bar{\theta}^T = (1 - T_{+1})((t^{-1} - 1)\theta)$$

is trivial in $\text{coker}(1 - T_{+1}) = Q_{+1}(M)$. Therefore Ξ is well-defined.

Now we prove that $\Xi: Q_{-t}(M) \rightarrow Q_{+1}(M)$ is injective. Let B and C be two $(-t)$ -quadratic forms in $Q_{-t}(M)$, and suppose that $Z := (t^{-1} - 1)(B - C) = 0 \in Q_{+1}(M)$. That is, $Z \in \text{im}(1 - T_{+1})$, so, choosing a basis for $M \cong \Lambda^n$, there exists a matrix X over Λ such that

$$(t^{-1} - 1)(B - C) = Z = X - \bar{X}^T.$$

We assert that there exists a matrix Y over Λ such that

$$Z = (t^{-1} - 1)Y - (t - 1)\bar{Y}^T.$$

Write the entries of the matrix $Z = X - \bar{X}^T$ as z_{ij} . Since $X - \bar{X}^T$ is skew-Hermitian, we have $\bar{z}_{ij} = -z_{ji}$. Also z_{ij} is divisible by $t^{-1} - 1$ for every i, j . For $i < j$, define the entries y_{ij} of Y by the formula $y_{ij} := z_{ij}/(t^{-1} - 1)$. For $i > j$ define $y_{ij} := 0$. This way $(t^{-1} - 1)y_{ij} - (t - 1)\bar{y}_{ji} = z_{ij}$ for $i \neq j$. It remains to define the diagonal entries of Y . For each i , since the Laurent polynomial z_{ii} satisfies $z_{ii} = -\bar{z}_{ii}$, we have $z_{ii}(-1) = -z_{ii}(-1)$, and so $z_{ii}(-1) = 0$. Therefore z_{ii} is divisible by $t + 1$. Since $t^{-1} - 1$ and $(t + 1)$ are coprime, we deduce that $z_{ii} = (t + 1)(t^{-1} - 1)q$ for some polynomial q that satisfies $q = \bar{q}$. Now set $y_{ii} := q$. We compute:

$$(t^{-1} - 1)y_{ii} - (t - 1)\bar{y}_{ii} = (t^{-1} - 1 - (t - 1))q = (t^{-1} - t)q = (t + 1)(t^{-1} - 1)q = z_{ii}.$$

This concludes the proof of the assertion that $Z = (t^{-1} - 1)Y - (t - 1)\bar{Y}^T$ for some matrix Y . Using this assertion, as well as the definitions of X and Y , we obtain:

$$Z = (t^{-1} - 1)(B - C) = X - \bar{X}^T = (t^{-1} - 1)Y - (t - 1)\bar{Y}^T = (t^{-1} - 1)(Y + t\bar{Y}^T).$$

This implies that $B - C = Y + t\bar{Y}^T = Y - (-t)\bar{Y}^T$, which is zero in $Q_{-t}(M)$. Therefore the map $\Xi: Q_{-t}(M) \rightarrow Q_{+1}(M)$ is injective as desired. This completes the proof of (2).

Next we prove (3), that $1 + T_{+1}: Q_{+1}(M) \rightarrow Q^{+1}(M)$ is injective. For an arbitrary finitely generated free module M , the map $1 + T_\varepsilon$ fits into the following exact sequence:

$$0 \rightarrow \widehat{Q}^{-\varepsilon}(M) \rightarrow Q_\varepsilon(M) \xrightarrow{1+T_\varepsilon} Q^\varepsilon(M).$$

The group, $\widehat{Q}^{-\varepsilon}(M) := \ker(1 + T_\varepsilon)$ is a *hyperquadratic Q -group*; its purpose is to measure the difference between Hermitian and quadratic forms [Ran81]. We have

$$(10) \quad \widehat{Q}^{-\varepsilon}(M) = \ker(1 + T_\varepsilon: Q_\varepsilon(M) \rightarrow Q^\varepsilon(M)) = \frac{\ker(1 + T_\varepsilon)}{\text{im}(1 - T_\varepsilon)} = \frac{\ker(1 - T_{-\varepsilon})}{\text{im}(1 + T_{-\varepsilon})}.$$

We shall use the penultimate description, but include the last equation to show why the $-\varepsilon$ appears in the notation for the group $\widehat{Q}^{-\varepsilon}(M)$. Writing $M = \Lambda^n$, there is an isomorphism $\widehat{Q}^{-\varepsilon}(M) = \widehat{Q}^{-\varepsilon}(\Lambda^n) \cong \widehat{Q}^{-\varepsilon}(\Lambda)^n$ [CS11, Remark 3.4]. By (10) with $\varepsilon = +1$ and $M = \Lambda$, we have $\widehat{Q}^{-1}(\Lambda) = \ker(1 + T_{+1})/\text{im}(1 - T_{+1})$. For $p(t) \in \Lambda$, if $p(t) + p(t^{-1}) = 0$, then $p(t)$ must be of the form $p(t) = r(t) - r(t^{-1})$ for some $r(t) \in \Lambda$. It follows that $\widehat{Q}^{-1}(\Lambda) = 0$. Thus $\ker(1 + T_{+1}: Q_{+1}(M) \rightarrow Q^{+1}(M)) = 0$ and so $1 + T_{+1}$ is indeed injective. This concludes the proof of the third item of the lemma.

For (4), let $\psi \in Q_{-t}(M)$. Then

$$\begin{aligned} (1 + T_{+1}) \circ \Xi(\psi) &= (1 + T_{+1})(t^{-1} - 1)\psi = (t^{-1} - 1)\psi + (t - 1)\bar{\psi}^T \\ &= (t^{-1} - 1)(\psi - t\bar{\psi}^T) = \Omega \circ (1 + T_{-t})(\psi) \end{aligned}$$

so the diagram commutes, which proves (4) and completes the proof that $(1 + T_{-t}): Q_{-t}(M) \rightarrow Q^{-t}(M)$ is injective. \square

Next we state the algebraic cancellation result that we will use. The idea behind the proof of Theorem 7.4 below will be to engineer a situation in which we can apply cancellation.

Definition 7.2. The *Witt index* $\text{ind}(H, \theta)$ of an ε -quadratic form is the largest integer k such that a subform of (H, θ) is isometric to $H_\varepsilon(R)^{\oplus k}$.

Proposition 7.3. *Let $\varepsilon \in \Lambda$ be a central unit with $\varepsilon\bar{\varepsilon} = 1$, and let $(H, \theta), (H', \theta')$ be ε -quadratic forms over Λ . Assume that for some $n \geq 0$ there is an isometry*

$$(11) \quad (H, \theta) \oplus H_\varepsilon(\Lambda)^{\oplus n} \cong (H', \theta') \oplus H_\varepsilon(\Lambda)^{\oplus n}.$$

If $\text{ind}(H', \theta') \geq 3$, then there is an isometry

$$(12) \quad (H, \theta) \cong (H', \theta').$$

Proof. We apply a result due to Bass [Bas73, Corollary IV.3.6]. Given a ring R with involution, we write $R_0 = \{\sum_i x_i \bar{x}_i \mid x_i \in R\}$ for the norm subring of R , as well as $\text{maxspec}(R_0)$ for the set of all maximal ideals of R_0 under the Zariski topology, and $d_R := \dim \text{maxspec}(R_0)$. Detailed definitions of these notions are irrelevant: we need only know that $d_\Lambda = 2$; see e.g. [Kha17, Proposition 2.2] or [HT97, p. 439]. Let $(H, \theta), (H', \theta')$ be quadratic forms over R . If there is an isometry

$$(H, \theta) \oplus H_\varepsilon(R)^{\oplus n} \cong (H', \theta') \oplus H_\varepsilon(R)^{\oplus n}$$

and if $\text{ind}((H', \theta') \oplus H_\varepsilon(R)) \geq d_R + 2$, then there is an isometry $(H, \theta) \cong (H', \theta')$ [Bas73, Corollary IV.3.6]. In particular, this cancellation result holds if $\text{ind}(H', \theta') \geq d_R + 1$. As mentioned above, for $R = \Lambda$, we have $d_\Lambda = 2$. Since

$$\text{ind}((H', \theta') \oplus H_\varepsilon(\Lambda)) \geq 3 + 1 = d_\Lambda + 2,$$

the result of Bass says that (11) implies (12). \square

The next theorem is the main result of this subsection.

Theorem 7.4. *If $\Sigma \subseteq D^4$ is a \mathbb{Z} -surface of genus $g \geq 3$ whose boundary is an Alexander polynomial one knot $K \subseteq S^3$, then*

$$\lambda_{D_\Sigma^4} \cong \mathcal{H}_2^{\oplus g}.$$

Proof. We outline the strategy of the proof. Since the knot K has Alexander polynomial one, using Proposition 6.3, there exists an integer $n \geq 0$ and an isometry

$$(13) \quad \lambda_{D_\Sigma^4} \oplus \mathcal{H}_2^{\oplus n} \cong \lambda_{N_{\Sigma^t}} \oplus \mathcal{H}_2^{\oplus n}.$$

By Lemma 6.5, $\lambda_{N_{\Sigma^t}} \cong \mathcal{H}_2^{\oplus g}$, so in fact we have

$$(14) \quad \lambda_{D_\Sigma^4} \oplus \mathcal{H}_2^{\oplus n} \cong \mathcal{H}_2^{\oplus g+n}.$$

We cannot apply the cancellation result of Proposition 7.3 directly, since $\mathcal{H}_2^{\oplus n}$ is not hyperbolic over Λ . In order to apply cancellation, the idea is to use the relative intersection pairing

$$\lambda_{D_\Sigma^4}^\partial : H_2(D_\Sigma^4; \Lambda) \times H_2(D_\Sigma^4, \partial D_\Sigma^4; \Lambda) \rightarrow \Lambda$$

instead of the ‘‘absolute’’ intersection pairing $\lambda_{D_\Sigma^4}$. We will see, using (13) and appropriate bases, that the pairing $\lambda_{D_\Sigma^4}^\partial$ is represented by a $(-t)$ -Hermitian matrix over Λ . We will then apply the cancellation result of Proposition 7.3 with $\varepsilon = -t$ to these forms, before deducing the desired conclusion on the original absolute pairings. Note that the adjoints of the relative and absolute pairings fit into the following commutative diagram:

$$(15) \quad \begin{array}{ccc} H_2(D_\Sigma^4; \Lambda) & \xrightarrow{\hat{\lambda}} & \overline{\text{Hom}_\Lambda(H_2(D_\Sigma^4; \Lambda), \Lambda)} \\ & \searrow q & \nearrow \hat{\lambda}^\partial \\ & H_2(D_\Sigma^4, \partial D_\Sigma^4; \Lambda) & \end{array}$$

For the first step of the proof we will choose bases for $H_2(D_\Sigma^4; \Lambda)$ and $H_2(D_\Sigma^4, \partial D_\Sigma^4; \Lambda)$ and describe the map q with respect to these bases. First a short rank computation.

Claim 3. $H_2(D_\Sigma^4; \Lambda) \cong \Lambda^{2g}$.

Proof. Since $H_2(D_\Sigma^4; \Lambda)$ is free (by Lemma 3.2), we need only show that its rank is $b_1(\Sigma) = 2g$. Using $H_1(D_\Sigma^4) = \mathbb{Z}$, a Mayer-Vietoris argument shows that $b_2(D_\Sigma^4) = b_1(\Sigma)$ as well as $b_3(D_\Sigma^4) = 0$. Since we also have $b_4(D_\Sigma^4) = 0$ and $b_0(D_\Sigma^4) = 1$, we get $\chi(D_\Sigma^4) = b_1(\Sigma)$. The Euler characteristic can also be computed with $Q = \mathbb{Q}(t)$ coefficients. By Lemma 3.2, we have $b_i^Q(D_\Sigma^4) = 0$ for $i \neq 2$, and therefore $\text{rk}_\Lambda H_2(D_\Sigma^4; \Lambda) = \chi^Q(D_\Sigma^4) = b_1(\Sigma)$ as asserted. \square

Claim 4. There is a choice of bases with respect to which the homomorphism

$$q : H_2(D_\Sigma^4; \Lambda) \rightarrow H_2(D_\Sigma^4, \partial D_\Sigma^4; \Lambda)$$

is represented by the matrix $(t - 1) \text{Id}$.

Proof. We first base $H_2(D_\Sigma^4, \partial D_\Sigma^4; \Lambda)$ using the fact that the Alexander polynomial of K is one. We have an exact sequence as part of the long exact sequence of the pair $(D_\Sigma^4, \partial D_\Sigma^4)$:

$$H_2(D_\Sigma^4; \Lambda) \xrightarrow{q} H_2(D_\Sigma^4, \partial D_\Sigma^4; \Lambda) \xrightarrow{\delta} H_1(\partial D_\Sigma^4; \Lambda) \rightarrow 0.$$

Here $H_1(D_\Sigma^4; \Lambda) = 0$ because $\pi_1(D_\Sigma^4) \cong \mathbb{Z}$. In particular, the map q presents the Λ -module $H_1(\partial D_\Sigma^4; \Lambda)$. Use Lemma 3.2, Lemma 5.5, Claim 3, and the fact that $\Delta_K = 1$ to deduce that

$$H_2(D_\Sigma^4; \Lambda) \cong \Lambda^{2g} \cong H_2(D_\Sigma^4, \partial D_\Sigma^4; \Lambda) \text{ and } H_1(\partial D_\Sigma^4; \Lambda) \cong \bigoplus_{i=1}^{2g} \Lambda/(t-1).$$

Choose a set of generators $\{\gamma_k\} \subseteq H_1(\partial D_\Sigma^4; \Lambda)$, represented in the form

$$g_k \times \{\text{pt}\} \subseteq \Sigma \times \{\text{pt}\} \subseteq \Sigma \times S^1 \subseteq \partial D_\Sigma^4,$$

where $\{g_k\}$ is a symplectic basis of curves on Σ . Choose a basis $\{x_i\}$ for $H_2(D_\Sigma^4, \partial D_\Sigma^4; \Lambda)$ such that $\delta(x_i) = \gamma_i$ for $1 \leq i \leq 2g$. The homology classes x_k can be represented by embedded surfaces $S_i \subseteq D_\Sigma^4$ with $\partial S_i = \gamma_i$ for $1 \leq i \leq 2g$ [FNOP19, Section 10.3].

Next, we base $H_2(D_\Sigma^4; \Lambda)$. For each $i = 1, \dots, 2g$, define a closed surface representing a class $y_i \in H_2(D_\Sigma^4; \Lambda)$, as follows. Consider the torus $g_i \times S^1 = \partial(\bar{v}\Sigma|_{g_i})$, and surger it along γ_i using the surface S_i . That is, remove an annular neighbourhood $g_i \times (p, p') \subseteq g_i \times S^1$ of $\gamma_i = g_i \times \{\text{pt}\}$, leaving $B := g_i \times (S^1 \setminus (p, p'))$. Then glue $-S_i$ to $g_i \times \{p\}$ and a push-off S'_i of S_i to $g_i \times \{p'\}$. Call the resulting surface T_i , and set $y_i := [T_i] \in H_2(D_\Sigma^4; \Lambda)$ for $1 \leq i \leq 2g$. Then, for $i = 1, \dots, 2g$, in $H_2(D_\Sigma^4, \partial D_\Sigma^4; \Lambda)$ we have

$$(16) \quad q(y_i) = q([T_i]) = [-S_i] + [B \cup S'_i] = -x_i + tx_i = (t-1)x_i.$$

Here the t arises because the annulus B wraps around a meridian of Σ . This process is illustrated in Figure 3.

Consider the following diagram of short exact sequences

$$\begin{array}{ccccccccc} 0 & \longrightarrow & \Lambda^{\oplus 2g} & \xrightarrow{(t-1)\text{Id}} & \Lambda^{\oplus 2g} & \xrightarrow{\text{proj}} & (\Lambda/(t-1))^{\oplus 2g} & \longrightarrow & 0 \\ & & \downarrow y & & \downarrow x & & \downarrow \gamma & & \\ 0 & \longrightarrow & H_2(D_\Sigma^4; \Lambda) & \xrightarrow{q} & H_2(D_\Sigma^4, \partial D_\Sigma^4; \Lambda) & \longrightarrow & H_1(\partial D_\Sigma^4; \Lambda) & \longrightarrow & 0, \end{array}$$

where the map y (resp. x) sends the canonical basis $\{e_i\}_{i=1}^{2g}$ of Λ^{2g} to $\{y_i\}_{i=1}^{2g}$ (resp. to $\{x_i\}_{i=1}^{2g}$) and γ maps the $[e_i]$ to the γ_i . This diagram commutes thanks to (16). By construction x and γ are isomorphisms and therefore so is y by the five lemma. Thus $\{y_i\}_{i=1}^{2g}$ forms a basis of $H_2(D_\Sigma^4; \Lambda)$ and q is represented by $(t-1)\text{Id}$. This completes the proof of Claim 4. \square

Recall from (14) that there is an isometry $\alpha: \lambda_{D_\Sigma^4} \oplus \mathcal{H}_2^{\oplus n} \cong \mathcal{H}_2^{\oplus(g+n)}$. Denote the matrices of $\lambda_{D_\Sigma^4}$ and $\lambda_{D_\Sigma^4}^\partial$ with respect to the previously described bases by L and L^∂ respectively. Also represent the isometry α by a matrix A with respect to these bases. Therefore we have a congruence

$$(17) \quad A^T \left(L \oplus \left(\begin{array}{cc} 0 & t-1 \\ t^{-1}-1 & 0 \end{array} \right)^{\oplus n} \right) \bar{A} = \left(\begin{array}{cc} 0 & t-1 \\ t^{-1}-1 & 0 \end{array} \right)^{\oplus(g+n)}.$$

Our aim is now to factor out a $(t^{-1}-1)$ from both sides of this equation. Recall from (15) that the adjoints of the pairings $\lambda_{D_\Sigma^4}$ and $\lambda_{D_\Sigma^4}^\partial$ are related by $\widehat{\lambda}_{D_\Sigma^4}^\partial \circ q = \widehat{\lambda}_{D_\Sigma^4}$. With respect to




FIGURE 3. Representing $y_i \in H_2(D_\Sigma^4; \Lambda)$ by an immersed sphere T_i in D_Σ^4 obtained by surgering the torus $g_i \times S^1$, using a surface S_i representing the relative homology class x_i . The top left picture shows a neighbourhood of a portion of g_i , intersected with a carefully chosen 3-dimensional subspace that contains the intersections of Σ and S_i with this neighbourhood. Bottom, a 4-dimensional picture of T_i , consisting of three 3-dimensional slices at certain special values of the fourth coordinate (time), in which Σ , $-S_i$, S'_i , and most of B appear. The annulus B joins $-S_i$ and S'_i together; in intermediate time values only a line of B appears, including in the middle slice. The top right picture shows a cross section of T_i , looking along g_i .

our bases (and recalling (16)), for every $x, y \in H_2(D_\Sigma^4; \Lambda)$, we therefore obtain

$$\begin{aligned} x^T L^\partial (t^{-1} - 1) \bar{y} &= x^T L^\partial \overline{(t-1)y} = \lambda_{D_\Sigma^4}^\partial(x, q(y)) = \widehat{\lambda}^\partial(q(y))(x) \\ &= \widehat{\lambda}(y)(x) = \lambda(x, y) = x^T L \bar{y}. \end{aligned}$$

It follows that $L^\partial(t^{-1} - 1) = L$. Combine this with $\mathcal{H}_2 = \begin{pmatrix} 0 & t^{-1} \\ t^{-1} & 0 \end{pmatrix} = \begin{pmatrix} 0 & -t \\ 1 & 0 \end{pmatrix} (t^{-1} - 1)$ to rewrite (17) as:

$$A^T \left(L^\partial (t^{-1} - 1) \oplus \begin{pmatrix} 0 & -t \\ 1 & 0 \end{pmatrix} (t^{-1} - 1) \right) \bar{A} = \begin{pmatrix} 0 & -t \\ 1 & 0 \end{pmatrix}^{\oplus g+n} (t^{-1} - 1).$$

As $(t^{-1} - 1) \cdot \text{Id}$ is diagonal with constant diagonal coefficients, it is central and thus we obtain

$$(t^{-1} - 1) \cdot A^T \left(L^\partial \oplus \begin{pmatrix} 0 & -t \\ 1 & 0 \end{pmatrix}^{\oplus n} \right) \bar{A} = (t^{-1} - 1) \begin{pmatrix} 0 & -t \\ 1 & 0 \end{pmatrix}^{\oplus (g+n)}.$$

As $(t^{-1} - 1) \cdot \text{Id}$ is nondegenerate over Λ , it follows that

$$(18) \quad A^T \left(L^\partial \oplus \begin{pmatrix} 0 & -t \\ 1 & 0 \end{pmatrix}^{\oplus n} \right) \bar{A} = \begin{pmatrix} 0 & -t \\ 1 & 0 \end{pmatrix}^{\oplus (g+n)}.$$

In particular L^∂ stabilises, via $(-t)$ -Hermitian matrices, to a $(-t)$ -Hermitian form, so it follows that L^∂ is itself $(-t)$ -Hermitian.

In order to apply cancellation, we refine (18) to a statement about quadratic forms. Set

$$\Psi := \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}.$$

Then $(1 + T_{-t})\Psi = \begin{pmatrix} 0 & -t \\ 1 & 0 \end{pmatrix}$, so Ψ is a quadratic refinement for $\begin{pmatrix} 0 & -t \\ 1 & 0 \end{pmatrix}$. Therefore $\Psi^{\oplus (g+n)}$ is a quadratic refinement for $\begin{pmatrix} 0 & -t \\ 1 & 0 \end{pmatrix}^{\oplus (g+n)}$, which is the right hand side of (18). Thus $\Psi^{\oplus (g+n)}$ is a quadratic refinement for

$$A^T \left(L^\partial \oplus \begin{pmatrix} 0 & -t \\ 1 & 0 \end{pmatrix}^{\oplus n} \right) \bar{A}.$$

Rearranging, we deduce that $A^{-T}\Psi^{\oplus (g+n)}\bar{A}^{-1}$ is a quadratic refinement for the $(-t)$ -Hermitian matrix $L^\partial \oplus \begin{pmatrix} 0 & -t \\ 1 & 0 \end{pmatrix}^{\oplus n}$. Now, corresponding to this decomposition, write:

$$A^{-T}\Psi^{\oplus (g+n)}\bar{A}^{-1} = \begin{pmatrix} \Psi^\partial & U \\ V & \Psi_{\text{ind}} \end{pmatrix}$$

where this defined the block entries on the right hand side. Since this is a quadratic refinement of the block diagonal $(-t)$ -Hermitian matrix $L^\partial \oplus \begin{pmatrix} 0 & -t \\ 1 & 0 \end{pmatrix}^{\oplus n}$, we deduce that $V - t\bar{U}^T = 0$ and therefore

$$(19) \quad A^{-T}\Psi^{\oplus (g+n)}\bar{A}^{-1} = \begin{pmatrix} \Psi^\partial & 0 \\ 0 & \Psi_{\text{ind}} \end{pmatrix} + \begin{pmatrix} 0 & U \\ t\bar{U}^T & 0 \end{pmatrix} = \Psi^\partial \oplus \Psi_{\text{ind}} \in Q_{-t}(\Lambda^{\oplus 2(g+n)}).$$

This procedure therefore endows both the $(-t)$ -Hermitian matrices L^∂ and $\begin{pmatrix} 0 & -t \\ 1 & 0 \end{pmatrix}^{\oplus n}$ with quadratic refinements, which we denoted by Ψ^∂ and Ψ_{ind} respectively. Rearranging (19) we obtain the following isometry of quadratic forms:

$$(20) \quad A^T(\Psi^\partial \oplus \Psi_{\text{ind}})\bar{A} = \Psi^{\oplus (g+n)} \in Q_{-t}(\Lambda^{\oplus 2(g+n)}).$$

In order to apply cancellation, we need that Ψ_{ind} agrees with the quadratic refinement $\Psi_{\text{triv}} = \Psi^{\oplus n}$ of $\begin{pmatrix} 0 & -t \\ 1 & 0 \end{pmatrix}^{\oplus n}$ in the Q -group $Q_{-t}(\Lambda^{2n})$. That both are quadratic refinements of the form $\begin{pmatrix} 0 & -t \\ 1 & 0 \end{pmatrix}^{\oplus n}$ in $Q^{-t}(\Lambda^{2n})$ can be restated as

$$(1 + T_{-t})(\Psi_{\text{ind}}) = (1 + T_{-t})(\Psi_{\text{triv}}) \in Q^{-t}(\Lambda^{2n}).$$

By Lemma 7.1, $(1 + T_{-t})$ is injective, and so $\Psi_{\text{ind}} = \Psi_{\text{triv}}$ in $Q_{-t}(\Lambda^{2n})$, as desired.

Now recall that Ψ^∂ denotes the quadratic refinement of L^∂ (the matrix for the relative intersection form), and that we set $\Psi := \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}$, as well as $\Psi_{\text{triv}} = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}^{\oplus n}$. Since Ψ_{triv} and Ψ_{ind} agree in $Q_{-t}(\Lambda^{2n})$, we deduce from (20) that

$$(21) \quad A^T \left(\Psi^\partial \oplus \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}^{\oplus n} \right) \bar{A} = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}^{\oplus (g+n)} \in Q_{-t}(\Lambda^{2(g+n)}).$$

Since $\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$, we see that $\begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}$ is congruent to the hyperbolic $(-t)$ -quadratic form $H_{-t}(\Lambda)$. We have established the following isometry of $(-t)$ -quadratic forms:

$$(22) \quad \Psi^\partial \oplus H_{-t}(\Lambda)^{\oplus n} \cong H_{-t}(\Lambda)^{\oplus g} \oplus H_{-t}(\Lambda)^{\oplus n} \in Q_{-t}(\Lambda^{2(g+n)}).$$

Since $g \geq 3$, we now apply the cancellation result of Proposition 7.3 to (22) and symmetrise the result to deduce that L^∂ is congruent to $H^{-t}(\Lambda)^{\oplus g}$, and therefore to $\begin{pmatrix} 0 & -t \\ 1 & 0 \end{pmatrix}^{\oplus g}$. Multiplying both sides of this congruence by $t^{-1} - 1$ shows that L is congruent to $\mathcal{H}_2^{\oplus g}$. This concludes the proof of the theorem. \square

7.2. Surfaces in S^4 and D^4 . We collect and prove our results concerning (rel. boundary) ambient isotopy of surfaces in D^4 and S^4 . Since the proofs of these results were already outlined in the introduction or elsewhere, we proceed swiftly.

Theorem 7.5. *Let $\Sigma_0, \Sigma_1 \subseteq D^4$ be \mathbb{Z} -surfaces of genus g with boundary K .*

- (1) *Suppose there is an isometry $F \in \text{Iso}(\lambda_{D_{\Sigma_0}^4}, \lambda_{D_{\Sigma_1}^4})$ and write $\partial F = h_K \oplus h_\Sigma$. If h_K is represented by an orientation-preserving homeomorphism that is the identity on ∂E_K , then Σ_0 and Σ_1 are topologically ambiently isotopic rel. boundary.*
- (2) *If $\Delta_K = 1, g = 1, 2$ and $\lambda_{D_{\Sigma_0}^4} \cong \lambda_{D_{\Sigma_1}^4}$, then Σ_0 and Σ_1 are topologically ambiently isotopic rel. boundary.*
- (3) *If $\Delta_K = 1$ and $g \neq 1, 2$, then Σ_0 and Σ_1 are topologically ambiently isotopic rel. boundary.*

Proof. The first assertion is the second item of Theorem 5.8. When $\Delta_K = 1$, we have $H_1(E_K; \Lambda) = 0$, so $h_K = \text{Id}$ is automatic, and the second assertion follows from the first. For the third assertion, additionally use that $\lambda_{D_{\Sigma_0}^4} \cong \lambda_{D_{\Sigma_1}^4}$ when $g \neq 1, 2$ by Theorem 7.4. \square

Theorem 7.6. *Let $\Sigma_0, \Sigma_1 \subseteq S^4$ be closed \mathbb{Z} -surfaces of genus g .*

- (1) *For $g = 1, 2$, if $\lambda_{\Sigma_0} \cong \lambda_{\Sigma_1}$, then Σ_0 and Σ_1 are topologically ambiently isotopic.*
- (2) *For $g \neq 1, 2$, the surfaces Σ_0 and Σ_1 are topologically ambiently isotopic.*

Proof. Both assertions follow from Theorem 7.5 by removing a (D^4, D^2) -pair from (S^4, Σ_i) . \square

7.3. Pushed-in Seifert surfaces for Alexander polynomial one knots. We prove Theorem 1.9 from the introduction, which states that any two pushed-in Seifert surfaces of the same genus for an Alexander polynomial one knot are topologically ambiently isotopic in D^4 .

Recall that a *sublagrangian* for a nonsingular ε -Hermitian form (H, λ) is a direct summand $M \subseteq H$ such that $\lambda(M \times M) = 0$ (i.e. $M \subseteq M^\perp$). A sublagrangian is a *Lagrangian* if $M = M^\perp$. For a nonsingular ε -quadratic form (H, λ, μ) , the definitions are identical, with the additional requirement that $\mu(M) = 0$.

Lemma 7.7. *Let (H, λ) be a nonsingular ε -Hermitian form on a free Λ -module H . If $M \subseteq H$ is a half-rank sublagrangian, then M is a Lagrangian.*

Proof. We know that $M \subseteq M^\perp$ and must show that $M = M^\perp$. Equivalently, we must show that $M/M^\perp = 0$. We will show that M/M^\perp is both torsion and torsion-free.

Claim 5. The inclusion $M^\perp \hookrightarrow H$ is split and $M^\perp \subseteq H$ is a half-rank summand.

Proof. Consider the exact sequence $0 \rightarrow M^\perp \rightarrow H \xrightarrow{\iota^* \widehat{\lambda}} M^*$. It now suffices to show that $\iota^* \widehat{\lambda}$ is surjective: since M^* is free and $\text{rk}_\Lambda(M^*) = \text{rk}_\Lambda(M) = \frac{1}{2} \text{rk}_\Lambda(H)$, the result would then follow. Since λ is nonsingular, $\widehat{\lambda}$ is surjective and so it suffices to prove that ι^* is injective. This occurs if and only if $\text{Ext}_\Lambda^1(H/M, \Lambda) = 0$. But since M is a summand, it follows that $H \cong M \oplus H/M$. Since H is free, H/M is projective over Λ and therefore free over Λ [Lam06, Chapter V, Corollary 4.12]. Thus $\text{Ext}_\Lambda^1(H/M, \Lambda) = 0$, so $\iota^* \widehat{\lambda}$ is surjective and thus $M^\perp \hookrightarrow H$ is a split injection, as claimed. \square

The fact that M/M^\perp is torsion follows from the claim because $M^\perp \subseteq M$ and $\text{rk}_\Lambda(M^\perp) = \text{rk}_\Lambda(M)$. We now show that M/M^\perp is torsion-free. Since submodules of torsion-free modules are torsion-free, it suffices to show that $H/M^\perp \supseteq M/M^\perp$ is torsion-free. But now by the claim, we know that H/M^\perp is (isomorphic to) a submodule of the free module $H \cong M^\perp \oplus H/M^\perp$ and so it is indeed torsion-free. This concludes the proof of the lemma. \square

Given a Seifert matrix A for a genus g Seifert surface S , we write $(H, A) := (\mathbb{Z}^{2g}, A)$. Use $F \subseteq D^4$ to denote the result of pushing S into D^4 . After some choice of bases, the equivariant intersection form $(H_2(D_F^4; \Lambda), \lambda_{D_F^4})$ is isometric to $(H \otimes_{\mathbb{Z}} \Lambda, (1-t)A^T + (1-t^{-1})A)$ [COT04, proof of Lemma 5.4] (see also [CFT18, Theorem 1.3]).

Remark 7.8. Observe that $(H \otimes_{\mathbb{Z}} \Lambda, A - tA^T, A)$ defines a $(-t)$ -quadratic form. Furthermore, if A is a Seifert matrix for an Alexander polynomial one knot K , then this quadratic form is nonsingular since $\det(A - tA^T) \doteq \Delta_K(t) \doteq 1$.

Recall that a *metaboliser* for a Seifert matrix (\mathbb{Z}^{2g}, A) is a half-rank direct summand $L \subseteq \mathbb{Z}^{2g}$ such that $x^T A y = 0$ for all $x, y \in L$. Since Alexander polynomial one knots are (in particular) algebraically slice, their Seifert matrices admit metabolisers.

Proposition 7.9. *Let $(H, A) := (\mathbb{Z}^{2g}, A)$ be a Seifert matrix for an Alexander polynomial one knot K . If $L \subseteq H$ is a metaboliser for (H, A) , then $L \otimes_{\mathbb{Z}} \Lambda$ is a Lagrangian for $(H \otimes_{\mathbb{Z}} \Lambda, A - tA^T, A)$ and so this latter $(-t)$ -quadratic form is hyperbolic.*

Proof. We verify the three points in the definition of a Lagrangian. Since A vanishes on L , the extended form (also denoted A) vanishes on $M := L \otimes_{\mathbb{Z}} \Lambda$. It follows that the $(-t)$ -Hermitian form $A - tA^T$ vanishes on $M \times M$. Since $L \subseteq H$ is a summand, so is $M \subseteq H \otimes_{\mathbb{Z}} \Lambda$. Thus $M \subseteq (H \otimes_{\mathbb{Z}} \Lambda, A - tA^T)$ is a sublagrangian of a $(-t)$ -Hermitian form, with $H \otimes_{\mathbb{Z}} \Lambda$ a free module. We showed in Lemma 7.7 that since M is half-rank, this forces $M^\perp = M$.

The last sentence of the proposition follows because it is known that if a nonsingular ε -quadratic form (H, λ, μ) admits a Lagrangian, then it is isometric to the standard hyperbolic form [CLM, Lemma 7.92]. \square

Proposition 7.10. *If A is a Seifert matrix for an Alexander polynomial one knot K , then*

$$(H \otimes_{\mathbb{Z}} \Lambda, (1-t)A^T + (1-t^{-1})A) \cong \mathcal{H}_2^{\oplus g}.$$

Proof. Using Proposition 7.9, we deduce the following sequence of isometries:

$$\begin{aligned} (H \otimes_{\mathbb{Z}} \Lambda, A - tA^T, A) &\cong H_{-t}(\Lambda)^{\oplus g} = \left(\Lambda^{2g}, \begin{pmatrix} 0 & 1 \\ -t & 0 \end{pmatrix}^{\oplus g}, \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}^{\oplus g} \right) \\ &\cong \left(\Lambda^{2g}, \begin{pmatrix} 0 & -t \\ 1 & 0 \end{pmatrix}^{\oplus g}, \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}^{\oplus g} \right). \end{aligned}$$

Multiplying both sides by $(1 - t^{-1})$ then gives the assertion. \square

We are now ready to prove Theorem 1.9 from the introduction.

Theorem 7.11. *If $F_0, F_1 \subseteq D^4$ are genus g pushed-in Seifert surfaces for an Alexander polynomial one knot K , then they are topologically ambiently isotopic rel. boundary.*

Proof. Let A_0 and A_1 be Seifert matrices for F_0 and F_1 . As mentioned above, the equivariant intersection form $(H_2(D_{F_i}^4; \Lambda), \lambda_{D_{F_i}^4})$ is isometric to $(H \otimes_{\mathbb{Z}} \Lambda, (1-t)A_i^T + (1-t^{-1})A_i)$. By Proposition 7.10, both forms are isometric to $\mathcal{H}_2^{\oplus g}$. Since K is an Alexander polynomial one knot, the result now follows from Theorem 7.5. \square

8. RIM SURGERY ON SURFACES WITH KNOT GROUP \mathbb{Z} .

We prove Theorems 1.7 and 1.8 which concern rim surgery on surfaces in 4-manifolds whose knot group is \mathbb{Z} . In Subsection 8.1, we review the definition of rim surgery, in Subsection 8.2, and in Subsection 8.3, we prove the main results.

8.1. Knot surgery and twist rim surgery. We review some facts about knot surgery and rim surgery. References include [FS97, Kim06, KR08b, BS16].

Let Z be a compact, oriented 4-manifold containing a locally flat embedded torus T with trivial normal bundle, and let $J \subseteq S^3$ be a knot. Use μ_T to denote the meridian of $T \subseteq Z$ and μ_J, λ_J for the meridian and 0-framed longitude of J . Let $\varphi: \partial\nu(T) \rightarrow S^1 \times \partial E_J$ be any diffeomorphism such that $\varphi_*(\mu_T) = \lambda_J$. The 4-manifold obtained by *knot surgery* along J and φ is defined as

$$Z_J(\varphi) = (Z \setminus \nu(T)) \cup_{\varphi} (S^1 \times E_J).$$

Given a compact 4-manifold W and a locally flat embedded orientable surface $\Sigma \subseteq W$, we assume either that W and Σ are closed or that $\Sigma \subseteq W$ is a properly embedded. As we now describe, rim surgery arises from a particular type of knot surgery on the surface exterior $W_{\Sigma} := W \setminus \nu(\Sigma)$. Choose a simple closed curve $\alpha \subseteq \Sigma$ and a trivialisation of the normal bundles over α and Σ so that $(\nu(\Sigma), \nu(\alpha)) = (\Sigma \times D^2, \alpha \times I \times D^2)$; in other words the normal bundle of α inside Σ is $\alpha \times I$. This way, it is understood that $\mu_{\Sigma} = \{\text{pt}\} \times \partial D^2$, and the *rim torus* T is $\partial(\nu(\Sigma)|_{\alpha}) = \alpha \times \mu_{\Sigma}$. The rim torus is framed: if we write $\nu(\Sigma)|_{\alpha \times I} = \alpha \times I \times D^2$, then a framing of νT is given by the I -direction and the radial direction in the polar coordinates of D^2 .

In order to perform a knot surgery on W_{Σ} along J using the rim torus T , for integers m and n we consider the homeomorphism $\varphi_n^m: \partial\nu(T) \rightarrow S^1 \times \partial E_J$ determined by

$$\begin{aligned} \varphi_n^m(\alpha) &= m\mu_J + n\lambda_J + S^1, \\ \varphi_n^m(\mu_{\Sigma}) &= \mu_J, \\ \varphi_n^m(\mu_T) &= \lambda_J. \end{aligned}$$

Gluing $\Sigma \times D^2$ back into the result of this knot surgery produces a manifold $W_J(\varphi_n^m)$ that is homeomorphic to W , and in fact if W is smooth then $W_J(\varphi_n^m)$ is diffeomorphic to W ; see [Kim06, Lemma 2.4], in which a specific diffeomorphism is constructed. Thus, we obtain a new embedding $\Sigma_n^m(\alpha, J) \subseteq W$.

Definition 8.1. Let $\alpha \subseteq \Sigma$ be a simple closed curve, let J be a knot, and let $m, n \in \mathbb{Z}$ be integers. The n -roll m -twist rim surgery of a locally flat, properly embedded, orientable surface $\Sigma \subseteq W$ is the image $\Sigma_n^m(\alpha, J)$ of Σ under the homeomorphism $W \cong W_J(\varphi_n^m)$ mentioned above. If W is smooth and Σ is smoothly embedded, then since $W \cong W_J(\varphi_n^m)$ is a diffeomorphism, $\Sigma_n^m(\alpha, J)$ is smoothly embedded.

The exterior $W_{\Sigma_n^m(\alpha, J)}$ of the rim surgery $\Sigma_n^m(\alpha, J)$ is the knot surgery on W_Σ along the homeomorphism φ_n^m . The proof of the following lemma can be found in [Kim06, Proposition 3.3].

Lemma 8.2. *For a compact 4-manifold W and a locally flat, embedded, orientable surface $\Sigma \subseteq W$, we assume either that W and Σ are closed or that $\Sigma \subseteq W$ is properly embedded. Let $\alpha \subseteq \Sigma$ be a simple closed curve, let J be a knot, and let $m, n \in \mathbb{Z}$. If $\pi_1(W_\Sigma) = \mathbb{Z}_d$ with $d = \pm 1 \pmod{m}$, then*

$$\pi_1(W_\Sigma) = \pi_1(W_{\Sigma_n^m(\alpha, J)}) = \mathbb{Z}_d.$$

In our case, the knot group is infinite cyclic, so we take $d = 0$ and restrict to n -roll 1-twist surgeries. We write $\Sigma_n(\alpha, J) := \Sigma_n^1(\alpha, J)$ and $\varphi_n := \varphi_n^1$.

Remark 8.3. We note that if $J = U$ is the unknot then $\Sigma_n(\alpha, U) = \Sigma$, by [KR08b, Lemma 2.2].

8.2. The Λ -intersection form. Now we focus on the case where the ambient manifold N has a boundary and show that the Λ -intersection form of $N_{\Sigma_n(\alpha, J)}$ agrees with that of N_Σ .

We start with an observation concerning the rim torus.

Lemma 8.4. *Let $\Sigma \subseteq N = X \setminus \mathring{D}^4$ be a \mathbb{Z} -surface. The exterior of a rim torus T satisfies*

$$H_1(N_\Sigma \setminus \nu(T)) = \mathbb{Z}\langle \mu_\Sigma \rangle \oplus \langle \mathbb{Z}\mu_T \rangle.$$

Proof. Consider the Mayer-Vietoris sequence of $N_\Sigma = (N_\Sigma \setminus \nu(T)) \cup \bar{\nu}(T)$:

$$(23) \quad \cdots \rightarrow H_2(N_\Sigma) \xrightarrow{\partial} H_1(T \times S^1) \xrightarrow{\theta} H_1(\bar{\nu}(T)) \oplus H_1(N_\Sigma \setminus \nu(T)) \rightarrow H_1(N_\Sigma) \rightarrow 0.$$

We show that $\partial = 0$. Note that

$$H_1(T \times S^1) \cong H_1(T) \oplus H_1(S^1) \xrightarrow{\theta} H_1(\bar{\nu}(T)) \cong H_1(T)$$

is projection onto the first factor, so for $x \neq 0$ in $H_1(T)$, $\theta(x, n) \neq 0$. Therefore in the image of ∂ , the first coordinate vanishes. Now for a closed surface $S \subseteq N$, the definition of the connecting homomorphism implies that

$$\partial([S]) = (0, Q_N([T], [S])) \in H_1(T \times S^1) \cong H_1(T) \oplus H_1(S^1).$$

Since T is isotopic to a torus in $\Sigma \times S^1 \subseteq \partial N_\Sigma$, we deduce that $Q_N([T], [S]) = 0$. It follows that $\partial = 0$ as asserted. The Mayer-Vietoris sequence in (23) then reduces to

$$0 \rightarrow \mathbb{Z}\langle \mu_T \rangle \rightarrow H_1(N_\Sigma \setminus \nu(T)) \rightarrow H_1(N_\Sigma) \rightarrow 0.$$

Since $H_1(N_\Sigma) = \mathbb{Z}\langle \mu_\Sigma \rangle$ is free, this sequence splits, establishing the lemma. \square

Next, we extend the coefficient system on $\pi_1(N_\Sigma)$ over $\pi_1(N_{\Sigma_n(\alpha, J)})$.

Remark 8.5. The map $\pi_1(N_\Sigma) \rightarrow \mathbb{Z}$ restricts to a map $\pi_1(N_\Sigma \setminus \nu(T)) \rightarrow \mathbb{Z}$ that sends μ_Σ to 1 and μ_T to 0 (recall from Lemma 8.4 that $H_1(N_\Sigma \setminus \nu(T)) \cong \mathbb{Z}\langle\mu_\Sigma\rangle \oplus \mathbb{Z}\langle\mu_T\rangle$). Consider the map $\pi_1(S^1 \times E_J) \rightarrow \mathbb{Z}$ that sends the class of the S^1 -factor to -1 and that is the abelianisation on $\pi_1(E_J)$. We verify that these maps extend to a map $\pi_1(N_{\Sigma_n(\alpha, J)}) \rightarrow \mathbb{Z}$. We are identifying the 3-torus $\partial\bar{\nu}(T) \subseteq \partial N_\Sigma \setminus \nu(T)$ with $S^1 \times \partial E_J$ via the homeomorphism φ_n . We have the identifications $[\alpha] \sim [S^1][\mu_J][\lambda_J]^n$, $[\mu_\Sigma] \sim [\mu_J]$ and $[\mu_T] \sim [\lambda_J]$. The loop $\alpha \subseteq \Sigma$ is mapped to zero under $\pi_1(N_\Sigma) \rightarrow \mathbb{Z}$; the same is true for $[S^1][\mu_J][\lambda_J]^n$ under the map $\pi_1(S^1 \times E_J) \rightarrow \mathbb{Z}$. Similarly, both μ_Σ and μ_J (resp μ_T and λ_J) are sent to 1 (resp. 0) under the respective maps to \mathbb{Z} .

Arguing as in [KR08b, Proof of Lemma 2.3], we construct a degree one map

$$\Psi: N_{\Sigma_n(\alpha, J)} \rightarrow N_\Sigma.$$

Using obstruction theory, one can construct a degree one map $E_J \rightarrow E_U$, where $U \subseteq S^3$ is the unknot, that takes meridian to meridian and longitude to longitude. Cross it with the identity to obtain a degree one map $S^1 \times E_J \rightarrow S^1 \times E_U$. This can be glued to the identity map $\text{Id}_{N_\Sigma \setminus \nu(T)}$ to obtain a degree one map $N_{\Sigma_n(\alpha, J)} \rightarrow N_{\Sigma_n(\alpha, U)}$. By [KR08b, Lemma 2.2], there is a homeomorphism $N_{\Sigma_n(\alpha, U)} \cong N_\Sigma$ and we have therefore obtained a degree one map

$$\Psi: N_{\Sigma_n(\alpha, J)} \rightarrow N_{\Sigma_n(\alpha, U)} \cong N_\Sigma.$$

By construction, Ψ is the identity on ∂N_Σ . Furthermore, since the fundamental groups of both $N_{\Sigma_n(\alpha, J)}$ and N_Σ are generated by the meridians of the surface, Ψ induces an isomorphism on fundamental groups: indeed by construction Ψ takes meridian to meridian. In particular Ψ induces homomorphisms on the Λ -homology groups.

In order to prove that Ψ in fact induces an isometry of the equivariant intersection forms, we compute the Λ -homology of $S^1 \times E_J$.

Lemma 8.6. *Let $J \subseteq S^3$ be a knot. With respect to the coefficient system on $S^1 \times E_J$ introduced in Remark 8.5, we have $H_2(S^1 \times E_J; \Lambda) = 0$ and $H_1(S^1 \times E_J; \Lambda) = \mathbb{Z}$. Additionally, the degree one map Ψ induces an isometry*

$$\Psi_*: (H_2(N_{\Sigma_n(\alpha, J)}; \Lambda), \lambda_{N_{\Sigma_n(\alpha, J)}}) \xrightarrow{\cong} (H_2(N_\Sigma; \Lambda), \lambda_{N_\Sigma}).$$

Proof. We use the Künneth spectral sequence with

$$E_{p,q}^2 = \bigoplus_{q_1+q_2=q} \text{Tor}_p(H_{q_1}(S^1; \Lambda), H_{q_2}(E_J; \Lambda))$$

and which converges to $H_*(S^1 \times E_J; \Lambda \otimes_\Lambda \Lambda) \cong H_*(S^1 \times E_J; \Lambda)$, where $\Lambda \otimes_\Lambda \Lambda$ is a right $\mathbb{Z}[\pi_1(E_K) \times \pi_1(S^1)]$ -module via the diagonal action induced by $p \otimes q \cdot (e, s) = pe \otimes qs$ for $p, q \in \Lambda$ and $(e, s) \in \pi_1(E_K) \times \pi_1(S^1)$. Write $\varepsilon: \Lambda \rightarrow \mathbb{Z}, p(t) \mapsto p(1)$ for the augmentation map and \mathbb{Z}_ε for the resulting Λ -module structure on \mathbb{Z} . By definition of the coefficient system in Remark 8.5, we have $H_1(S^1; \Lambda) = 0$ and $H_0(S^1; \Lambda) = \Lambda/(t-1) = \mathbb{Z}_\varepsilon$. We also have $H_0(E_J; \Lambda) = \mathbb{Z}$ and $H_i(E_J; \Lambda) = 0$ for $i \geq 2$. This implies that $E_{0,2}^2 = 0$. In fact, we also deduce that $E_{1,0}^2 = \mathbb{Z}$ and $E_{2,0}^2 = 0$ because

$$E_{i,0}^2 = \text{Tor}_i^{\mathbb{Z}[\mathbb{Z}]}(\mathbb{Z}, \mathbb{Z}) = H_i(\mathbb{Z}; \mathbb{Z}) = H_i(S^1).$$

We claim that $\text{Tor}_i^\Lambda(\mathbb{Z}_\varepsilon, H_1(E_J; \Lambda)) = 0$ for $i = 0, 1$. These Tor groups are computed as the homology of the complex obtained tensoring the resolution $0 \rightarrow \Lambda \xrightarrow{t-1} \Lambda \rightarrow \mathbb{Z}_\varepsilon \rightarrow 0$ with the

Alexander module $H_1(E_J; \Lambda)$. Since multiplication by $t - 1$ induces an isomorphism on the Alexander module [Lev77, Proposition 1.2], these Tor groups vanish and the claim is proved.

Using the claim, we deduce that $E_{0,1}^2 = 0$ and $E_{1,1}^2 = 0$. It follows that $d_{2,0} = 0$ and now the first assertion is a consequence of a standard spectral sequence computation.

We now assert that Ψ induces an isomorphism on $H_2(-; \Lambda)$. Recall the decompositions $N_{\Sigma_n(\alpha, J)} = (N_\Sigma \setminus \nu(T)) \cup_{\varphi_n} (S^1 \times E_J)$ and $N_\Sigma \cong N_{\Sigma_n(\alpha, U)} = (N_\Sigma \setminus \nu(T)) \cup_{\varphi_n} (S^1 \times E_U)$. By construction, Ψ restricts to the identity on $N_\Sigma \setminus \nu(T)$ and $\partial\bar{\nu}(T)$ and therefore induces the identity on the Λ -homology of these spaces. Next, $\Psi_*: H_1(S^1 \times E_J; \Lambda) \rightarrow H_1(S^1 \times E_U; \Lambda)$ is also an isomorphism: indeed, the first assertion shows that both modules are isomorphic to $\mathbb{Z} = \Lambda/(t - 1)$, and this term comes from $H_0(E_J; \Lambda) = \mathbb{Z}_\varepsilon$, on which Ψ does indeed induce an isomorphism. Since we showed that $H_2(S^1 \times E_J; \Lambda) = 0$, the assertion now follows from the five lemma applied to the following commutative diagram, where Λ -coefficients are understood:

$$\begin{array}{ccccccccc} H_2(\partial\bar{\nu}(T)) & \longrightarrow & H_2(N_\Sigma \setminus \nu(T)) & \longrightarrow & H_2(N_{\Sigma_n(\alpha, J)}) & \longrightarrow & H_1(\partial\bar{\nu}(T)) & \longrightarrow & H_1(N_\Sigma \setminus \nu(T)) \oplus \mathbb{Z} & \longrightarrow & 0 \\ \Psi_* \downarrow = & & \Psi_* \downarrow = & & \Psi_* \downarrow & & \Psi_* \downarrow = & & \Psi_* \downarrow \cong & & \\ H_2(\partial\bar{\nu}(T)) & \longrightarrow & H_2(N_\Sigma \setminus \nu(T)) & \longrightarrow & H_2(N_\Sigma) & \longrightarrow & H_1(\partial\bar{\nu}(T)) & \longrightarrow & H_1(N_\Sigma \setminus \nu(T)) \oplus \mathbb{Z} & \longrightarrow & 0. \end{array}$$

Since Ψ induces a Λ -isomorphism on $H_2(-; \Lambda)$, it induces one on $H_2(-; \Lambda)^*$, and therefore on $H^2(-; \Lambda)$ (recall the UCSS argument from Lemma 3.2) and thus on the second relative Λ -homology groups (because $\deg(\Psi) = 1$). We conclude that Ψ_* in fact induces an isometry of the Λ -intersection forms. \square

8.3. Topological triviality of rim surgery on surfaces with knot group \mathbb{Z} . Next we prove Theorem 1.8 from the introduction.

Theorem 8.7. *Let $\Sigma \subseteq N = X \setminus \mathring{D}^4$ be a \mathbb{Z} -surface, let $\alpha \subseteq \Sigma$ be a simple closed curve, let J be a knot, and let $n \in \mathbb{Z}$ be an integer. There is a rel. boundary orientation-preserving homeomorphism of pairs*

$$\Phi: (N, \Sigma_n(\alpha, J)) \xrightarrow{\cong} (N, \Sigma).$$

that induces the same isometry as the degree one map Ψ on the equivariant intersection forms of the surface exteriors. If $N = D^4$, then the surfaces are topologically isotopic rel. boundary.

Proof. By Lemma 8.2 both surfaces have knot group \mathbb{Z} . Lemma 8.6 states that the degree one map $\Psi: N_{\Sigma_n(\alpha, J)} \rightarrow N_\Sigma$ induces an isometry $\Psi_*: \lambda_{N_{\Sigma_n(\alpha, J)}} \cong \lambda_{N_\Sigma}$. As Ψ is the identity on ∂N_Σ , we see that $\partial\Psi$ induces the identity isometry between the boundary Blanchfield forms. The result now follows from Theorem 5.8. \square

The remaining result we owe a proof of is Theorem 1.7 from the introduction. To prove this result, which involves obtaining an isotopy between closed surfaces, we need to control the map $F_{\mathbb{Z}}$ discussed in Lemma 5.10. This will rely on the following result.

Lemma 8.8. *Let $\Sigma \subseteq N$ be a \mathbb{Z} -surface, let $\alpha \subseteq \Sigma$ be a simple closed curve with associated rim torus T , and let J be a knot. The following sequences of inclusion induced maps are exact:*

$$\begin{array}{ccccccc} 0 \rightarrow \mathbb{Z}\langle \alpha \times \mu_T \rangle & \rightarrow & H_2(N_\Sigma \setminus \nu(T); \Lambda) & \rightarrow & H_2(N_{\Sigma_n(\alpha, J)}; \Lambda) & \rightarrow & 0, \\ 0 \rightarrow \mathbb{Z}\langle \alpha \times \mu_T \rangle & \rightarrow & H_2(N_\Sigma \setminus \nu(T); \Lambda) & \rightarrow & H_2(N_\Sigma; \Lambda) & \rightarrow & 0. \end{array}$$

Proof. As we noted in Lemma 8.2 that $\pi_1(N_{\Sigma_n(\alpha, J)}) = \mathbb{Z}$ and since $N_{\Sigma_n(\alpha, J)}$ has ribbon boundary, we know that $H_1(N_{\Sigma_n(\alpha, J)}; \Lambda) = 0$ as well as $H_3(N_{\Sigma_n(\alpha, J)}; \Lambda) = 0$, again by Lemma 3.2.

Consequently, the Mayer-Vietoris sequence for $N_{\Sigma_n(\alpha, J)} = N_\Sigma \setminus \nu(T) \cup_{\varphi_n} (S^1 \times E_J)$ with Λ coefficients gives

$$\begin{aligned} 0 \rightarrow H_2(\partial\bar{\nu}(T); \Lambda) &\rightarrow H_2(N_\Sigma \setminus \nu(T); \Lambda) \oplus H_2(S^1 \times E_J; \Lambda) \rightarrow H_2(N_{\Sigma_n(\alpha, J)}; \Lambda) \\ &\rightarrow H_1(\partial\bar{\nu}(T); \Lambda) \rightarrow H_1(N_\Sigma \setminus \nu(T); \Lambda) \oplus H_1(S^1 \times E_J; \Lambda) \rightarrow 0. \end{aligned}$$

Since there is a homotopy equivalence $\bar{\nu}(T) \simeq \alpha \times \mu_\Sigma$, the discussion of coefficient systems from Remark 8.5 implies that $H_i(\partial\bar{\nu}(T); \Lambda) = H_i(\alpha \times \mu_\Sigma \times \mu_T; \Lambda) = H_i(\alpha \times \mu_T)$ for $i = 1, 2$. We saw in Lemma 8.6 that $H_2(E_J \times S^1; \Lambda) = 0$ and $H_1(E_J \times S^1; \Lambda) = \mathbb{Z}$. The previous sequence therefore reduces to

$$(24) \quad \begin{aligned} 0 \rightarrow \mathbb{Z}\langle \alpha \times \mu_T \rangle &\rightarrow H_2(N_\Sigma \setminus \nu(T); \Lambda) \rightarrow H_2(N_{\Sigma_n(\alpha, J)}; \Lambda) \rightarrow \mathbb{Z}\langle \alpha \rangle \oplus \mathbb{Z}\langle \mu_T \rangle \\ &\rightarrow H_1(N_\Sigma \setminus \nu(T); \Lambda) \oplus \mathbb{Z} \rightarrow 0. \end{aligned}$$

When $J = U$ is the unknot, $H_1(S^1 \times E_J; \Lambda) = H_1(\alpha) = \mathbb{Z}\langle \alpha \rangle$, and therefore $\mathbb{Z}\langle \mu_T \rangle$ surjects onto $H_1(N_\Sigma \setminus \nu(T); \Lambda)$. In particular, as an abelian group, $H_1(N_\Sigma \setminus \nu(T); \Lambda)$ is cyclic.

Claim 6. The map $\mathbb{Z}\langle \mu_T \rangle \xrightarrow{\iota} H_1(N_\Sigma \setminus \nu(T); \Lambda)$ is an isomorphism.

Proof. Since we already noted that, as an abelian group, $H_1(N_\Sigma \setminus \nu(T); \Lambda)$ is cyclic, it suffices to show that $H_1(N_\Sigma \setminus \nu(T); \Lambda) \otimes_{\mathbb{Z}} \mathbb{Q} \neq 0$. Equivalently, we must establish that

$$H := H_1(N_\Sigma \setminus \nu(T); \mathbb{Q}[t^{\pm 1}]) \neq 0.$$

By way of contradiction, assume that $H = 0$. Use once again $\varepsilon: \Lambda \rightarrow \mathbb{Z}$ to denote the augmentation map and write \mathbb{Z}_ε and \mathbb{Q}_ε for the resulting Λ -module structures. Since $\text{Tor}_2^\Lambda(\mathbb{Z}_\varepsilon, \mathbb{Q}_\varepsilon) = 0$, the universal coefficient spectral sequence applied to $H_1(N_\Sigma \setminus \nu(T); \mathbb{Q}) = H_1(N_\Sigma \setminus \nu(T); \mathbb{Q}[t^{\pm 1}] \otimes_\Lambda \mathbb{Q}_\varepsilon)$ produces the short exact sequence

$$0 \rightarrow H \otimes_\Lambda \mathbb{Q}_\varepsilon \rightarrow H_1(N_\Sigma \setminus \nu(T); \mathbb{Q}) \rightarrow \text{Tor}_1^\Lambda(\mathbb{Z}_\varepsilon, \mathbb{Q}_\varepsilon) \rightarrow 0.$$

We assumed that $H = 0$, so $H \otimes_\Lambda \mathbb{Q}_\varepsilon = 0$, and thus $H_1(N_\Sigma \setminus \nu(T); \mathbb{Q}) \cong \text{Tor}_1^\Lambda(\mathbb{Z}, \mathbb{Q})$. Using Lemma 8.4, $H_1(N_\Sigma \setminus \nu(T); \mathbb{Q}) = \mathbb{Q}^2$. On the other hand, we have $\text{Tor}_1^\Lambda(\mathbb{Z}_\varepsilon, \mathbb{Q}_\varepsilon) = H_1(S^1; \mathbb{Q}) = \mathbb{Q}$. This is a contradiction and the claim is established. \square

Using Claim 6, the penultimate map in (24) is a surjection $\mathbb{Z}^2 \rightarrow \mathbb{Z}^2$ and therefore an isomorphism. The first short exact sequence in the statement of the lemma now follows from the one displayed in (24). The exactness of the second sequence follows from exactness of the first by taking $J = U$ and recalling that $\Sigma_n(\alpha, U) = \Sigma$ by Remark 8.3. \square

Theorem 8.9. *Let $\Sigma \subseteq X$ be a closed \mathbb{Z} -surface, let $\alpha \subseteq \Sigma$ be a simple closed curve, let J be a knot, and let $n \in \mathbb{Z}$. The surfaces Σ and $\Sigma_n(\alpha, J)$ are topologically ambiently isotopic.*

Proof. After an ambient isotopy, we may assume that the surfaces $\Sigma_0 := \Sigma_n(\alpha, J)$ and $\Sigma_1 := \Sigma$ coincide on a disc $D^2 \subseteq \Sigma \cap \Sigma_n(\alpha, J)$ that does not intersect $\alpha \subseteq \Sigma$. Assume that the normal bundles also coincide over this D^2 . Consider the preimage $\tilde{D}^2 \times \mathbb{R}^2 \subseteq \nu\Sigma_i$. This is homeomorphic to an open 4-ball \tilde{D}^4 . Remove this $(\tilde{D}^4, \tilde{D}^2)$ from (X, Σ) and $(X, \Sigma_n(\alpha, J))$ to obtain properly embedded surfaces $\tilde{\Sigma} \subseteq N$ and $\tilde{\Sigma}_n(\alpha, J) \subseteq N$ with the unknot as a common boundary. By construction, $\tilde{\Sigma}_n(\alpha, J)$ is obtained by rim surgery on $\tilde{\Sigma}$. Apply Theorem 8.7 to obtain a rel. boundary homeomorphism of pairs $\Phi: (N, \tilde{\Sigma}_n(\alpha, J)) \rightarrow (N, \tilde{\Sigma})$. Recall that on $H_2(-; \Lambda)$, we have $\Phi_* = \Psi_*$, where Ψ is the degree one map described above Lemma 8.6. Construct a homeomorphism of pairs $(X, \Sigma_n(\alpha, J)) \rightarrow (X, \Sigma)$ by gluing Φ with the identity homeomorphism $(D^4, D^2) \rightarrow (D^4, D^2)$.

We refine this argument to obtain the required ambient isotopy. Use F to denote the isometry induced by these homeomorphisms on $H_2(-; \Lambda)$ of the surface exteriors. In Lemma 8.6, we argued that this isometry fits into the following commutative diagram with exact rows (here we also used Lemma 8.8 to simplify the Mayer-Vietoris sequences):

$$(25) \quad \begin{array}{ccccccc} 0 & \longrightarrow & \mathbb{Z}\langle \alpha \times \mu_T \rangle & \longrightarrow & H_2(N_{\tilde{\Sigma}} \setminus \nu(T); \Lambda) & \longrightarrow & H_2(N_{\tilde{\Sigma}_n(\alpha, J)}; \Lambda) \longrightarrow 0 \\ & & \downarrow \text{Id} & & \downarrow \text{Id} & & \downarrow F = \Psi_* \\ 0 & \longrightarrow & \mathbb{Z}\langle \alpha \times \mu_T \rangle & \longrightarrow & H_2(N_{\tilde{\Sigma}} \setminus \nu(T); \Lambda) & \longrightarrow & H_2(N_{\tilde{\Sigma}}; \Lambda) \longrightarrow 0. \end{array}$$

Recall that $\Sigma_0 := \Sigma_n(\alpha, J)$ and $\Sigma_1 := \Sigma$, so $X_{\Sigma_0} \cong N_{\tilde{\Sigma}_n(\alpha, J)}$ and $X_{\Sigma_1} \cong N_{\tilde{\Sigma}}$. In Lemma 5.10, $F_{\mathbb{Z}}$ was defined by the following commutative diagram with exact rows

$$(26) \quad \begin{array}{ccccccc} 0 & \longrightarrow & \mathbb{Z}^{2g} & \longrightarrow & H_2(X_{\Sigma_0}) & \xrightarrow{p_0} & H_2(X) \longrightarrow 0 \\ & & \downarrow (F \otimes_{\Lambda} \text{Id}_{\mathbb{Z}})_1 & & \downarrow F \otimes_{\Lambda} \text{Id}_{\mathbb{Z}} & & \downarrow F_{\mathbb{Z}} \\ 0 & \longrightarrow & \mathbb{Z}^{2g} & \longrightarrow & H_2(X_{\Sigma_1}) & \xrightarrow{p_1} & H_2(X) \longrightarrow 0. \end{array}$$

By (26), every element x of $H_2(X)$ can be represented by a class in $H_2(X_{\Sigma_0})$, and by (25) every class here can be represented by a surface S in $N_{\tilde{\Sigma}_n(\alpha, J)} \setminus \nu(T) \subseteq N_{\tilde{\Sigma}_n(\alpha, J)} \cong X_{\Sigma_0}$. Since F is induced by the degree one map Ψ , mapping our surface to the same surface S in $N_{\tilde{\Sigma}} \setminus \nu(T) \subseteq N_{\tilde{\Sigma}} \cong X_{\Sigma_1}$ yields $(F \otimes_{\Lambda} \text{Id}_{\mathbb{Z}})([S]) \in H_2(X_{\Sigma_1})$. The inclusion induced map $p_1: H_2(X_{\Sigma_1}) \rightarrow H_2(X)$ then sends $(F \otimes_{\Lambda} \text{Id}_{\mathbb{Z}})([S])$ to x . Therefore $F_{\mathbb{Z}}: H_2(X) \rightarrow H_2(X)$ is indeed the identity, so the first item of Theorem 5.11 implies that Σ_0 and Σ_1 are topologically ambiently isotopic. \square

APPENDIX A. STABLE ISOTOPY FOR SURFACES IN A TOPOLOGICAL 4-MANIFOLD WITH BOUNDARY

The next result is an extension of [BS16, Theorem 5] to the topological case, and allowing nonempty boundary, but restricting the ambient 4-manifolds somewhat. Since we need the given extension, we provide details of the proof. The main ideas are due to [BS16]. We fill in some details in their argument for constructing a map to S^1 in the course of the proof. The case with nonempty boundary was also stated as [JZ18, Proposition 2.13] in the smooth category.

Theorem A.1. *Let N be an oriented, connected, simply connected, compact topological 4-manifold with boundary S^3 . Let Σ be a compact, connected, orientable surface with one boundary component. Let $\Sigma_0, \Sigma_1 \subseteq N$ be two properly embedded, locally flat, oriented surfaces homeomorphic to Σ , and suppose that $\partial\Sigma_0 = \partial\Sigma_1$. Assume that for each i , $\pi_1(N \setminus \Sigma_i) \cong \mathbb{Z}$ is infinite cyclic generated by a meridian to Σ_i . Then some finite number of trivial 1-handle stabilisations results in ambiently isotopic surfaces Σ'_0 and Σ'_1 with $\partial\Sigma_i = \partial\Sigma'_i$ for $i = 0, 1$.*

We begin with a lemma on framings, some of which is a recollection from Section 5. Note that any locally flat embedded surface in a 4-manifold admits a normal bundle with linear structure group [FQ90, Theorem 9.3].

Lemma A.2. *For $i = 0, 1$ the normal bundle of Σ_i is trivial and for any choice of framing there is a well-defined induced (homotopy class of) framing on the normal bundle of $\partial\Sigma_i$. The surface Σ_i is null-homologous in $H_2(N, \partial N)$, and the induced framing on $\partial\Sigma_i$ equals the Seifert framing.*

Proof. Since the surface Σ_i has nonempty boundary and is oriented, its normal bundle is trivial. Two choices of framings on Σ_i differ by a map $\Sigma_i \rightarrow SO(2)$. The difference between the two framings on the boundary $\partial\Sigma_i$ is governed by the homotopy class of the composite

$$S^1 \xrightarrow{\cong} \partial\Sigma_i \rightarrow \Sigma_i \rightarrow SO(2) \cong S^1.$$

Since the boundary determines a commutator in $\pi_1(\Sigma_i)$, the displayed map is null-homotopic. It follows that the two framings induced on $\partial\Sigma_i$ agree up to homotopy. This proves that the induced framing is well-defined. The second sentence was already shown in Lemmas 5.1 and 5.2. \square

Proof of Theorem A.1. Push the boundary of Σ_1 off the boundary of Σ_0 , using the Seifert framing, to arrange that $\partial\Sigma_0$ and $\partial\Sigma_1$ in $\partial N \cong S^3$ are parallel circles. We may assume that there is a collar $\partial N \times I$ with $\partial N \times \{0\} = \partial N$, and an embedding $g: S^1 \times I \times I \hookrightarrow \partial N \times I$ corresponding to the Seifert framing push-off, with

$$\begin{aligned} g(S^1 \times I \times \{i\}) &= \Sigma_i \cap (\partial N \times I), \quad i = 0, 1, \text{ and} \\ g(S^1 \times \{t\} \times I) &\subseteq \partial N \times \{t\} \text{ for all } t \in I. \end{aligned}$$

Let $A \subseteq \partial N \cong S^3$ be the annulus

$$A := g(S^1 \times \{0\} \times I)$$

connecting $\partial\Sigma_0$ and $\partial\Sigma_1$, arising from the trace of the push. Note that A induces the Seifert framing on $\partial\Sigma_i$, for $i = 0, 1$, and that by Lemma A.2 this equals the framing induced by some choice of framing of the normal bundle of Σ_i .




FIGURE 4. A schematic diagram indicating the relation of $g(S^1 \times I \times I) \subseteq \partial N \times I$ to $\partial\Sigma_0$, $\partial\Sigma_1$, and the annulus $A \subseteq S^3$ that joins them.

By an isotopy, arrange further that Σ_0 and Σ_1 intersect transversely in their interiors [FQ90, Theorem 9.5], and since Σ is compact we may also assume there are finitely many intersection points.

Cap off each $\partial\Sigma_i$ with a Seifert surface, and by a small isotopy of the capped-off Σ_1 , arrange that the capped-off surfaces are disjoint in $\partial N \times I$. By Lemma A.2 (and since $H_2(\partial N) = H_2(S^3) = 0$), the capped-off surfaces are null-homologous in $H_2(N)$, and so intersect algebraically zero times. By tubing Σ_0 to itself, that is by 1-handle stabilisations, arrange that Σ_0 and Σ_1 are disjoint. To achieve this, pair up points with opposite signs in $\Sigma_0 \cap \Sigma_1$ and for each pair $\{p, q\}$ choose a path γ in Σ_1 connecting the two intersection points and away from the other intersection points. We can choose these paths to be mutually disjoint, but this is not obligatory. For each pair of points $\{p, q\}$, remove two open discs from Σ_0 , one for each pair of points, and add a tube, coming from the normal circle bundle of Σ_1 restricted to γ , as

shown in Figure 5. In case different paths on Σ_1 intersect, vary the radii of the tubes to keep them disjoint. This stabilises Σ_0 to a surface disjoint from Σ_1 .




FIGURE 5. Stabilising Σ_0 to Σ'_0 by adding a 1-handle in a neighbourhood of an arc γ in Σ_1 .

Perform the same number of trivial 1-handle stabilisations to Σ_1 , so that the surfaces are still abstractly homeomorphic. We will abuse notation and still refer to the resulting surfaces as Σ_0 and Σ_1 . Let

$$N_{\Sigma_0, \Sigma_1} := N \setminus (\nu\Sigma_0 \cup \nu\Sigma_1).$$

We will construct an oriented 3-dimensional relative Seifert manifold $Y \subseteq N_{\Sigma_0, \Sigma_1}$, that is a rel. boundary cobordism between Σ_0 and Σ_1 , embedded in N and with $\partial Y = S$, where

$$S := \Sigma_0 \cup A \cup -\Sigma_1.$$

Claim. There is a locally flat, embedded, compact, orientable 3-manifold Y with $\partial Y = S$ and corners at $\Sigma_i \cap A$, as described in the preceding paragraph.

Proof. The strategy to construct Y is as follows: define a suitable map $\partial N_{\Sigma_0, \Sigma_1} \rightarrow S^1$, extend it to a map $N_{\Sigma_0, \Sigma_1} \rightarrow S^1$ while controlling the restriction to $\partial N_{\Sigma_0, \Sigma_1}$, and then take Y to be the inverse image of a transverse regular point in S^1 .

The first step is to construct a map $\alpha: N_{\Sigma_0, \Sigma_1} \rightarrow S^1$. Recall that $N_{\Sigma_i} := N \setminus \nu\Sigma_i$. We can express $N = N_{\Sigma_0} \cup N_{\Sigma_1}$, with $N_{\Sigma_0, \Sigma_1} = N_{\Sigma_0} \cap N_{\Sigma_1}$. A portion of the resulting Mayer-Vietoris sequence (with \mathbb{Z} coefficients) is

$$H_2(N) \xrightarrow{\delta} H_1(N_{\Sigma_0, \Sigma_1}) \rightarrow H_1(N_{\Sigma_0}) \oplus H_1(N_{\Sigma_1}) \rightarrow H_1(N).$$

By hypothesis $H_1(N) = 0$ and $H_1(N_{\Sigma_i}) \cong \mathbb{Z}$, so that we obtain a short exact sequence

$$0 \rightarrow \text{im } \delta \rightarrow H_1(N_{\Sigma_0, \Sigma_1}) \rightarrow \mathbb{Z} \oplus \mathbb{Z} \rightarrow 0.$$

Since $\mathbb{Z} \oplus \mathbb{Z}$ is free abelian, this splits and we have that

$$H_1(N_{\Sigma_0, \Sigma_1}) \cong \mathbb{Z} \oplus \mathbb{Z} \oplus \text{im } \delta.$$

By the hypotheses of Theorem A.1, the first two summands are generated by meridians μ_{Σ_0} and μ_{Σ_1} to the surfaces Σ_0 and Σ_1 respectively. Consider the dual element

$$\alpha := \mu_{\Sigma_0}^* - \mu_{\Sigma_1}^* \in \text{Hom}(H_1(N_{\Sigma_0, \Sigma_1}), \mathbb{Z}) \cong H^1(N_{\Sigma_0, \Sigma_1}) \cong [N_{\Sigma_0, \Sigma_1}, S^1],$$

sending the $\text{im } \delta$ summand identically to 0. We will also write $\alpha: N_{\Sigma_0, \Sigma_1} \rightarrow S^1$ for a corresponding representing map. We use that since N_{Σ_0, Σ_1} is a topological 4-manifold, it is

homotopy equivalent to a CW-complex [FNOP19, Theorem 4.5], and therefore we may identify $H^1(N_{\Sigma_0, \Sigma_1}) \cong [N_{\Sigma_0, \Sigma_1}, S^1]$.

Now we consider the restriction to $\partial N_{\Sigma_0, \Sigma_1}$ under the inclusion-induced map

$$\iota^*: H^1(N_{\Sigma_0, \Sigma_1}) \rightarrow H^1(\partial N_{\Sigma_0, \Sigma_1}) \cong [\partial N_{\Sigma_0, \Sigma_1}, S^1],$$

where $\iota: \partial N_{\Sigma_0, \Sigma_1} \rightarrow N_{\Sigma_0, \Sigma_1}$ is the inclusion map. We have a decomposition

$$\partial N_{\Sigma_0, \Sigma_1} = \Sigma_0 \times S^1 \cup \partial_0 N_{\Sigma_0, \Sigma_1} \cup \Sigma_1 \times S^1.$$

An elementary Mayer-Vietoris argument yields that

$$H^1(\partial N_{\Sigma_0, \Sigma_1}) \cong \mathbb{Z} \oplus \mathbb{Z} \oplus \mathbb{Z}^{2g} \oplus \mathbb{Z}^{2g}.$$

The first two summands are generated by duals to the meridians, $\mu_{\Sigma_0}^*$ and $\mu_{\Sigma_1}^*$. By our work with framings above, for $i = 0, 1$ there is a framing of the normal bundle of Σ_i , determining an identification of the tubular neighbourhood $\bar{\nu}\Sigma_i$ with $\Sigma_i \times D^2$, that agrees on $\partial\Sigma_i$ with a corresponding identification induced from the Seifert framing on $\partial\Sigma_i$. This determines an identification

$$\partial\bar{\nu}\Sigma_i \setminus \nu\partial\Sigma_i \cong \Sigma_i \times S^1.$$

The two \mathbb{Z}^{2g} summands of $H^1(\partial N_{\Sigma_0, \Sigma_1})$ are generated by dual classes to curves of the form $\gamma_k \times \{-1\}$ where γ_k is simple closed curve forming part of a symplectic basis for $H_1(\Sigma_i)$, for some i . Here we use the chosen framing of the normal bundle of Σ_i to fix representatives for the \mathbb{Z}^{2g} summands.

The restriction $\iota^*\alpha \in H^1(\partial N_{\Sigma_0, \Sigma_1})$ is $(1, -1, x, y)$, for some $x, y \in \mathbb{Z}^{2g}$. But by changing the choice of framing of $\nu\Sigma_i$ along the curves γ_k , we can arrange that $x = y = 0$. By changing the framing of the normal bundle of Σ_i by some number of full twists along some basis curve γ_k in $H_1(\Sigma_i)$, we change the meaning of $\gamma_k \times \{-1\}$ in the previous paragraph.

Let us explain the operation of ‘‘changing the framing’’ in more detail. Such a change is occasioned by the action of $[\cup_k \gamma_k, SO(2)]$ on the set of framings of the normal bundle, to alter the given framing on the union of the curves $\{\gamma_k\}$ by some number of full twists for each curve. Any such alteration automatically extends over the 2-skeleton since the attaching map of the 2-cell of Σ_i is a commutator in the γ_k times $\partial\Sigma_i$. Any two choices of extension over the 2-cells are homotopic, since $\pi_2(SO(2)) = 0$. Therefore we have a well-defined notion of altering the framing along the curves γ_k . This changes the entry of (x, y) corresponding to γ_k , since the map $\alpha: N_{\Sigma_0, \Sigma_1} \rightarrow S^1$ now sends $\gamma_k \times \{-1\}$ to a curve representing a different element of $H_1(S^1)$. By Lemma A.2, changing the choice of framing on a basis element for $H_1(\Sigma_i)$ does not change the induced framing on the boundary.

Now we define a map $f: \partial N_{\Sigma_0, \Sigma_1} \rightarrow S^1$. On $\Sigma_i \times S^1 \subseteq \partial N_{\Sigma_0, \Sigma_1}$, define the map to S^1 by the projection $f|_{\Sigma_i \times S^1}: \Sigma_i \times S^1 \rightarrow S^1$ onto the second factor. On the remainder of $\partial N_{\Sigma_0, \Sigma_1}$, namely

$$\partial_0 N_{\Sigma_0, \Sigma_1} := \partial N \setminus (\nu\Sigma_0 \cup \nu\Sigma_1),$$

or in other words the link exterior $S^3 \setminus (\nu\partial\Sigma_0 \cup \nu\partial\Sigma_1)$, define a Pontryagin-Thom style collapse map $f|_{\partial_0 N_{\Sigma_0, \Sigma_1}}: \partial_0 N_{\Sigma_0, \Sigma_1} \rightarrow S^1$ by choosing a tubular neighbourhood $A \times [-1, 1] \subseteq \partial_0 N_{\Sigma_0, \Sigma_1}$ and sending $(a, x) \mapsto e^{\pi i x} \in S^1$ for $a \in A$ and $x \in [-1, 1]$, then sending $\partial_0 N_{\Sigma_0, \Sigma_1} \setminus (A \times [-1, 1])$ to $-1 \in S^1$. Then $f|_{\partial_0 N_{\Sigma_0, \Sigma_1}}^{-1}(\{1\}) = A \cap \partial_0 N_{\Sigma_0, \Sigma_1}$. Note that we may assume that our maps to S^1 agree on the torus overlaps $\partial\Sigma_i \times S^1$, since the framings on $\partial\Sigma_i$ all agree up to homotopy.

This completes the construction of a map $f: \partial N_{\Sigma_0, \Sigma_1} \rightarrow S^1$. Note that $f: \partial N_{\Sigma_0, \Sigma_1} \rightarrow S^1$ corresponds to the element

$$(1, -1, 0, 0) \in \mathbb{Z} \oplus \mathbb{Z} \oplus \mathbb{Z}^{2g} \oplus \mathbb{Z}^{2g}.$$

Therefore the cohomology classes $\iota^* \alpha$ and f agree in $H^1(\partial N_{\Sigma_0, \Sigma_1}) \cong [\partial N_{\Sigma_0, \Sigma_1}, S^1]$, and so by a homotopy of α supported in a collar of $\partial N_{\Sigma_0, \Sigma_1}$ we obtain a map $F: N_{\Sigma_0, \Sigma_1} \rightarrow S^1$ with $F|_{\partial N_{\Sigma_0, \Sigma_1}} = f: \partial N_{\Sigma_0, \Sigma_1} \rightarrow S^1$.

The inverse image under F of a transverse regular point in S^1 yields a 3-dimensional relative Seifert manifold, locally flatly embedded in $N_{\Sigma_0, \Sigma_1} = N \setminus (\nu \Sigma_0 \cup \nu \Sigma_1)$. See [FNO19, Section 10.2] for information on map transversality in the topological category. Add collars $\Sigma_0 \times I$ and $\Sigma_1 \times I$ in $\nu \Sigma_0$ and $\nu \Sigma_1$ respectively, to obtain the 3-manifold Y that we seek. This completes the proof of the claim. \square

Morse theory on Y gives rise to a Heegaard decomposition relative to the annulus A . The Heegaard surface can be obtained from both Σ_0 and Σ_1 by 1-handle stabilisations and ambient isotopy.

Since by our assumptions $\pi_1(N_{\Sigma_i})$ is generated by a meridian to Σ_i , Boyle's [Boy88] proof shows that every 1-handle stabilisation is a trivial stabilisation. He applied [Hud72, Theorem 4] of Hudson for the statement that D^1 cores of handle additions that are homotopic rel. endpoints are in fact smoothly ambiently isotopic fixing the endpoints. To apply Boyle's work to topologically embedded surfaces in a compact 4-manifold N , remove a point from N and smooth $N \setminus \{\text{pt}\}$ in such a way that Σ is smoothly embedded. Then Boyle's application of Hudson's result yields a smooth ambient isotopy, which gives rise to a topological ambient isotopy once the point is added back to N . We note that Boyle works in S^4 , but his proof applies to any oriented ambient 4-manifold.

We therefore have that after finitely many trivial stabilisations, Σ_0 and Σ_1 are ambiently isotopic in N relative to the constant isotopy on the boundary, as desired. \square

REFERENCES

- [Bas73] Hyman Bass. Unitary algebraic K -theory. In *Algebraic K-theory, III: Hermitian K-theory and geometric applications (Proc. Conf., Battelle Memorial Inst., Seattle, Wash., 1972)*, pages 57–265. Lecture Notes in Math., Vol. 343, 1973.
- [BF14] Maciej Borodzik and Stefan Friedl. On the algebraic unknotting number. *Trans. London Math. Soc.*, 1(1):57–84, 2014.
- [BF15] Maciej Borodzik and Stefan Friedl. The unknotting number and classical invariants, I. *Algebr. Geom. Topol.*, 15(1):85–135, 2015.
- [Boy86] Steven Boyer. Simply-connected 4-manifolds with a given boundary. *Trans. Amer. Math. Soc.*, 298(1):331–357, 1986.
- [Boy88] Jeffrey Boyle. Classifying 1-handles attached to knotted surfaces. *Trans. Amer. Math. Soc.*, 306(2):475–487, 1988.
- [Bre93] Glen E. Bredon. *Topology and geometry*, volume 139 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1993.
- [BS16] R. İnanç Baykur and Nathan Sunukjian. Knotted surfaces in 4-manifolds and stabilizations. *J. Topol.*, 9(1):215–231, 2016.
- [CFT18] Anthony Conway, Stefan Friedl, and Enrico Toffoli. The Blanchfield pairing of colored links. *Indiana Univ. Math. J.*, 67(6):2151–2180, 2018.
- [CLM] Diarmuid Crowley, Wolfgang Lueck, and Tibor Macko. Surgery theory: Foundations (11-2020 notes). <http://thales.doa.fmph.uniba.sk/macko/sb-www-19-11-2020.pdf>.
- [COT04] Tim D. Cochran, Kent E. Orr, and Peter Teichner. Structure in the classical knot concordance group. *Comment. Math. Helv.*, 79(1):105–123, 2004.

- [CP19] Anthony Conway and Mark Powell. Characterisation of homotopy ribbon discs. *ArXiv 1902.05321*, 2019.
- [Cro02] Diarmuid Crowley. On the classification of highly connected manifolds in dimensions 7 and 15. 2002. Indiana University.
- [CS11] Diarmuid Crowley and Joerg Sixt. Stably diffeomorphic manifolds and $l_{2q+1}(\mathbb{Z}[\pi])$. *Forum Math.*, 23(3):483–538, 2011.
- [FLNP17] Stefan Friedl, Constance Leidy, Matthias Nagel, and Mark Powell. Twisted Blanchfield pairings and decompositions of 3-manifolds. *Homology Homotopy Appl.*, 19(2):275–287, 2017.
- [FM12] Benson Farb and Dan Margalit. *A primer on mapping class groups*, volume 49 of *Princeton Mathematical Series*. Princeton University Press, Princeton, NJ, 2012.
- [FNOP19] Stefan Friedl, Matthias Nagel, Patrick Orson, and Mark Powell. A survey of the foundations of four-manifold theory in the topological category. *ArXiv 1910.07372*, 2019.
- [FP17] Stefan Friedl and Mark Powell. A calculation of Blanchfield pairings of 3-manifolds and knots. *Moscow Math. J.*, 17:59–77, 2017.
- [FQ90] Michael H. Freedman and Frank Quinn. *Topology of 4-manifolds*, volume 39 of *Princeton Mathematical Series*. Princeton University Press, Princeton, NJ, 1990.
- [FS97] Ronald Fintushel and Ronald J. Stern. Surfaces in 4-manifolds. *Math. Res. Lett.*, 4(6):907–914, 1997.
- [GS99] Robert E. Gompf and András I. Stipsicz. *4-manifolds and Kirby calculus*, volume 20 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 1999.
- [HK95] Jonathan A. Hillman and Akio Kawauchi. Unknotting orientable surfaces in the 4-sphere. *J. Knot Theory Ramifications*, 4(2):213–224, 1995.
- [HKT09] Ian Hambleton, Matthias Kreck, and Peter Teichner. Topological 4-manifolds with geometrically two-dimensional fundamental groups. *J. Topol. Anal.*, 1(2):123–151, 2009.
- [HT97] Ian Hambleton and Peter Teichner. A non-extended Hermitian form over $\mathbf{Z}[\mathbf{Z}]$. *Manuscripta Math.*, 93(4):435–442, 1997.
- [Hud72] John Hudson. Embeddings of bounded manifolds. *Proc. Cambridge Philos. Soc.*, 72:11–20, 1972.
- [JMZ20] András Juhász, Maggie Miller, and Ian Zemke. Transverse invariants and exotic surfaces in the 4-ball. *ArXiv 2001.07191*, 2020.
- [JZ18] András Juhász and Ian Zemke. Stabilization distance bounds from link Floer homology. *ArXiv:1810.09158*, 2018.
- [Kaw94] Akio Kawauchi. Splitting a 4-manifold with infinite cyclic fundamental group. *Osaka J. Math.*, 31(3):489–495, 1994.
- [Kaw13] Akio Kawauchi. Splitting a 4-manifold with infinite cyclic fundamental group, revised. *J. Knot Theory Ramifications*, 22(14):1350081, 9, 2013.
- [Kha17] Qayum Khan. Cancellation for 4-manifolds with virtually abelian fundamental group. *Topology Appl.*, 220:14–30, 2017.
- [Kim06] Hee Jung Kim. Modifying surfaces in 4-manifolds by twist spinning. *Geom. Topol.*, 10:27–56, 2006.
- [Kir97] Problems in low-dimensional topology. In Rob Kirby, editor, *Geometric topology (Athens, GA, 1993)*, volume 2 of *AMS/IP Stud. Adv. Math.*, pages 35–473. Amer. Math. Soc., Providence, RI, 1997.
- [Knu91] Max-Albert Knus. *Quadratic and Hermitian forms over rings*, volume 294 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, 1991. With a foreword by I. Bertuccioni.
- [KPT20] Daniel Kasprowski, Mark Powell, and Peter Teichner. Algebraic criteria for stable diffeomorphism of spin 4-manifolds. *ArXiv 2006.06127*, 2020.
- [KR08a] Hee Jung Kim and Daniel Ruberman. Smooth surfaces with non-simply-connected complements. *Algebr. Geom. Topol.*, 8(4):2263–2287, 2008.
- [KR08b] Hee Jung Kim and Daniel Ruberman. Topological triviality of smoothly knotted surfaces in 4-manifolds. *Trans. Amer. Math. Soc.*, 360(11):5869–5881, 2008.
- [Kre79] Matthias Kreck. Isotopy classes of diffeomorphisms of $(k - 1)$ -connected almost-parallelizable $2k$ -manifolds. In *Algebraic topology, Aarhus 1978 (Proc. Sympos., Univ. Aarhus, Aarhus, 1978)*, volume 763 of *Lecture Notes in Math.*, pages 643–663. Springer, Berlin, 1979.
- [Kre99] Matthias Kreck. Surgery and duality. *Ann. of Math. (2)*, 149(3):707–754, 1999.

- [Lam06] T. Y. Lam. *Serre's problem on projective modules*. Springer Monographs in Mathematics. Springer-Verlag, Berlin, 2006.
- [Lev77] Jerome Levine. Knot modules. I. *Trans. Amer. Math. Soc.*, 229:1–50, 1977.
- [Liv82] Charles Livingston. Surfaces bounding the unlink. *Michigan Math. J.*, 29(3):289–298, 1982.
- [Mar13] Thomas E. Mark. Knotted surfaces in 4-manifolds. *Forum Math.*, 25(3):597–637, 2013.
- [MvdKV88] Bruce Magurn, Wilberd van der Kallen, and Leonid Vaserstein. Absolute stable rank and Witt cancellation for noncommutative rings. *Invent. Math.*, 91(3):525–542, 1988.
- [Oba17] Takahiro Oba. Surfaces in D^4 with the same boundary and fundamental group. *ArXiv 2006.06127*, 2017.
- [Per86] Bernard Perron. Pseudo-isotopies et isotopies en dimension quatre dans la catégorie topologique. *Topology*, 25(4):381–397, 1986.
- [Qui86] Frank Quinn. Isotopy of 4-manifolds. *J. Differential Geom.*, 24(3):343–372, 1986.
- [Ran81] Andrew Ranicki. *Exact sequences in the algebraic theory of surgery*, volume 26 of *Mathematical Notes*. Princeton University Press, Princeton, N.J.; University of Tokyo Press, Tokyo, 1981.
- [Sun15] Nathan S. Sunukjian. Surfaces in 4-manifolds: concordance, isotopy, and surgery. *Int. Math. Res. Not. IMRN*, (17):7950–7978, 2015.
- [SW00] Richard Stong and Zhenghan Wang. Self-homeomorphisms of 4-manifolds with fundamental group \mathbb{Z} . *Topology Appl.*, 106(1):49–56, 2000.

MAX-PLANCK-INSTITUT FÜR MATHEMATIK, BONN, GERMANY
Email address: anthonyconway@gmail.com

DEPARTMENT OF MATHEMATICAL SCIENCES, DURHAM UNIVERSITY, LOWER MOUNTJOY, UNITED KINGDOM
Email address: mark.a.powell@durham.ac.uk