

A survey of the foundations of four-manifold theory in the topological category

Stefan Friedl

Matthias Nagel

Patrick Orson

Mark Powell

FAKULTÄT FÜR MATHEMATIK, UNIVERSITÄT REGENSBURG, GERMANY

Email address: `sfriedl@gmail.com`

Email address: `matthinagel@gmail.com`

CALIFORNIA POLYTECHNIC STATE UNIVERSITY, SAN LUIS OBISPO, CA, USA

Email address: `patrickorson@gmail.com`

SCHOOL OF MATHEMATICS AND STATISTICS, UNIVERSITY OF GLASGOW, UK

Email address: `mark.powell@glasgow.ac.uk`

ABSTRACT. This survey aims to provide a guide to the literature on topological 4-manifolds. Foundational theorems on 4-manifolds are stated, especially in the topological category. Precise references are given, with indications of the strategies employed in the proofs. Where appropriate we give statements for manifolds of all dimensions.

Many intuitively plausible theorems which are standard results in differential topology are either extraordinarily deep results in the topological category, are open, or are known to be false. Hence one must proceed with caution. This book seeks to help 4-manifold topologists navigate potential pitfalls, and to apply the many powerful results that do exist with confidence.

CHAPTER 1

Introduction

Our aim in this survey is to provide readers who have trained in algebraic topology and perhaps differential topology or Riemannian geometry, with a guide to the literature on topological manifolds. We aim to state some foundational theorems in topological manifolds, with a strong bias towards dimension four. If we were considering manifolds with a smooth atlas, many of these statements would be familiar textbook-level tools. In the topological category, navigating which of these tools can still be used, and where to find proofs of these facts, can be a challenging endeavour. On the other hand, particularly in dimension four, some of the results we describe are not familiar results in the smooth category and often no result of the sort holds for smooth manifolds. This second type of statement demonstrates one of the attractions of working with 4-manifolds in the topological category, where major classification statements can be achieved.

Our hope is that with the statements from this book the “working topologist” will be equipped to handle most situations. We make no claims of originality.

1.1. HIGH-DIMENSIONAL TOPOLOGICAL MANIFOLDS

Though we have a strong bias in this survey towards thinking about 4-dimensional manifolds, in order to do this in the topological category one must have a good understanding of manifold topology in higher dimensions.

Spectacular results on the classification of smooth manifolds in dimensions ≥ 5 arose from Surgery Theory, due to Smale [Sma62], Kervaire–Milnor [KM63], Browder [Bro72], Novikov [Nov64], Sullivan [Sul66], and Wall [Wal99], among others. These methods were extended to topological manifolds by Newman [New66a], Kirby [Kir69], and Siebenmann [KS77] in dimensions at least five, and to dimension four by Freedman and Quinn [Fre82, Qui82, FQ90]. To paraphrase Andrew Ranicki,

Smooth manifolds in dimensions at least five exhibit a beautiful correspondence between geometry and algebra. Topological manifolds in dimension at least four are in this sense like smooth high-dimensional manifolds, but even more so.

To what does “even more so” refer? One instance is the principle that topological manifolds are governed by their homotopy type which, in turn, is often governed by algebraic invariants. Many specific instances of this principle hold uniformly across all

dimensions for topological manifolds and we highlight a few of these now to demonstrate the point. Most famous is the (topological) Poincaré conjecture, which characterises the sphere S^n , up to homeomorphism, in terms of algebraic topological invariants. This result is known to be true in all dimensions, due to Newman, Freedman, and Perelman [New66b, Fre82, MT07]. Locally flat topological embeddings of spheres in spheres are also well understood in these terms. In codimension zero, every orientation preserving homeomorphism of S^n is isotopic to the identity, due to Fisher, Kirby, and Quinn [Fis60, Kir69, Qui82]. For codimension 1, we have the Schoenflies conjecture, that every locally flat embedding $S^{n-1} \subseteq S^n$ is trivial. This is true in all dimensions, and due to Brown, Mazur, and Morse [Bro60, Maz59, Mor60]. A codimension two locally flat embedding $S^{n-2} \subseteq S^n$ is topologically unknotted if and only if the complement is homotopy equivalent to S^1 , due to Papakyriakopoulos [Pap57], Stallings [Sta63], and Freedman-Quinn [FQ90, Theorem 11.7A]. Finally for high codimension, every knot $S^{n-k} \subseteq S^n$ is trivial, for $k \geq 3$, which is again due to Stallings [Sta63], cf. [Zee63].

For manifold classification results, there is also an especially close correspondence between topology and algebra in the topological category in dimensions at least four. Wall [Wal62], Freedman, and Quinn [FQ90] proved that topological manifolds of dimension $2n$ that are $(n-1)$ -connected are classified up to homeomorphism by their intersection forms on the n th homology groups, together with a quadratic extension. The Borel conjecture holds in many cases [AFW15, Lüc10]. In particular for every $n \geq 1$, every homotopy equivalence $M^n \rightarrow T^n = (S^1)^n$ is homotopic to a homeomorphism, see [FQ90, p. 205] and [Lüc10]. Finally, an important example, though somewhat more specialised, is that the Surgery Exact Sequence becomes an exact sequence of abelian groups in the topological category, due to Quinn [Qui70] and Nicas [Nic79], with a purely algebraic formulation due to Ranicki [Ran92].

These examples demonstrate a tight connection between the study of topological manifolds and the associated homotopy types, and related algebra. In dimension four, many of the significant classification results we state in this survey will also hew to this principle.

1.2. A FOCUS ON DIMENSION FOUR

In the Ranicki quotation paraphrased above, the only dimension not common to both ranges is dimension four. Indeed, most of the deep classification results that are known about 4-manifolds are only possible in the topological category. Studying topological 4-manifolds combines the visual nature of low-dimensional topology, with the ability to apply powerful high-dimensional methods to classification problems.

An additional special feature of dimension four is that here the contrast between the smooth and topological manifold categories is starkest. Indeed, the indications are that smooth 4-manifolds fail in every way imaginable to exhibit a close correspondence to their underlying homotopy type. Although in this survey we focus very heavily on topological 4-manifolds, we will hint at this dramatic divergence in some places, for example in Chapter 8.

The geometric topologist who wishes to study topological 4-manifolds, with all the rich rewards suggested above, must navigate the challenge that even basic results from differential topology are either false, unknown, or extraordinarily deep results in the absence of a smooth atlas. For example it is not true in general that topological submanifolds admit tubular neighbourhoods, and in cases where this is known, such as for codimension two submanifolds, the proofs use all the available technology developed by Kirby-Siebenmann and Freedman-Quinn. As another example, it was not known until Quinn's work in 1982 that connected sum of topological 4-manifolds is a well-defined operation.

1.3. WHAT IS IN THIS SURVEY?

This brings us, at last, to one of the main practical purposes of this survey. We seek to clarify exactly which of the familiar tools of geometric topology are available in the topological category, and to provide a precise guide for where to find proofs. Below, we give a sample of the statements discussed in this book. (Here, and throughout the book, “manifold” refers to what is often called a “topological manifold”; see Chapter 2 for a precise definition.)

- (1) Existence and uniqueness of collar neighbourhoods (Theorem 2.16).
- (2) The Isotopy Extension Theorem 2.20.
- (3) Existence of CW structures (Theorem 3.16).
- (4) Multiplicativity of the Euler characteristic under finite covers (Corollary 3.18).
- (5) The Annulus Theorem 4.1 and the Stable Homeomorphism Theorem 4.3.
- (6) Connected sum of two oriented connected 4-manifolds is well-defined (Theorem 4.12).
- (7) Existence and uniqueness of tubular neighbourhoods of submanifolds (Theorems 5.5 and 5.6).
- (8) Stiefel-Whitney classes for topological manifolds (Chapter 7).
- (9) Intersection forms of compact, connected, oriented 4-manifolds are even (Proposition 8.4).
- (10) Noncompact connected 4-manifolds admit a smooth structure (Theorem 9.1).
- (11) When the Kirby-Siebenmann invariant of a connected 4-manifold vanishes, both connected sum with copies of $S^2 \times S^2$ and taking the product with \mathbb{R} yield smoothable manifolds (Theorem 9.9).
- (12) Transversality for submanifolds and for maps (Theorems 10.3 and 10.10).
- (13) Codimension one and two homology classes can be represented by submanifolds (Theorem 10.17).
- (14) Classification of 4-manifolds up to homeomorphism with trivial and cyclic fundamental groups (Chapter 12).
- (15) Compact orientable manifolds that are homeomorphic are stably diffeomorphic (Theorem 13.2 and Corollary 13.4).
- (16) Multiplicativity of signatures under finite covers (Theorem 8.6).
- (17) The definition of Reidemeister torsion for compact manifolds and some of its key technical properties (Section 14.3).

- (18) Obstructions to concordance of knots and links (Theorem 15.2).
- (19) Poincaré duality for compact manifolds (with possibly non-empty boundary) with twisted coefficients (Theorems A.15 and A.16).

Remark 1.1. We make a remark here about a notable *omission* from this survey. We have not written in any great detail about non-compact 4-manifolds. This is itself a rich topic, exhibiting some of the most exciting and distinctive features of dimension 4. However, it was decided to be beyond the scope of the current survey.

Conventions.

- (1) Given a subset A of a topological space X we denote the interior by $\text{Int } A$.
- (2) For $n \in \mathbb{N}_0$, $x \in \mathbb{R}^n$ and $r \geq 0$ we write $D_r^n(x) = \{y \in \mathbb{R}^n \mid \|y - x\| \leq r\}$. We write $D_r^n = D_r^n(0)$ and we write $D^n = D_1^n(0)$ for the closed unit ball in \mathbb{R}^n . We refer to $\text{Int } D^n = \{x \in \mathbb{R}^n \mid \|x\| < 1\}$ as the open n -ball.
- (3) Unless indicated otherwise I denotes the interval $I = [0, 1]$.
- (4) All maps between topological spaces are understood to be continuous.
- (5) A topological space X is called *simply connected* if it is nonempty, path-connected and if the fundamental group is trivial.
- (6) On several occasions we use cup and cap products and we cite several results from [Dol95, Bre97, Hat02, Fri23]. Different books on algebraic topology often have different sign conventions for cup and cap products, but in all statements that we give, the sign conventions are irrelevant, so it is not a problem to mix results from different sources.

Acknowledgments. Thanks to Steve Boyer, Anthony Conway, Jim Davis, Mauricio Gómez López, Fabian Hebestreit, Jonathan Hillman, Min Hoon Kim, Alexander Kupers, Markus Land, Tye Lidman, Chuck Livingston, Ciprian Manolescu, Erik Pedersen, George Raptis, Arunima Ray, and Eamonn Tweedy for very helpful conversations.

We are particularly grateful to Gerrit Herrmann for providing us with detailed notes which form the basis of Appendix A.

Jim Davis brought the Ranicki quotation above to our attention, and discussed its significance with us. He also provided crucial assistance with the proof of Theorem 9.2.

We wish to thank the referee for an extraordinarily thorough and helpful report.

SF was supported by the SFB 1085 ‘Higher Invariants’ at the University of Regensburg, funded by the Deutsche Forschungsgemeinschaft (DFG). MN gratefully acknowledges support by the SNSF Grant 181199. MP was supported by an NSERC Discovery Grant, EPSRC New Investigator grant EP/T028335/2 and EPSRC New Horizons grant EP/V04821X/2. SF wishes to thank the Université du Québec à Montréal, Durham University, Glasgow University, and Cal Poly for hospitality.

Contents

Chapter 1. Introduction	3
1.1. High-dimensional topological manifolds	3
1.2. A focus on dimension four	4
1.3. What is in this survey?	5
Chapter 2. Manifolds	9
2.1. Definition of manifolds	9
2.2. Definition of submanifolds	10
2.3. Immersions	13
2.4. Collar neighbourhoods	14
2.5. The Isotopy Extension Theorem	15
Chapter 3. CW structures, triangulations, and handle structures on manifolds	17
Chapter 4. The Annulus Theorem and the Stable Homeomorphism Theorem	27
4.1. The Stable Homeomorphism Theorem	27
4.2. The connected sum operation	31
4.3. The Product Structure Theorem	33
Chapter 5. Tubular neighbourhoods	35
5.1. Tubular neighbourhoods: existence and uniqueness	35
5.2. Normal vector bundles	37
5.3. Tubular neighbourhoods: proofs	40
Chapter 6. Background on bundle structures	43
6.1. Topological, smooth, and piecewise linear \mathbb{R}^n -bundles	43
6.2. Microbundles	48
Chapter 7. Stiefel-Whitney classes	53
Chapter 8. Intersection forms and smooth 4-manifolds	63
8.1. Intersection forms	63
8.2. Intersection forms of spin manifolds	64
8.3. Twisted intersection forms and twisted signatures	65
8.4. Intersection forms of smooth 4-manifolds	67
Chapter 9. Smoothing 4-manifolds	71

9.1. Smoothing noncompact 4-manifolds.....	71
9.2. The Kirby-Siebenmann invariant and stable smoothing of 4-manifolds.....	72
Chapter 10. Topological transversality.....	81
10.1. Transversality for submanifolds.....	82
10.2. Transversality for maps.....	85
10.3. Representing homology classes by submanifolds.....	87
Chapter 11. Tubing of surfaces.....	95
Chapter 12. Classification results for 4-manifolds.....	99
12.1. Simply connected 4-manifolds.....	99
12.2. Non simply-connected 4-manifolds.....	100
Chapter 13. Stable smoothing of homeomorphisms.....	105
13.1. Kreck's modified surgery.....	107
13.2. Stable diffeomorphism of homeomorphic orientable 4-manifolds.....	110
13.3. Nonorientable 4-manifolds and stable diffeomorphism.....	112
13.4. Homeomorphic 4-manifolds with boundary.....	114
Chapter 14. Reidemeister torsion in the topological category.....	115
14.1. The simple homotopy type of a manifold.....	115
14.2. The cellular chain complex and Poincaré triads.....	119
14.3. Reidemeister torsion.....	120
Chapter 15. Obstructions to being topologically slice.....	123
15.1. The Fox-Milnor Theorem.....	123
15.2. A proof of the Fox-Milnor Theorem.....	124
Appendix A. Poincaré Duality with twisted coefficients.....	131
A.1. Twisted homology and cohomology groups.....	131
A.2. The Poincaré Duality Theorem.....	135
A.3. Preparations for the proof of the Twisted Poincaré Duality Theorem.....	137
A.4. The main technical theorem.....	139
A.5. Proof of the Twisted Poincaré Duality Theorem.....	144
Bibliography.....	147

CHAPTER 2

Manifolds

In this chapter we introduce the very basic notions of a manifold, of a submanifold, of locally flat embeddings and of immersions. We also state two fundamental results, namely the Collar Neighbourhood Theorem 2.16 and the Isotopy Extension Theorem 2.20.

2.1. DEFINITION OF MANIFOLDS

In the literature the notion of a “manifold” gets defined differently, depending on the preferences of the authors. Thus we state here what we mean by a manifold.

Definition 2.1. Let X be a topological space.

- (1) We say that X is *second countable* if there exists a countable basis for the topology.
- (2) An *n -dimensional chart for X at a point $x \in X$* is a homeomorphism $\Phi: U \rightarrow V$ where U is an open neighbourhood of x and
 - (i) V is an open subset of \mathbb{R}^n or
 - (ii) V is an open subset of the half-space $H_n = \{(x_1, \dots, x_n) \in \mathbb{R}^n \mid x_n \geq 0\}$ and $\Phi(x)$ lies on $E_{n-1} = \{(x_1, \dots, x_n) \in \mathbb{R}^n \mid x_n = 0\}$.

In the former case we say that Φ is a *chart of type (i)*; in the latter case we say that Φ is a *chart of type (ii)*.

- (3) We say that X is an *n -dimensional manifold* if X is second countable and Hausdorff, and if for every $x \in X$ there exists an n -dimensional chart $\Phi: U \rightarrow V$ at x .
- (4) We say that a point x on a manifold is a *boundary point* if x admits a chart of type (ii). (A point cannot admit charts of both types [Hat02, Theorem 2B.3].) We denote the set of all boundary points of X by ∂X .
- (5) An *atlas* for a manifold X consists of a family of charts such that the domains cover all of X . An atlas $\{\Phi_i: U_i \rightarrow V_i\}_{i \in I}$ is *smooth* if all transition maps $\Phi_i \circ \Phi_j^{-1}: \Phi_j(U_i \cap U_j) \rightarrow \Phi_i(U_i \cap U_j)$ are smooth. A *smooth manifold* is a manifold together with a smooth atlas. Usually one suppresses the choice of a smooth atlas from the notation.

To avoid misunderstandings we want to stress once again that what we call a “manifold” is often referred to as a “topological manifold”.

Definition 2.2. An *orientation* of an n -manifold M is a choice of generators $\alpha_x \in H_n(M, M \setminus \{x\}; \mathbb{Z})$ for each $x \in M \setminus \partial M$ such that for every $x \in M \setminus \partial M$ there exists an open neighbourhood $U \subseteq M \setminus \partial M$ of x and a class $\beta \in H_n(M, M \setminus U; \mathbb{Z})$ such that β projects to α_y for each $y \in U$.

Using the cross product one can show that the product of two oriented manifolds admits a natural orientation. Furthermore, the boundary of an oriented manifold also comes with a natural orientation. The proof of the latter statement is slightly delicate; we refer to [GH81, Chapter 28] or to [Fri23, Chapter 125.5] for details.

The following theorem [Hat02, Theorem 2B.3] is one of the foundational results on n -manifolds.

Theorem 2.3. (Invariance of Domain Theorem) *If $U \subseteq \mathbb{R}^n$ is an open subset and if $h: U \rightarrow \mathbb{R}^n$ is an injective map, then $h(U)$ is an open subset of \mathbb{R}^n .*

We conclude this section with the following lemma.

Lemma 2.4. *Let M be a connected manifold of dimension $n \geq 2$. Then for any two sets of pairwise disjoint points $\{x_1, \dots, x_m\}, \{y_1, \dots, y_m\} \in M \setminus \partial M$, there exists a homeomorphism $f: M \rightarrow M$ with $f(x_i) = y_i$ for $i = 1, \dots, m$.*

PROOF. Since M is connected we see that $M \setminus \partial M$ is path-connected. Thus there exist points $x_1 = t_0, t_1, \dots, t_{k+1} = y_1$ in $M \setminus \partial M$ such that there are charts $(U_i, \psi_i: U_i \rightarrow \text{Int } D^n)$ for $i = 0, \dots, k$ and both t_i, t_{i+1} are contained in U_i . It is elementary to show that given any two points $a, b \in \text{Int } D^n$ there exists a homeomorphism $f: \text{Int } D^n \rightarrow \text{Int } D^n$ with $f(a) = b$ and which is the identity outside of a compact subset. It is now clear that one can find a homeomorphism $f: M \rightarrow M$ with $f(x_1) = y_1$ such that f is the identity outside of a compact subset. We now consider the image of the remaining points in $M \setminus \{y_1\}$ and we restart the engine. \square

2.2. DEFINITION OF SUBMANIFOLDS

We move on to the definition of a submanifold. Again there are many different definitions in the literature, so let us define carefully what we mean by a submanifold.

Definition 2.5. Let M be an n -dimensional manifold. We say a subset $X \subseteq M$ is a k -dimensional submanifold if given any $x \in X$ one of the following holds:

(α) there exists a chart $\Phi: U \rightarrow V$ of type (i) for M and x such that

$$\Phi(U \cap X) \subseteq \{(0, \dots, 0, x_1, \dots, x_k) \mid x_1, \dots, x_k \in \mathbb{R}\},$$

(β) there exists a chart $\Phi: U \rightarrow V$ of type (ii) for M and x such that $\Phi(x)$ lies in E_{n-1} and

$$\Phi(U \cap X) \subseteq \{(0, \dots, 0, x_1, \dots, x_k) \in \mathbb{R}^n \mid x_k \geq 0\},$$

(γ) there exists a chart $\Phi: U \rightarrow V$ of type (i) for M and x such that $\Phi(x)$ lies in E_{n-1} and

$$\Phi(U \cap X) \subseteq \{(0, \dots, 0, x_1, \dots, x_k) \in \mathbb{R}^n \mid x_k \geq 0\}.$$

If for every $x \in X$ we can find charts as in (α) and (β), and $X \subseteq M$ is a closed subset, then we call X a *proper* submanifold.




FIGURE 2.1. Definition of submanifolds.

The following proposition is a straightforward consequence of the definitions. The proposition often makes it possible to reduce arguments about manifolds with boundary to the case of closed manifolds.

Proposition 2.6.

- (1) Let N be an n -manifold, possibly disconnected. Let A and B be collections of components of ∂N such that $A \cap B = \emptyset$. Let $f: A \rightarrow B$ be a homeomorphism. Then the quotient N/\sim under the relation $a \sim f(a)$ is an n -manifold with boundary $\partial(N/\sim) = \partial N \setminus (A \cup B)$. Moreover, the image of A in N/\sim is a submanifold.
- (2) Let M be an n -manifold. Its double $DM := M \cup M$, where the boundaries are identified via $\text{Id}_{\partial M}$, is an n -manifold with empty boundary. Moreover, $M \subseteq DM$ is a submanifold.

Definition 2.7. A map $f: X \rightarrow M$ from a k -manifold to an m -manifold M is called a (proper) *locally flat embedding* if f is a homeomorphism onto its image and if the image is a (proper) submanifold of M .

Remark 2.8.

- (1) Note that if M is a k -manifold and U is an open subset of \mathbb{R}^k , then it follows from the Invariance of Domain Theorem 2.3 that the image of any injective map $f: U \rightarrow M$ is an open subset of M . In particular $f(U)$ is a submanifold of M . Put differently, f is locally flat.
- (2) In point set topology, one often defines a *topological embedding* to be a map $f: X \rightarrow Y$ of topological spaces that is a homeomorphism to its image. The image of a topological embedding is not necessarily a submanifold and such an image is sometimes called *wild* due to the bizarre properties that such objects can exhibit. For example, the famous Alexander horned sphere [Ale24] is not a submanifold of S^3 under Definition 2.5, but it is the image of a wild topological embedding $S^2 \rightarrow S^3$.

- (3) In the literature a compact subset F of 4-manifold is often called a *locally flat surface* if F is homeomorphic to a compact 2-dimensional manifold with $\partial F = F \cap \partial M$ and if F has the following properties.
- (a) Given any $x \in F \setminus \partial F$ there exists a topological embedding $\varphi: D^2 \times D^2 \rightarrow M \setminus \partial M$ with $\varphi(D^2 \times D^2) \cap F = \varphi(D^2 \times \{0\})$ and with $x \in \varphi(D^2 \times \{0\})$.
 - (b) Given any $x \in \partial F$ there exists a topological embedding $\varphi: D_{\geq 0}^2 \times D^2 \rightarrow M$ such that $\varphi(D_{\geq 0}^2 \times D^2) \cap F = \varphi(D_{\geq 0}^2 \times \{0\})$, $\varphi(D_{\geq 0}^2 \times D^2) \cap \partial M = \varphi(\partial_{y=0} D_{\geq 0}^2 \times D^2)$, and with $x \in \varphi(D_{\geq 0}^2 \times \{0\})$. Here, we used the following abbreviations $D_{\geq 0}^2 = \{(x, y) \in D^2 \mid y \geq 0\}$ and $\partial_{y=0} D^2 = \{(x, 0) \in D^2\}$.
- It follows easily from the definitions that $F \subseteq M$ is a locally flat surface if and only if F is proper 2-dimensional submanifold of M .

The following proposition gives examples of embeddings $D^2 \rightarrow D^4$ that are not locally flat.

Proposition 2.9. *Given a knot $K \subseteq S^3$ the corresponding cone*

$$\text{Cone}(K) := \{r \cdot Q \mid Q \in K \text{ and } r \in [0, 1]\} \subseteq D^4.$$

is locally flat if and only if K is the unknot.

PROOF. Consider the specific unknot U that is the equator of the equator $U = S^1 \subseteq S^2 \subseteq S^3 = \partial D^4$. Taking the cone radially inwards to the origin of D^4 exhibits $\text{cone}(U)$ as a locally flatly (properly) embedded disc. Any other unknotted $K \subseteq S^3$ is related to U by a homeomorphism of S^3 . By the Alexander trick 4.4(1), this homeomorphism extends radially inwards to a homeomorphism of D^4 fixing the origin. Thus the cone on any other unknot K is locally flatly embedded, as we obtain a chart as in Definition 2.5(1) at the origin of D^4 .

Conversely, suppose $K \subseteq S^3$ is a knot such that $C := \text{Cone}(K)$ is locally flat. This implies that there is a chart $\Phi: U \rightarrow D^4$ where U is an open neighbourhood of the cone point 0, such that $\Phi(0) = 0$ and such that $\Phi(D^4 \cap C) = D^2 \times \{0\}$. We set $\Psi := \Phi^{-1}: D^4 \rightarrow U$. We introduce the following notation.

- (i) Given $J \subseteq [0, 1]$ we write $D_J := \{v \in D^4 \mid \|v\| \in J\}$.
- (ii) Given $J \subseteq [0, 1]$ we write $N_J := \Psi(D_J)$.

An elementary argument shows that there exist $s_1 < t_1 < s_2 < t_2 < s_3$ such that $D_{[0, s_1]} \subseteq N_{[0, t_1]} \subseteq D_{[0, s_2]} \subseteq N_{[0, t_2]} \subseteq D_{[0, s_3]}$. We make the following observations:

- (1) For any $J \subseteq [0, 1]$ we have homeomorphisms $D_J \setminus C \cong (S^3 \setminus K) \times J$ and $N_J \setminus C \xrightarrow{\Phi} D_J \setminus \text{Cone}(U) \cong (S^3 \setminus U) \times J$.
- (2) For any inclusion $J \subseteq J'$ of intervals the inclusion induced map $D_J \rightarrow D_{J'}$ is a homotopy equivalence.

We consider the following commutative diagram where all maps are induced by inclusions

$$\begin{array}{ccc}
 \pi_1(S^3 \setminus K) \cong \pi_1(D_{\{s_2\}} \setminus C) & \xrightarrow{\quad\quad\quad} & \pi_1(N_{[t_1, t_2]} \setminus C) \cong \mathbb{Z} \\
 & \searrow \cong & \swarrow \\
 & \pi_1(S^3 \setminus K) \cong \pi_1(D_{[s_1, s_3]} \setminus C). &
 \end{array}$$

Since the inclusion $D_{\{s_2\}} \setminus C \rightarrow D_{[s_1, s_3]} \setminus C$ is a homotopy equivalence we see that the left diagonal map is an isomorphism. Thus we see that we have an automorphism of $\pi_1(S^3 \setminus K)$ that factors through \mathbb{Z} . Since the abelianisation of $\pi_1(S^3 \setminus K)$ is isomorphic to \mathbb{Z} we see that $\pi_1(S^3 \setminus K) \cong \mathbb{Z}$. It follows from the Loop Theorem that K is in fact the unknot [Ro190, Theorem 4.B.1]. \square

We conclude this section with the following lemma, which provides us with examples of locally flat embeddings:

Lemma 2.10. *Let M be a connected manifold. Then any two points $x \neq y \in M \setminus \partial M$ are connected by a locally flat embedded arc.*

PROOF. By Lemma 2.4 we only have to deal with the case that x and y lie in a subspace that is homeomorphic to an open n -ball. But this case is trivial. \square

2.3. IMMERSIONS

We define immersions and generic immersions in the topological category cf. [KPRT22, Section 2]. In the smooth category immersions are required to have injective derivative at each point. In the topological category we cannot make such a definition, but instead define immersions as follows.

Definition 2.11. A continuous map $F: \Sigma^k \rightarrow M^n$ between manifolds of dimensions $k \leq n$ is an *immersion* if for each $p \in \Sigma$ there is a codimension zero submanifold $U \subseteq \Sigma$ containing p such that $F|_U: U \rightarrow M$ is a locally flat embedding.

Recall that a continuous map is said to be *proper* if the inverse image of every compact set in the codomain is compact. With this notion we can define proper immersions.

Definition 2.12. A proper continuous map $F: \Sigma^k \rightarrow M^n$ between manifolds of dimensions $k \leq n$ is a *proper immersion* if for each $p \in \Sigma$ there is a codimension zero submanifold $U \subseteq \Sigma$ containing p such that $U \subseteq \Sigma$ is an open subset, $F|_U: U \rightarrow M$ is a locally flat embedding, and every for $u \in U$ we can find a chart for $F(u)$ in M as in (α) and (β) of Definition 2.5.

We now consider surfaces in 4-manifolds, that is we restrict to $k = 2$ and $n = 4$. We take M to be a connected 4-manifold. The *singular set* of an immersion $F: \Sigma \rightarrow M$ is

$$\mathcal{S}(F) := \{m \in M \mid |F^{-1}(m)| \geq 2\}.$$

Definition 2.13. Let Σ be a surface, possibly noncompact. A continuous, proper map $F: \Sigma \rightarrow M$ is said to be a *generic immersion*, denoted $F: \Sigma \looparrowright M$, if it is a proper immersion and the singular set is a closed, discrete subset of M consisting only of transverse double points, each of whose preimages lies in the interior of Σ . The requirement that all singular points be transverse double points means that whenever $m \in \mathcal{S}(F)$, there are exactly two points $p_1, p_2 \in \Sigma$ with $F(p_1) = m = F(p_2)$, and there are disjoint charts φ_i around p_i , for $i = 1, 2$, where φ_1 and φ_2 are as in (2.1), with respect to the same chart Ψ around m and the standard inclusions

$$\begin{aligned} \iota_1: \mathbb{R}^2 &= \mathbb{R}^2 \times \{0\} \hookrightarrow \mathbb{R}^2 \times \mathbb{R}^2 = \mathbb{R}^4 \text{ and} \\ \iota_2: \mathbb{R}^2 &= \{0\} \times \mathbb{R}^2 \hookrightarrow \mathbb{R}^2 \times \mathbb{R}^2 = \mathbb{R}^4. \end{aligned}$$

$$(2.1) \quad \begin{array}{ccc} \mathbb{R}^2 & \xrightarrow{\iota_i} & \mathbb{R}^4 \\ \downarrow \varphi_i & & \downarrow \Psi \\ \Sigma & \xrightarrow{F} & M \end{array}$$

Typically one prefers to work with generic immersions of surfaces in 4-manifolds than arbitrary immersions. As the name suggests we can always arrange by a homotopy that maps of surfaces are generic immersions; see Theorem 10.6.

2.4. COLLAR NEIGHBOURHOODS

We discuss the existence of a collar neighbourhood of the boundary of a manifold. First we recall the definition of a neighbourhood.

Definition 2.14 (Neighbourhood). Let X be a space. A *neighbourhood* of a subset $A \subseteq X$ is a set $U \subseteq X$ for which there is an open set V satisfying $A \subseteq V \subseteq U$.

Next we give our definition of a collar neighbourhood.

Definition 2.15 (Collar neighbourhood). Let M be a manifold and let B be a union of components of ∂M . A *collar neighbourhood* is a map $\Phi: B \times [0, r] \rightarrow M$ for some $r > 0$ with the following three properties:

- (1) Φ is a locally flat embedding,
- (2) for all $x \in B$ we have $\Phi(x, 0) = x$,
- (3) we have $\Phi^{-1}(B \times [0, r]) \cap \partial M = B$.

Often, by a slight abuse of language, we identify $B \times [0, r]$ with its image $\Phi(B \times [0, r])$ and we refer to $B \times [0, r]$ also as a collar neighbourhood.

Note that a collar neighbourhood of a union B of components of ∂M is a neighbourhood of B . Now we can state the Collar Neighbourhood Theorem in the formulation of [**Arm70**, Theorem 1]. The existence of collars is originally due to Brown [**Bro62**], and there is another easier proof due to Connelly [**Con71**].

Theorem 2.16. (Collar Neighbourhood Theorem) *Let M be an n -manifold, let C be a compact submanifold of ∂M and let $f: C \times [0, 2] \rightarrow M$ be a map with $f(x, 0) = x$ for all $x \in C$. We assume that we are in one of the following two settings:*

- (1) C is closed as a manifold and f is a locally flat embedding;
- (2) $C \subseteq \partial M$ is a codimension zero submanifold and the restriction of f to $(\partial C \times [0, 2]) \cup (C \times \{2\})$ is a locally flat embedding.

Then there exists a collar neighbourhood $g: \partial M \times [0, 1] \rightarrow M$ with $g|_{C \times [0, 1]} = f$.

An *isotopy* of a space Y is a continuous one parameter family of maps $H = \{H_t\}_{t \in [0, 1]}: Y \times [0, 1] \rightarrow Y$ such that $H_t: Y \rightarrow Y$ is a homeomorphism for all $t \in [0, 1]$. To formulate a uniqueness result for collar neighbourhoods it helps to introduce the following definition.

Definition 2.17. Let $f, g: X \rightarrow Y$ be two maps between topological spaces and let Z be a subset of X . We say f and g are *ambiently isotopic rel. Z* if there exists an isotopy $H = \{H_t\}_{t \in [0, 1]}: Y \times [0, 1] \rightarrow Y$ such that $H_0 = \text{Id}$, $H_t|_Z = \text{Id}_Z$ for all t , and such that $H_1 \circ f = g$.

Theorem 2.18. *Let M be a manifold. Given two collar neighbourhoods $\Phi: \partial M \times [0, 2] \rightarrow M$ and $\Psi: \partial M \times [0, 2] \rightarrow M$, their restrictions $\Phi|_{\partial M \times [0, 1]}$ and $\Psi|_{\partial M \times [0, 1]}$ are ambiently isotopic rel. $\partial M \times \{0\}$.*

PROOF. The theorem is due to [Arm70, Theorem 2], although Armstrong comments that the proof was told to him by Lashof. See also [KS77, Essay I, Theorem A.2, p. 40]. \square

2.5. THE ISOTOPY EXTENSION THEOREM

In the smooth setting the Isotopy Extension Theorem gets used frequently and often subconsciously. In the non-smooth setting the formulation of the Isotopy Extension Theorem requires some care.

Definition 2.19. Let X be a k -dimensional manifold and let M be a compact m -dimensional manifold. Let $h: X \times [0, 1] \rightarrow M$ be a homotopy.

- (1) We say h is *locally flat* if for every $(x, t) \in X \times [0, 1]$ there exists a neighbourhood $[t_0, t_1]$ of t and level-preserving embeddings $\alpha: D^k \times [t_0, t_1] \rightarrow X \times [0, 1]$ and $\beta: D^k \times D^{m-k} \times [t_0, t_1] \rightarrow M \times [0, 1]$ to neighbourhoods of (x, t) and $(h_t(x), t)$ respectively, such that the following diagram commutes:

$$\begin{array}{ccc} D^k \times \{0\} \times [t_0, t_1] & \hookrightarrow & D^k \times D^{m-k} \times [t_0, t_1] \\ \downarrow \alpha & & \downarrow \beta \\ X \times [0, 1] & \xrightarrow{(x, t) \mapsto (h_t(x), t)} & M \times [0, 1]. \end{array}$$

- (2) We say h is *proper* if for every $t \in [0, 1]$ we have $h_t(X) \cap \partial M = h_t(\partial X)$.

This definition allows us to formulate the following useful theorem [EK71, Corollary 1.4] (see [Lee69, p. 530] for a related result.)

Theorem 2.20. (Isotopy Extension Theorem) *Let $h: X \times [0, 1] \rightarrow M$ be a locally flat proper isotopy of a compact manifold X into a manifold M . Then h can be covered by an ambient isotopy of M , i.e. there exists an isotopy $H: M \times [0, 1] \rightarrow M$ such that $H_0 = \text{Id}$ and $h_t = H_t \circ h_0$ for all $t \in [0, 1]$.*

Using the Isotopy Extension Theorem 2.20 we can now prove the following refinement of the Collar Neighbourhood Theorem 2.16.

Theorem 2.21. (Collar Neighbourhood Theorem for Proper Submanifolds) *Let M be a manifold and let $X \subseteq M$ be a proper submanifold. There exists a collar neighbourhood $\partial M \times [0, 1]$ such that $(\partial M \times [0, 1]) \cap X$ is a collar neighbourhood for $\partial X \subseteq X$.*

PROOF. By the earlier Collar Neighbourhood Theorem 2.16 we can pick a collar neighbourhood $\partial M \times [0, 2]$ for ∂M and we can also pick a collar neighbourhood $\partial X \times [0, 2]$ for ∂X . Given $t \in [0, 1]$ we consider the obvious homeomorphisms

$$f_t: M = (M \setminus (\partial M \times [0, 2])) \cup (\partial M \times [0, 2]) \rightarrow (M \setminus (\partial M \times [0, 2])) \cup (\partial M \times [t, 2])$$

and

$$g_t: X = (X \setminus (\partial X \times [0, 2])) \cup (\partial X \times [0, 2]) \rightarrow (X \setminus (\partial X \times [0, 2])) \cup (\partial X \times [t, 2]).$$

Next we consider the following proper locally flat isotopy:

$$h: X \times [0, 1] \rightarrow M$$

$$(x, t) \mapsto \begin{cases} (y, s) \in \partial M \times [0, t], & \text{if } x = (y, s) \text{ with } y \in \partial X, s \in [0, t], \\ f_t(g_t^{-1}(x)), & \text{otherwise.} \end{cases}$$

Note that the collar neighbourhood $\partial M \times [0, 1]$ is of the desired form for the proper submanifold $h_1(X)$. By the Isotopy Extension Theorem 2.20 we can extend h to a isotopy H of M . Thus $H_1^{-1}(\partial M \times [0, 1])$ is the desired collar neighbourhood for M . \square




FIGURE 2.2. Illustration of the proof of the Collar Neighbourhood Theorem 2.21.

CHAPTER 3

CW structures, triangulations, and handle structures on manifolds

In this chapter we will discuss the existence of various types of structures on compact manifolds. The results of this chapter are summarised in the following diagram. We use the following colour code.

- (1) A green arrow from A to B means that existence of structure A implies the existence of structure B .
- (2) A dashed blue arrow from A to B means that existence of structure A implies the existence of structure B , under the hypothesis that is written next to the blue arrow.




FIGURE 3.1. Existence of structures on a compact manifold.

In the following we will explain some of the structures mentioned in the diagram and we explain the various arrows.

Definition 3.1. An n -dimensional *PL manifold* is a simplicial complex X such that each $x \in X$ admits a neighbourhood that is PL homeomorphic to the standard PL n -ball.

Given a topological n -manifold M , a *PL structure* is a triangulation of M such that the resulting simplicial complex is an n -dimensional PL manifold.

Remark 3.2. Note that Dedecker [Ded62] and Zeeman [Zee62] showed that a PL-manifold can also be defined as a topological manifold together with a “piecewise linear” atlas. We refer to the the above references for precise definitions.

Definition 3.3. Let $CAT = TOP, PL,$ or $Diff$. Let M be a CAT -manifold and let W be a (possibly empty) union of components of ∂M . A *CAT-handle structure rel. W* is a CAT -isomorphism to a CAT -manifold that is obtained from $W \times [0, 1]$ by iteratively attaching handles of dimension $0, 1, 2, \dots$ along CAT -gluing maps.

With these definitions we can now formulate the theorems which in particular contain all the results mentioned in the diagram in Figure 3.1.

Theorem 3.4.

- (1) *Every compact smooth manifold admits a smooth handle structure rel. any union of boundary components.*
- (2) *Every smooth manifold admits a PL structure.*
- (3) *Every manifold that admits a smooth handle structure also admits a PL handle structure.*

PROOF.

- (1) In [Mil63, Section 3] and [Hir94, Section 6.4] it was shown that every compact smooth manifold admits a smooth handle decomposition.
- (2) In [Mun66, Theorem 10.6] and [Whi57, Chapter IV.12] it was shown that every smooth manifold admits a PL structure.
- (3) Every manifold that admits a smooth handle structure is by definition smooth. It follows from (2) and Theorem 3.5 (1) that every smooth manifold admits a PL handle structure. One might expect that one can turn every smooth handle structure into a PL handle structure by suitable choices of PL structures on handles and by applying isotopies to the attaching maps. But it is not clear to us whether or not this approach can be made to work. \square

Theorem 3.5.

- (1) *Every compact PL manifold admits a PL handle structure rel. any fixed union of boundary components.*
- (2) *Every compact PL manifold of dimension ≤ 7 admits a smooth structure, which is unique up to isotopy in dimensions ≤ 6 .*
- (3) *Every manifold that admits a PL handle structure also admits a topological handle structure.*

PROOF.

- (1) This statement is proved in [RS68, Proposition 6.8].
- (2) The 0-dimensional case is clear and the 1-dimensional case can be proved fairly easily by hand. The 2-dimensional case can also be done by hand: one picks obvious charts for the interiors of the 2-simplices and one can easily find charts that cover the interiors of the 1-simplices. For each 0-simplex one can pick a single chart by combining the various 2-simplices and adjusting the angle. The transition maps of such an atlas will all be smooth. Alternatively [Hat13, Theorem A] shows that every 2-dimensional topological manifold admits a smooth structure. Furthermore, [Hat13,

Theorem B] says that every homeomorphism between smooth surfaces is isotopic to a diffeomorphism.

We turn to manifolds of dimension 3, 4, 5, 6, and 7. By smoothing theory for PL manifolds, due to Munkres [Mun59, Mun60b] and Hirsch-Mazur [HM74, Part II, Theorem 4.2], isotopy classes of smooth structures on a compact PL manifold M are in one to one correspondence with homotopy classes of sections of a fibre bundle over M with fibre PL/O . It follows from work of Munkres [Mun60a], Cerf [Cer68], and Kervaire–Milnor [KM63] that PL/O is 6-connected. It then follows from obstruction theory that every compact PL manifold of dimension ≤ 7 admits a smooth structure, which is unique up to isotopy in dimensions ≤ 6 .

We remark that uniqueness of smooth structures in dimension 3 was proven earlier by Munkres [Mun59, p. 333], [Mun60a], [Mun60b, Theorems 6.2 and 6.3], and independently by Whitehead [Whi61, Corollary 1.18]. See also [Thu97, Chapter 3.10] for a more detailed discussion.

- (3) This statement follows immediately from the definitions. □

Theorem 3.6.

- (1) *Every manifold of dimension ≥ 5 admits a topological handle structure rel. any fixed union of boundary components.*
- (2) (a) *Every compact manifold of dimension 1, 2 or 3 admits a PL structure.*
 (b) *In dimension ≤ 4 a manifold that admits a simplicial structure also admits a PL structure.*
 (c) *Given any $n \geq 4$ there exists a closed n -manifold that does not admit a simplicial structure.*

PROOF.

- (1) For manifolds of dimension ≥ 6 this statement was proven by Kirby-Siebenmann [KS77, Essay III, Theorem 2.1, p. 104]. Quinn [Qui82, Theorem 2.3.1] extended this result to manifolds of dimension 5.
- (2) (a) Radó [Rad26] showed in 1926 that every compact 2-manifold has a simplicial structure. (Uniqueness was proved by Papakyriakopoulos [Pap46] in 1946.) Moise [Moi52, Moi77] proved the analogous result for 3-manifolds. We refer to [Bin59, Theorems 6 and 8], [Ham76, Theorem 2] and [Sha84] for alternative proofs of the 3-dimensional case.
 (b) By (a) we only need to consider the case of a 4-dimensional manifold M . We will defer the proof to Proposition 3.11 below. In fact, in Proposition 3.11 we will show the stronger fact that every triangulation of a manifold M of dimension n at most 4 is in fact a PL-structure. The inductive nature of the proof of Proposition 3.11 means that we need to consider all $n \leq 4$, even though only $n = 4$ is logically required at this point.
 (c) Casson [AM90, p. xvi] showed in the 1980s that there exist closed 4-manifolds that do not have a simplicial structure. It is now known that in every dimension

$n \geq 5$, there exists a closed n -manifold that does not admit a simplicial structure. This question was reduced to a problem about homology 3-spheres [Mat78, GS80], which was then solved by Manolescu [Man16]. Manolescu's examples are necessarily nonorientable.

We remark that orientable topological manifolds in dimension $n \geq 5$ that do not admit a PL structure were constructed much earlier by Siebenmann in [KS77, Annex C, Section 2]. However each of Siebenmann's manifolds admits a simplicial structure. \square

In order to prove Theorem 3.6 (2b), we will need some facts about polyhedral homology manifolds. We are grateful to Arunima Ray for assistance with this proof.

Definition 3.7. A locally finite n -dimensional simplicial complex P is an n -dimensional polyhedral homology manifold if for every $x \in P$ and for any subdivision of P with x as a vertex,

$$H_*(\text{Lk}(x); \mathbb{Z}) \cong H_*(S^{n-1}; \mathbb{Z}).$$

Remark 3.8.

- (1) It is immediate from the definition that if P is an n -dimensional polyhedral homology manifold, then so is any subdivision.
- (2) There is an extensions of the definition for polyhedral homology manifolds with nonempty boundary, but we will not give it here.

We need two facts about polyhedral homology manifolds. The first is that triangulated topological manifolds are examples of polyhedral homology manifolds.

Proposition 3.9 ([GS80, Proposition 1.2]). *Let M be a topological manifold with empty boundary, with a simplicial structure. Then M is a polyhedral homology manifold.*

The second fact we need, which will drive the induction in the proof of Proposition 3.11, is that the polyhedral homology manifold property descends to links of vertices.

Proposition 3.10. *Let P be an n -dimensional polyhedral homology manifold. Then for all vertices $v \in P$, the link $\text{Lk}(v)$ is a polyhedral homology manifold.*

PROOF. If $n = 1$, the link of a vertex consists of a collection of points, so is certainly a polyhedral homology manifold. So from now on we let $n \geq 2$. A lemma of Maunder [Mau80, Lemma 5.4.7, p. 188] implies the following. Let K be a simplicial complex, let $x \in K$ be a vertex, and let $L := \text{Lk}_K(x)$, and let $y \in L$ be a vertex. Let z be the midpoint of the line segment xy . Then

$$\tilde{H}_*(\text{Lk}_K(z); \mathbb{Z}) \cong \tilde{H}_{*-1}(\text{Lk}_L(y); \mathbb{Z})$$

where $\text{Lk}_K(z)$ is understood to be the link of z in the simplicial complex obtained from K by subdividing at z .

To apply this, let $K := P$ and let $x \in P$. Then since $z \in P$ is a vertex in some subdivision of P ,

$$\tilde{H}_*(S^{n-2}; \mathbb{Z}) \cong \tilde{H}_{*+1}(S^{n-1}; \mathbb{Z}) \cong \tilde{H}_{*+1}(\text{Lk}_K(z); \mathbb{Z}) \cong \tilde{H}_*(\text{Lk}_L(y); \mathbb{Z}).$$

Since this holds for every vertex y of L , it follows that $L = \text{Lk}_P(x)$ is a polyhedral homology manifold. \square

Now we prove the main result of our digression into polyhedral homology manifolds. In particular note that the case $n = 4$ (b) implies Theorem 3.6 (2b).

Proposition 3.11. *Let $n \leq 4$.*

- (a) *Let P be a finite $(n - 1)$ -dimensional polyhedral homology manifold with $H_*(P; \mathbb{Z}) \cong H_*(S^{n-1}; \mathbb{Z})$. If $n = 4$, suppose that $\pi_1(P) = \{1\}$. Then P is a PL $(n - 1)$ -manifold and is PL homeomorphic to S^{n-1} .*
- (b) *Every triangulation of a topological n -manifold M without boundary is a PL structure.*

PROOF. We work inductively on n , starting with $n = 1$. In this proof all homology groups are with \mathbb{Z} coefficients, which will be omitted for brevity.

We also make the following observation before we begin. Let X be a simplicial complex such that for every vertex $x \in X$, the link $\text{Lk}_X(x)$ is PL homeomorphic to the standard PL $(n - 1)$ -sphere. Then for each x , the star of x is the cone on the link of x , and so the star is PL homeomorphic to the standard n -ball. Hence X is a PL n -manifold without boundary.

The case $n = 1$ (a). This means that P is dimension 0, so is a finite collection of points. Thus P is certainly a PL manifold. Also $H_*(P) \cong H_*(S^0)$, so we deduce that P consists of two points, that is $P \cong S^0$.

The case $n = 1$ (b). By Proposition 3.9, M is a 1-dimensional polyhedral homology manifold, so the link $\text{Lk}(v)$ is dimension 0, with $H_*(\text{Lk}(v)) \cong H_*(S^0)$ for every vertex $v \in M$. Hence by the case $n = 1$ (a), $\text{Lk}(v) \cong S^0$, and so the triangulation of M is a PL structure.

The case $n = 2$ (a). Let P be a finite 1-dimensional polyhedral homology manifold with $H_*(P) \cong H_*(S^1)$. We have that P is a graph. By definition of a polyhedral homology manifold, links of vertices have $H_*(\text{Lk}(v)) \cong H_*(S^0)$, for every vertex $v \in P$. Hence vertices have valency two, so P is a PL 1-manifold. Since $H_*(P) \cong H_*(S^1)$, we must have that $P \cong S^1$, by the classification of compact PL 1-manifolds.

The case $n = 2$ (b). By Proposition 3.9, M is a 2-dimensional polyhedral homology manifold, so by Proposition 3.10 the link $\text{Lk}(v)$ is a polyhedral homology manifold of dimension 1 with $H_*(\text{Lk}(v)) \cong H_*(S^1)$ for every vertex $v \in M$. Hence by the case $n = 2$ (a), $\text{Lk}(v) \cong S^1$, and so the triangulation of M is a PL structure.

The case $n = 3$ (a). Now P is a 2-dimensional finite polyhedral homology manifold, with $H_*(P) \cong H_*(S^2)$. By Proposition 3.10, $\text{Lk}(v)$ is a polyhedral homology manifold and

$H_*(\text{Lk}(v)) \cong H_*(S^1)$ by the definition of a polyhedral homology manifold, for every vertex v of P . Then by the case $n = 1$ (a), $\text{Lk}(v)$ is a PL manifold and $\text{Lk}(v) \cong S^1$. Hence P is a PL manifold with $H_*(P) \cong H_*(S^2)$. Thus $P \cong S^2$ by the classification of compact PL 2-manifolds.

The case $n = 3$ (b). By Proposition 3.9, M is a 3-dimensional polyhedral homology manifold, so by Proposition 3.10 the link $\text{Lk}(v)$ is a polyhedral homology manifold of dimension 2 with $H_*(\text{Lk}(v)) \cong H_*(S^2)$ for every vertex $v \in M$. Hence by the case $n = 3$ (a), $\text{Lk}(v) \cong S^2$. Thus the triangulation of M is a PL structure, as desired.

The case $n = 4$ (a). Now P is a 3-dimensional finite polyhedral homology manifold, with $H_*(P) \cong H_*(S^3)$. In addition, since $n = 4$ we have by the hypotheses of the proposition that $\pi_1(P) = \{1\}$. By Proposition 3.10, for every vertex v of P , $\text{Lk}(v)$ is a polyhedral homology manifold with $H_*(\text{Lk}(v)) \cong H_*(S^2)$. Hence by the case $n = 3$ (a), $\text{Lk}(v)$ is a PL 2-manifold and $\text{Lk}(v) \cong S^2$. Hence P is a PL 3-manifold. Since P is simply-connected, compact, and has the homology of S^3 the Poincaré conjecture in dimension three [Per02, Per03b, Per03a], implies that $P \cong S^3$.

The case $n = 4$ (b). Let M be a 4-dimensional topological manifold with a simplicial structure. By Proposition 3.9, M is a 4-dimensional polyhedral homology manifold, so by Proposition 3.10 for every vertex $v \in M$, the link $\text{Lk}(v)$ is a polyhedral homology manifold of dimension 3 with $H_*(\text{Lk}(v)) \cong H_*(S^3)$. Moreover, for a triangulation of a topological manifold of dimension $n \geq 3$, a standard argument, similar to the proof of Proposition 2.9 (see [GS80, Theorem 1.5] for a proof), shows that $\pi_1(\text{Lk}(v)) = \{1\}$ for every vertex v . Hence we can apply the case $n = 3$ (a), to deduce that $\text{Lk}(v)$ is a PL manifold and $\text{Lk}(v) \cong S^3$. Thus the triangulation of M is a PL structure, as desired. \square

We continue by recalling the definition of a CW structure.

Definition 3.12. A *CW complex* is a topological space X together with a filtration

$$\emptyset = X_{-1} \subseteq X_0 \subseteq X_1 \subseteq X_2 \subseteq \cdots$$

such that $X = \varinjlim X_n$ and such that for each $n \geq 0$, the space X_n arises as a pushout

$$\begin{array}{ccc} \coprod_{j \in \mathcal{J}_n} S^{n-1} & \longrightarrow & X_{n-1} \\ \downarrow & & \downarrow \\ \coprod_{j \in \mathcal{J}_n} D^n & \longrightarrow & X_n \end{array}$$

where \mathcal{J}_n indexes the discs D^n . The interiors $\text{Int } D^n$ of the discs are called the *n -cells*. For $n \geq 0$, a CW complex X is said to be *n -dimensional* if $X_n \setminus X_{n-1} \neq \emptyset$ and $X_i = X_{i+1}$ for all $i \geq n$. We say a topological space X *admits a CW structure* if X admits such a filtration.

The above results can be used to prove the following theorem on the existence of CW structures.

Theorem 3.13.

- (1) *Every PL manifold (and thus every smooth manifold) admits a CW structure.*
- (2) *For $n \leq 3$ every compact n -manifold admits the structure of a finite n -dimensional CW complex.*
- (3) *Let $n \geq 5$ and let M be a compact n -manifold. Then M is homeomorphic to the mapping cylinder of some map $f: \partial M \rightarrow X$, where X is a finite CW complex.*
- (4) *For $n \geq 5$, every closed n -manifold admits the structure of a finite n -dimensional CW complex.*

PROOF.

- (1) Every simplicial complex is evidently a CW complex. The statement for PL manifold thus holds by definition. For smooth manifolds this statement now follows from Theorem 3.4 (2). Also note that it follows from [Mil63, Theorem 3.5] and the existence of a handle decomposition that every compact smooth n -dimensional manifold is homotopy equivalent to a compact n -dimensional CW complex. For most purposes it suffices to know that a compact smooth manifold has the homotopy type of a compact CW complex.
- (2) In Theorem 3.6 we saw that every compact n -manifold of dimension ≤ 3 admits a PL structure. It follows from (1) that every such manifold admits a CW structure.
- (3) Let M be a manifold with (possibly empty) boundary of dimension ≥ 5 . In Theorem 3.6 we saw that M admits a topological handle structure rel. $\partial M \times I$. Kirby-Siebenmann [KS77, Essay III, Theorem 2.2, p. 107] then says that M is homeomorphic to the mapping cylinder of some map $f: \partial M \rightarrow X$, where X is a finite CW complex. Thus (2) holds.
- (4) If M is a closed manifold of dimension ≥ 5 , then it follows from (3) that M admits the structure of a finite n -dimensional CW complex. \square

It is not clear to us whether Theorem 3.13 suffices to show that every compact high-dimensional manifold admits a CW structure. Put differently, to the best of our knowledge the following question is open for manifolds with nonempty boundary.

Question 3.14. *Let $n \geq 5$. Does every compact n -manifold have a CW structure?*

The following question also seems to still be open, even in the closed case.

Question 3.15. *Does every compact 4-manifold have a CW structure?*

In Theorem 3.13 we showed that many compact manifolds admit a CW structure, but we saw that there are still open cases. In many applications it suffices to know that a topological space is homotopy equivalent to a finite CW complex. This leads us to the following theorem.

Theorem 3.16.

- (1) Every connected closed nonempty n -manifold is homotopy equivalent to an n -dimensional finite CW complex.
- (2) Every connected compact nonempty n -manifold with nonempty boundary is homotopy equivalent to an $(n - 1)$ -dimensional finite CW complex.

In the proof of Theorem 3.16 we will use the following theorem proven by Wall [Wal66, Corollary 5.1].

Theorem 3.17. *Let X be a finite connected CW complex with fundamental group $\pi = \pi_1(X)$. Suppose that there is an integer $n \geq 3$ such that $H^i(X; \mathbb{Z}[\pi]) = 0$ for all $i > n$. (Here $H^i(X; \mathbb{Z}[\pi]) = 0$ denotes cohomology with twisted coefficients which we will introduce in Appendix A.1.) Then X is homotopy equivalent to an n -dimensional finite CW complex.*

PROOF OF THEOREM 3.16. We start out with the following claim.

Claim. Every compact manifold is homotopy equivalent to a finite CW complex.

We provide two different proofs of the claim.

- (a) Let M be a compact n -manifold. If $n \geq 6$, then it follows also from Theorem 3.13 (3) and (4) together with the Cellular Approximation Theorem [Bre97, Theorem IV.11.4] that M is homotopy equivalent to a finite CW complex. If $n < 5$, then we just replace M by $M \times D^6$ and apply the above argument. (see also Kirby-Siebenmann [KS69, Section 1 (III), p. 744].)
- (b) We now provide a second proof, which is of a very different flavour. Before we can provide this alternative argument we need to introduce one definition: a space X is called an *absolute neighbourhood retract (ANR)* if X is metrisable (recall that a space is metrisable if it admits the structure of a metric space inducing the given topology) and if whenever $X \subseteq Y$ is a closed subset of a metrisable space Y , then X is a neighbourhood retract of Y . That is, there is an open neighbourhood $U \subseteq Y$ containing X , with a map $r: U \rightarrow X$ such that the composition $X \rightarrow U \xrightarrow{r} X$ is equal to the identity on X .

Now we can return to the actual proof of the claim: It follows from the Dugundji Extension Theorem [Sak13, Theorem 6.1.1] and work of Hanner [Han51, Theorem 3.3] that every manifold (possibly noncompact) is an ANR, and West [Wes77, Corollary 5.3] showed that every compact ANR is homotopy equivalent to a finite CW complex.

With this claim it remains to prove the dimension statements of the theorem.

- (1) By Theorem 3.13 we only need to prove (1) in the case $n = 4$. But since the subsequent argument works for all $n \geq 4$ we also give it for all $n \geq 4$. We follow the argument provided in [Wes77, Corollary 5.4].

Let M be a compact connected nonempty n -manifold. By the claim, M is homotopy equivalent to a finite CW complex X .

First we consider the case that M is orientable. Since M is n -dimensional it follows from the Universal Poincaré Duality Theorem A.15 that for every $k > n$ and

every $\mathbb{Z}[\pi_1(X)]$ -module Λ we have

$$H^k(X; \Lambda) \cong H^k(M; \Lambda) \cong H_{n-k}(M, \partial M; \Lambda) = 0.$$

By Theorem 3.17, X is homotopy equivalent to an n -dimensional finite CW complex. Note that to apply Theorem 3.17 we have used that $n \geq 3$.

If M is nonorientable, then in the above one needs to apply Poincaré Duality for nonorientable manifolds. In the closed case, a proof can be found in [Sun17]. The case for manifolds with boundary can be proved by combining the ideas of [Sun17] and the Universal Poincaré Duality Theorem A.15.

- (2) Now we turn to the proof of (2). Let M be a compact connected n -manifold with nonempty boundary. We start with $n = 1, 2$, or 3 . We saw in Theorem 3.6 that in this dimension range every compact connected n -manifold admits a simplicial structure. It is well known that a compact connected n -manifold with nonempty boundary and a simplicial structure is homotopy equivalent to an $(n - 1)$ -dimensional simplicial complex: iteratively collapse top dimensional simplices starting with those that have a face on the boundary. In particular such a manifold is homotopy equivalent to an $(n - 1)$ -dimensional finite CW complex.

Now suppose that $n \geq 4$. By the claim M is homotopy equivalent to a finite CW complex X . Let $k \geq n$ and let Λ be a $\mathbb{Z}[\pi_1(X)]$ -module. By the Universal Poincaré Duality Theorem A.15 we have that

$$H^k(X; \Lambda) \cong H^k(M; \Lambda) \cong H_{n-k}(M, \partial M; \Lambda) = 0.$$

Here the last conclusion is obvious for $k > n$. For $k = n$ the conclusion follows from the fact that $\partial M \neq \emptyset$, that M is connected and the explicit calculation of 0-th twisted homology groups as given in [HS97, Chapter VI.3]. It follows from Theorem 3.17 that X is homotopy equivalent to an $(n - 1)$ -dimensional finite CW complex. Note here that to apply Theorem 3.17 we used $n \geq 4$. \square

Theorem 3.16 is strong enough to recover many familiar statements.

Corollary 3.18. *Let M be a compact, connected, nonempty manifold.*

- (1) *The group $\pi_1(M)$ is finitely presented.*
- (2) *All homology groups $H_k(M; \mathbb{Z})$ are finitely generated abelian groups, in particular it makes sense to define the Euler characteristic*

$$\chi(M) := \sum_n (-1)^n \cdot b_n(M).$$

- (3) *Let $p: \widetilde{M} \rightarrow M$ be a finite covering. Then*

$$\chi(\widetilde{M}) = [\widetilde{M} : M] \cdot \chi(M).$$

PROOF. The first two statements in the corollary are an immediate consequence of Theorem 3.16 and standard results on fundamental groups and homology groups of finite CW complexes. We turn to the final statement. Let X be a finite CW complex homotopy

equivalent to M . Use the fact that the Euler characteristic is multiplicative for finite covers of finite CW complexes and use that a k -fold cover \widetilde{M} of M induces a k -fold cover \widetilde{X} of X such that \widetilde{M} and \widetilde{X} are homotopy equivalent, to deduce the result.

Here is an alternative approach to proving the first two statements. Let M be a compact, connected manifold. Borsuk's Theorem that M is a Euclidean Neighbourhood Retract shows that M is a retract of a finite CW complex; see [Hat02, Appendix A, Corollary A.9], [Bre97, Appendix E], [Flo22], and [Fri23, Theorem 124.3 and Proposition 124.7]. This fact is nontrivial but it is much easier to prove than Theorem 3.16. Borsuk's Theorem implies immediately that the homology groups of M are finitely generated and that the fundamental group of M is finitely generated. In fact using a group theoretic lemma as in [Wal65, Lemma 1.3] or [FR01, Theorem 3.1], one actually obtains that $\pi_1(M)$ is finitely presented. But it is not clear how Borsuk's Theorem can be used to prove the third statement. \square

Remark 3.19. As pointed out above, every compact smooth manifold admits the structure of a finite CW complex. One can combine this fact with Theorem 9.9 below to obtain an alternative proof of Corollary 3.18 (1) and (2). More precisely: Theorem 9.9 says that for any compact 4-manifold M there is a closed orientable simply connected 4-manifold N such that the connected sum $M\#N$ admits a smooth structure. Using the well-known behaviour of the fundamental group and the homology groups under the connected sum operation one can now fairly easily provide an alternative proof of Corollary 3.18 (1) and (2).

CHAPTER 4

The Annulus Theorem and the Stable Homeomorphism Theorem

The Annulus Theorem and the Stable Homeomorphism Theorems are two (basically equivalent) fundamental results in the development of the theory of manifolds. For example, in high dimensions, the Stable Homeomorphism Theorem is an essential ingredient in the proof of the Product Structure Theorem 4.16 [KS77, Essay I, Theorem 5.1, p. 31], which itself underpins all the results of [KS77]. We state the Product Structure Theorem in Section 4.3. In dimension four, the Annulus Theorem is one of the many consequences of Quinn's controlled h -cobordism theorem [FQ90, Chapter 7]. In dimension 4 this theorem is used in the proofs of smoothing theorems (Chapter 9), existence and uniqueness of normal vector bundles (Chapter 5), and transversality (Chapter 10). We discuss these developments in the later chapters indicated. Later in this section (Section 4.2), we will discuss an application of the Annulus Theorem: showing that connected sum is a well-defined operation on connected, topological manifolds that are either oriented, or at least one of which is nonorientable. Here is the Annulus Theorem.

Theorem 4.1. (Annulus Theorem) *Let $n \in \mathbb{N}_0$ and let $f, g: D^n \rightarrow \mathbb{R}^n$ be two orientation-preserving locally flat embeddings. If $f(D^n) \subseteq \text{Int}(g(D^n))$, then $g(D^n) \setminus \text{Int}(f(D^n))$ is homeomorphic to $S^{n-1} \times [0, 1]$.*

For $n = 0, 1$ the Annulus Theorem is basically trivial. For $n = 2, 3$ the Annulus Theorem follows from the work of Radó [Rad26] and Moise [Moi52, Moi77] (see also [Edw84, p. 247]). The Annulus Theorem was proved for dimensions ≥ 5 by Kirby [Kir69], with a little help from Siebenmann, and in dimension 4 by Quinn [Qui82, p. 506], making use of the main results of [Fre82]; see also [Edw84, p. 247].

The known proofs of the Annulus Theorem 4.1 deduce it from the Stable Homeomorphism Theorem. In the next chapter we will state the Stable Homeomorphism Theorem 4.3 and we will explain the argument, provided in [BG64b], showing that the Annulus Theorem 4.1 can be deduced from the Stable Homeomorphism Theorem 4.3.

4.1. THE STABLE HOMEOMORPHISM THEOREM

We reduce the Annulus Theorem to the Stable Homeomorphisms Theorem stated in Theorem 4.3. This follows from work of Brown and Gluck [BG64b], but since it requires some work to find this deduction in [BG64b], we give the details here.

Definition 4.2. Let $n \in \mathbb{N}_0$. A homeomorphism $f: \mathbb{R}^n \rightarrow \mathbb{R}^n$ is said to be *stable* if there is a sequence of homeomorphisms $f_1, \dots, f_m: \mathbb{R}^n \rightarrow \mathbb{R}^n$ such that $f_m \circ \dots \circ f_1 = f$ and such

that for each i , the homeomorphism f_i is *somewhere the identity*, which means that there is an open nonempty set $U \subseteq \mathbb{R}^n$ such that $f_i|_U$ is the identity on U .

The key ingredient to the subsequent discussion is the following theorem.

Theorem 4.3. (Stable homeomorphism Theorem) *Let $n \in \mathbb{N}_0$. Every orientation preserving homeomorphism from \mathbb{R}^n to itself is stable.*

For $n \geq 5$ this was proven by Kirby [Kir69, p. 575] using the torus trick. Slightly more precisely, Kirby [Kir69] showed that the Stable Homeomorphism Theorem in dimensions at least five is a consequence of the surgery theoretic classification of PL homotopy tori, which was worked out around the same time by Wall [Wal69], [Wal99, Section 15A] and independently by Hsiang-Shaneson [HS69, p. 688], both proofs building on the work of Browder, Novikov, and Wall, which culminated in Wall's book [Wal99]. See also [HW69]. For $n = 4$ the Stable Homeomorphism Theorem was proven by Quinn [Qui82], see also [Edw84, p. 247].

Before we discuss consequences of the Stable Homeomorphism Theorem 4.3 we recall the two versions of the Alexander trick.

Lemma 4.4. (Alexander trick) *Let $n \in \mathbb{N}_0$.*

- (1) *Every homeomorphism of S^{n-1} can be extended radially to a homeomorphism of D^n that sends 0 to 0.*
- (2) *Let f and g be two homeomorphisms of D^n . If the restrictions of f and g to S^{n-1} are isotopic, then f and g are isotopic homeomorphisms of D^n .*
- (3) *The topological group $\text{Homeo}_\partial(D^n)$ of homeomorphisms of D^n fixing the boundary pointwise is contractible.*

PROOF. The extension in the first statement can be obtained by coning: $f(t \cdot x) = t \cdot f(x)$.

The proof of the second is an amusing exercise; see e.g. [Han89, Lemma 5.6] for a proof. The idea is also similar to the proof of the third statement, which we give now. To show that $\text{Homeo}_\partial(D^n)$ is contractible it suffices to give a homotopy

$$F: \text{Homeo}_\partial(D^n) \times [0, 1] \rightarrow \text{Homeo}_\partial(D^n)$$

with $F(f, 0) = \text{Id}_D^n$ and $F(f, 1) = f$, for all $f \in \text{Homeo}_\partial(D^n)$. To do this, we define

$$F(f, t)(x) = \begin{cases} t \cdot f\left(\frac{1}{t} \cdot x\right) & \|x\| \leq t \\ x & t \leq \|x\| \leq 1. \end{cases}$$

We omit the proof that this is well-defined and continuous. □

We can now prove the following almost immediate consequence of the Stable Homeomorphism Theorem 4.3.

Corollary 4.5. *Let $n \in \mathbb{N}_0$. Every orientation preserving self-homeomorphism of S^n is isotopic to the identity.*

PROOF. We identify S^n with $\mathbb{R}^n \cup \{\infty\}$. Let h be a self-homeomorphism of $S^n = \mathbb{R}^n \cup \{\infty\}$. By the Stable Homeomorphism Theorem 4.3 we know that h is stable. Thus we only have to consider the case that h fixes an open subset of $\mathbb{R}^n \cup \{\infty\} = S^n$, since a composition of homeomorphisms isotopic to the identity is isotopic to the identity. After an isotopy (using Theorem 2.20) we can assume that h fixes an open neighbourhood of ∞ , so in particular there exists $C > 0$ such that h is the identity on $\{x \in \mathbb{R}^n \mid \|x\| \geq C\}$. It then follows from Lemma 4.4 (2) that h is isotopic to the identity. \square

Now we begin showing how to deduce the annulus theorem from the stable homeomorphism theorem.

Definition 4.6. Let $n \in \mathbb{N}_0$. We say that two elements $f_0, f_1 \in \text{Emb}(D^n, \mathbb{R}^n)$ are *intertwined* if there exists an $h \in \text{Homeo}(\mathbb{R}^n, \mathbb{R}^n)$ with $h \circ f_0 = f_1$.

We will need the following straightforward technical lemma. See Definition 2.7 for the definition of a locally flat embedding.

Lemma 4.7. Let $n \in \mathbb{N}_0$. Let M be an n -dimensional manifold and let $f: D^n \rightarrow M$ be a locally flat embedding into $\text{Int } M = M \setminus \partial M$. Then there exists a locally flat embedding $F: \mathbb{R}^n \rightarrow M$ such that the restriction of F to D^n equals f .

PROOF. Let $f: D^n \rightarrow M$ be a locally flat embedding. By definition $f(D^n)$ is a submanifold of M . It is straightforward to see that $W := M \setminus f(\text{Int } D^n)$ is also a submanifold of M . By the Collar Neighbourhood Theorem 2.16 there exists a collar $f(S^{n-1}) \times [0, 1]$. The map

$$F: \mathbb{R}^n \rightarrow M$$

$$x \mapsto \begin{cases} f(x), & \text{if } x \in D^n, \\ (f(y), \frac{2}{\pi} \arctan(t-1)) & \text{if } x = t \cdot y \text{ with } t \in [1, \infty) \text{ and } y \in S^{n-1}, \end{cases}$$

is easily seen to be a locally flat embedding. \square

Denote the set of locally flat embeddings of D^n into \mathbb{R}^n by $\text{Emb}(D^n, \mathbb{R}^n)$.

Lemma 4.8. Let $n \in \mathbb{N}_0$. Any two elements $f_0, f_1 \in \text{Emb}(D^n, \mathbb{R}^n)$ are intertwined.

PROOF. It suffices to show that any $f \in \text{Emb}(D^n, \mathbb{R}^n)$ is intertwined with the standard embedding $D^n \subseteq \mathbb{R}^n$. So let $f \in \text{Emb}(D^n, \mathbb{R}^n)$. Apply Lemma 4.7 to extend f to a locally flat embedding $F: D_{\frac{3}{2}}^n \rightarrow \mathbb{R}^n$. Note that F restricts to a locally flat embedding of $S^{n-1} \times [\frac{1}{2}, \frac{3}{2}]$ into $S^n = \mathbb{R}^n \cup \{\infty\}$. Let \tilde{D}^n be another copy of D^n . By the generalised Schoenflies Theorem [Bro60, Theorem 5] there exists a homeomorphism $g: \tilde{D}^n \rightarrow S^n \setminus f(\text{Int } D^n)$. Using that the homeomorphisms of \tilde{D}^n act transitively on the interior of \tilde{D}^n , arrange that $g(0) = \infty$.

Note that $g^{-1} \circ f: S^{n-1} \rightarrow S^{n-1}$ is a homeomorphism. By Lemma 4.4 (1) this homeomorphism extends to a homeomorphism ϕ of D^n . Replace g by $g \circ \phi$ if necessary to obtain

that $f = g: S^{n-1} \rightarrow f(S^{n-1})$. Identify $S^n = \mathbb{R}^n \cup \{\infty\} = D^n \cup \tilde{D}^n$ in such a way that $0 \in \tilde{D}^n$ corresponds precisely to ∞ . Consider the map

$$F: S^n = D^n \cup \tilde{D}^n \rightarrow S^n$$

$$x \mapsto \begin{cases} f(x) & x \in D^n \\ g(x) & x \in \tilde{D}^n. \end{cases}$$

The maps f and g agree on the overlap, so the map is well-defined and is a homeomorphism. Note that F restricts to a homeomorphism of \mathbb{R}^n which has the property that the restriction to D^n equals f . This shows that $F \circ \text{Id} = f$, so f and the standard embedding are intertwined. \square

We continue with the following definition from [BG64b, p. 19].

Definition 4.9. Let $n \in \mathbb{N}_0$. Let $f_0, f_1 \in \text{Emb}(D^n, \mathbb{R}^n)$.

- (1) We say f_0 and f_1 are *strictly annularly equivalent* if $f_0(D^n) \subseteq \text{Int } f_1(D^n)$ and if there exists a map $F: S^{n-1} \times I \rightarrow \mathbb{R}^n$ that is a homeomorphism onto its image such that $F(x, 0) = f_0(x)$ and $F(x, 1) = f_1(x)$ for all $x \in S^{n-1}$.
- (2) We say f_0 and f_1 are *annularly equivalent* if there exists a sequence $f_0 = g_0, g_1, \dots, g_k = f_1$ of elements of $\text{Emb}(D^n, \mathbb{R}^n)$ such that any two successive g_i are strictly annularly equivalent.

Theorem 4.10. Let $n \in \mathbb{N}_0$. Let $f_0, f_1 \in \text{Emb}(D^n, \mathbb{R}^n)$ with $f_0(D^n) \subseteq \text{Int } f_1(D^n)$. If f_0 and f_1 are orientation preserving, then they are strictly annularly equivalent if and only if they are intertwined.

PROOF. If two such elements are strictly annularly equivalent, then they are intertwined by [BG64b, Theorem 5.2].

Now suppose that f_0 and f_1 are intertwined, that is there exists an $h \in \text{Homeo}(\mathbb{R}^n, \mathbb{R}^n)$ with $h \circ f_0 = f_1$. By the Stable Homeomorphism Theorem 4.3 we know that h is stable. Thus we know from [BG64b, Theorem 5.4] that the embeddings are annularly equivalent, i.e. there exist $h_0, \dots, h_k \in \text{Emb}(D^n, \mathbb{R}^n)$ such that $h_0 = f_0, h_k = f_1$ and for each i the maps h_i and h_{i+1} are strictly annularly equivalent. Since $f_0(D^n) \subseteq \text{Int } f_1(D^n)$, the embeddings of the boundary spheres $f_0(\partial D^n)$ and $f_1(\partial D^n)$ are disjoint. Therefore it follows from [BG64a, Theorem 3.5] that f_0 and f_1 are not only annularly equivalent, but are moreover strictly annularly equivalent. \square

Now we can easily prove the Annulus Theorem 4.1.

PROOF OF THE ANNULUS THEOREM 4.1. Let $f_0, f_1: D^n \rightarrow \mathbb{R}^n$ be two orientation-preserving locally flat embeddings with $f_0(D^n) \subseteq \text{Int}(f_1(D^n))$. By Lemma 4.8 and Theorem 4.10 the two maps f_0 and f_1 are strictly annularly equivalent. But this implies that $f_1(D^n) \setminus \text{Int}(f_0(D^n))$ is homeomorphic to $S^{n-1} \times [0, 1]$. \square

4.2. THE CONNECTED SUM OPERATION

Definition 4.11. Let M and N be connected nonempty oriented n -manifolds. Pick an orientation preserving locally flat embedding $\Phi_M: D^n \rightarrow M \setminus \partial M$ of an n -ball into M and an orientation reversing locally flat embedding $\Phi_N: D^n \rightarrow N \setminus \partial N$ of an n -ball into N . Define the *connected sum* $M\#N$ of M and N by

$$M\#N := (M \setminus \Phi_M(\text{Int}(D^n))) \cup_{\Phi_M(S^{n-1})=\Phi_N(S^{n-1})} (N \setminus \Phi_N(\text{Int}(D^n)))$$

where we glue the left hand side to the right hand side via the map

$$\Phi_N \circ \Phi_M^{-1}: \Phi_M(S^{n-1}) \xrightarrow{\cong} \Phi_N(S^{n-1}).$$

It follows from the Collar Neighbourhood Theorem 2.16 that the topological space $M\#N$ inherits the structure of an n -manifold; see [Lee11, Proposition 6.6] for details. Furthermore $M\#N$ can be oriented in such a way that $M \setminus \Phi_M(D^n)$ and $N \setminus \Phi_N(D^n)$ are oriented submanifolds.

Theorem 4.12. *Let $n \in \mathbb{N}_0$. The connected sum $M\#N$ of two connected oriented n -manifolds M and N is independent of the choice of embeddings of the n -balls.*

Remark 4.13.

- (1) In Proposition 8.3 we will see that the manifolds $\mathbb{C}P^2\#\overline{\mathbb{C}P^2}$ and $\mathbb{C}P^2\#\mathbb{C}P^2$ have non-isometric intersection forms, so they are not homeomorphic. Thus connected sum is not well-defined on orientable 4-manifolds, rather it depends on the choice of orientation.
- (2) If at least one of the two manifolds involved is nonorientable, then the connected sum is well-defined. This follows from the fact that one can show, say using the orientation cover of a nonorientable manifold, that in a nonorientable connected n -manifold M any two locally flat embeddings of $D^n \rightarrow M \setminus \partial M$ are ambiently isotopic. In the smooth category this argument is worked out in detail in [Fri23, Chapter 47.4]. Using the Annulus Theorem 4.1 one can translate the smooth argument to a topological argument.
- (3) As discussed in [BCF⁺21], in contrast to the case of orientable 3-dimensional manifolds, orientable 4-dimensional topological manifolds do not admit a unique decomposition as a connected sum of irreducible 4-manifolds. For example $\mathbb{C}P^2\#S^2 \times S^2$ and $\mathbb{C}P^2\#\mathbb{C}P^2\#\overline{\mathbb{C}P^2}$ are diffeomorphic.

The proof of Theorem 4.12 relies on the following two lemmas. The elementary proof of the first lemma is left to the reader.

Lemma 4.14. *Let $D_r^n(x)$ and $D_s^n(y)$ be two Euclidean balls in \mathbb{R}^n . There exists an orientation-preserving homeomorphism $f: \mathbb{R}^n \rightarrow \mathbb{R}^n$ with $f(D_r^n(x)) = D_s^n(y)$ such that f is the identity outside of some compact set.*

The next lemma is a consequence of the Annulus Theorem 4.1.

Lemma 4.15. *Let $\varphi, \psi: D^n \rightarrow \mathbb{R}^n$ be two orientation-preserving locally flat embeddings. If $\varphi(D^n) \subseteq \text{Int}(\psi(D^n))$, then there exists an orientation-preserving homeomorphism f of \mathbb{R}^n with $f(\varphi(D^n)) = \psi(D^n)$ such that f is the identity outside of some compact set.*

PROOF. By the Annulus Theorem 4.1 and the Collar Neighbourhood Theorem 2.16 we can find a locally flat embedding $\theta: S^{n-1} \times [-1, 2]$ such that $\theta(S^{n-1} \times [-1, 0]) \subseteq \varphi(D^n)$ is an (interior) collar for $\partial\varphi(D^n)$, such that $\theta(S^{n-1} \times [0, 1]) = \psi(D^n) \setminus \varphi(\text{Int } D^n)$ and such that $\theta(S^{n-1} \times [1, 2]) \subseteq \mathbb{R}^n \setminus \psi(\text{Int } D^n)$ is an (internal) collar for $\partial(\mathbb{R}^n \setminus \psi(\text{Int } D^n))$. It is now obvious that we can find a homeomorphism f with $f(\varphi(D^n)) = \psi(D^n)$ which is the identity outside of $\theta(S^{n-1} \times [-1, 2])$. \square

The subsequent proof is partly based on the sketch given in [Rol90, p. 42].

PROOF OF THEOREM 4.12. We have to show that the connected sum is independent of the choice of $\Phi_M: D^n \rightarrow M$ and $\Phi_N: D^n \rightarrow N$. In the following we show that the oriented homeomorphism type of the connected sum is independent of the choice of Φ_M . Basically the same argument then shows that the oriented homeomorphism type of the connected sum is independent of the choice of Ψ_N . Putting these two orientation-preserving homeomorphisms together gives independence of all choices.

After this preamble we now show that the oriented homeomorphism type of the connected sum is independent of the choice of Φ_M . So suppose we are given two orientation-preserving embeddings $\Phi_1: D^n \rightarrow M$ and $\Phi_2: D^n \rightarrow M$ and suppose we are given an orientation-reversing embedding $\Psi: D^n \rightarrow N$. For $i = 1, 2$ we introduce the following notation.

- (1) Write $D_i := \Phi_i(D^n)$.
- (2) Let $X_i := M \setminus \Phi_i(\text{Int } D^n)$ and let $Y := N \setminus \Psi(\text{Int } D^n)$,
- (3) Denote the restriction of Φ_i to S^{n-1} by φ_i and denote the restriction of Ψ to S^{n-1} by ψ .

Figure 4.1 hopefully makes it easier for the reader to internalise the notation. We have to show that there exists a homeomorphism

$$(X_1 \cup Y)/\varphi_1(x) \sim \psi(x) \rightarrow (X_2 \cup Y)/\varphi_2(x) \sim \psi(x)$$

where the gluing on both sides is given by taking $x \in S^{n-1}$.

Claim. There exists an orientation-preserving homeomorphism h of M so that $h(D_1) = D_2$.

To prove the claim, first note that it follows from Lemmas 4.7 and 4.14, together with our hypothesis that M is path connected, that there exists an orientation-preserving homeomorphism μ of M such that $\mu(D_1) \subseteq \text{Int } D_2$. Then apply Lemmas 4.7 and 4.15 to find an orientation-preserving homeomorphism ν of M such that $\nu(\mu(D_1)) = D_2$. This concludes the proof of the claim.

After replacing φ_1 by $h \circ \varphi_1$ we can assume that $\varphi_2^{-1} \circ \varphi_1$ is an orientation-preserving homeomorphism of S^{n-1} . By Corollary 4.5 we know that there exists an isotopy $H: S^{n-1} \times [0, 1] \rightarrow S^{n-1}$ from $\varphi_2^{-1} \circ \varphi_1$ to the identity.




FIGURE 4.1. Illustration for the proof of Theorem 4.12.

We write $C := \Psi(S^{n-1})$. By the Collar Neighbourhood Theorem 2.16 we can pick an (internal) collar $C \times [0, 1] \subseteq Y$ for ∂Y . It is straightforward to verify that

$$(X_1 \cup Y)/\varphi_1(x) \sim \psi(x) \rightarrow (X_2 \cup Y)/\varphi_2(x) \sim \psi(x)$$

$$p \mapsto \begin{cases} h(p), & \text{if } p \in X_1, \\ \psi(H(\psi^{-1}(q), t)) & \text{if } p = (q, t) \in C \times [0, 1] \\ p, & \text{if } p \in Y \setminus (C \times [0, 1]) \end{cases}$$

is a well-defined map and is an orientation-preserving homeomorphism. This shows that the connected sums defined using Φ_1 and Φ_2 give rise to manifolds of the same oriented homeomorphism type. \square

4.3. THE PRODUCT STRUCTURE THEOREM

The Product Structure Theorem [KS77, Essay I, Theorem 5.1, p. 31], is a key result for the development of topological manifold theory in high dimensions. It is a consequence of the Stable Homeomorphism Theorem 4.3, together with a more sophisticated torus trick. The Product Structure Theorem is used in [KS77] to deduce the existence of handle structures for manifolds of dimension $n \geq 6$, transversality and smoothing theory for $n \geq 5$, and the existence of a canonical simple homotopy type for all n . We will give some examples of the use of the Product Structure Theorem, for instance in Section 14.1 on the simple homotopy type of topological manifolds. Even though the Product Structure Theorem a priori only concerns high dimensional manifolds, it still appears in the development of the theory of 4-manifolds.

The Product Structure Theorem will be stated for upgrading either a smooth or PL structure on $M \times \mathbb{R}$ to one on M .

A *concordance* of (smooth, PL) structures Σ, Σ' on a manifold N is a (smooth, PL) structure Ω on $N \times I$ that restricts to Σ on $N \times \{0\}$ and restricts to Σ' on $N \times \{1\}$.

Theorem 4.16. (Product Structure Theorem) *Let M be a manifold of dimension $n \geq 5$. Let Σ be a (smooth, PL) structure on $M \times \mathbb{R}^s$, with $s \geq 1$. Let U be an open subset of M with a (smooth, PL) structure ρ on U such that $\rho \times \mathbb{R}^s = \Sigma|_{U \times \mathbb{R}^s}$. If $n = 5$ then suppose that $\partial M \subseteq U$.*

Then there is a (smooth, PL) structure σ on M extending ρ , together with a concordance of (smooth, PL) structures from Σ to $\sigma \times \mathbb{R}^s$, that is a product concordance in some neighbourhood of $U \times \mathbb{R}^s$ and that is a product near $M \times \mathbb{R}^s \times \{i\}$ for $i = 0, 1$.

Remark 4.17. The statement of the Product Structure Theorem was modelled on the Cairns-Hirsch Theorem [KS77, Essay I, Theorem 5.3, p. 37], which was proven in the early 1960s, and provided the analogous upgrade from PL structures to smooth structures. See [HM74] for a comprehensive treatment of smoothing theory for PL manifolds. The Cairns-Hirsch Theorem tells us that if M already has a PL structure ϖ , such that $\varpi \times \mathbb{R}^s$ is *Whitehead compatible* (see the discussion below [KS77, Essay I, Theorem 5.3, p. 37] for details) with a smooth structure Σ on $M \times \mathbb{R}^s$, then the smooth structure σ on M produced by Theorem 4.16 is Whitehead compatible with ϖ .

In Section 14.1 on the simple homotopy type of a manifold we make use of the following stronger local version [KS77, Essay I, Theorem 5.2, p. 36].

Theorem 4.18. (Local Product Structure Theorem) *Let M be a manifold of dimension $n \geq 5$.*

- (i) *Let W be an open neighbourhood of $M \times \{0\}$ in $M \times \mathbb{R}^s$, for some $s \geq 1$.*
- (ii) *Let Σ be a (smooth, PL) structure on W .*
- (iii) *Let $C \subseteq M \times \{0\}$ be a closed subset such that there is a neighbourhood $N(C)$ of $C \subseteq W$ on which the (smooth, PL) structure Σ is a product $\Sigma|_{N(C)} = \sigma \times \mathbb{R}^s$ for some (smooth, PL) structure σ on $N(C)$. If $n = 5$ then suppose that $\partial M \subseteq C$.*
- (iv) *Let $D \subseteq M \times \{0\}$ be another closed subset.*
- (v) *Let $V \subseteq W$ be an open neighbourhood of $D \setminus C$ in $M \times \mathbb{R}^s$.*

Then we have the following.

- (1) *A (smooth, PL) structure Σ' on W that equals Σ on $(W \setminus V) \cup ((C \times \mathbb{R}^s) \cap W)$ and is a product (smooth, PL) structure $\rho \times \mathbb{R}^s$ on $(N(D) \times \mathbb{R}^s) \cap W$ for some neighbourhood $N(D)$ of D and for some (smooth, PL) structure ρ on $N(D)$.*
- (2) *A concordance of (smooth, PL) structures from Σ to Σ' , that is a product concordance on some neighbourhood of $(W \setminus V) \cup ((C \times \mathbb{R}^s) \cap W)$ and that is a product near $W \times \{i\}$ for $i = 0, 1$.*

Note that the Concordance implies Isotopy Theorem [KS77, Essay I, Theorem 4.1, p. 25] means that the concordances in Theorems 4.16 and 4.18 can be upgraded to isotopies of (smooth, PL) structures under the same hypotheses on dimensions, that is if $n \geq 6$ or if $n = 5$ and the structures already agree on ∂M .

We start with a structure on W that is a product structure over C . We obtain a concordance of structures to a product structure on D , supported in V , and that is a product concordance over C . We therefore have a product structure in some neighbourhood of $C \cup D$.

CHAPTER 5

Tubular neighbourhoods

Every smooth submanifold of a smooth manifold admits a normal vector bundle and, by the smooth Tubular Neighbourhood Theorem, also admits a tubular neighbourhood [Hir94, Sections 5 & 6], [Wal16, Chapter 2.5]. However, in the topological category submanifolds may not admit normal vector bundles, a general problem we discuss further below and in Chapter 6 once we have developed the necessary language. Curiously, in the special case of 4-manifolds these general problems do not exist, and familiar smooth results hold true using an appropriate notion of normal vector bundles (Definition 5.12).

5.1. TUBULAR NEIGHBOURHOODS: EXISTENCE AND UNIQUENESS

In the literature one can find many different definitions of tubular neighbourhoods for smooth submanifolds. We will give a definition for manifolds that is modelled on the definition provided by Wall [Wal16] for smooth manifolds. To do so we first need one extra definition.

Definition 5.1. Let M be an n -dimensional manifold. We say a subset $W \subseteq M$ is a *k -dimensional submanifold with corners* if given any $p \in W$ there exists a chart of the type (1), (2) or (3) as in Definition 2.5 above, or if

(4) there exists a chart $\Phi: U \rightarrow V$ of type (ii) for M such that

$$\Phi(U \cap W) \subseteq \{(0, \dots, 0, x_1, \dots, x_k) \mid x_i \in \mathbb{R} \text{ with } x_{k-1} \geq 0 \text{ and } x_k \geq 0\}$$

and with $\Phi(p) \in \{(0, \dots, 0, x_1, \dots, x_{k-2}, 0, 0) \mid x_1, \dots, x_{k-2} \in \mathbb{R}\}$.

If W is an n -dimensional submanifold with corners we write

$$\partial_0 W := W \cap \overline{M \setminus W}, \quad \partial_1 W := W \cap \partial M,$$


and we note that

$$\text{Int } W = W \setminus \partial_0 W.$$

Remark 5.2. The complement of the interior of a codimension 0 submanifold with corners is a submanifold with corners, with the obvious corners.

Definition 5.3. Let M be an n -manifold and let X be a compact, proper, k -dimensional submanifold. A *tubular neighbourhood* for X is a pair $(N, p: N \rightarrow X)$ with the following properties:

(1) N is a neighbourhood of X ;

FIGURE 5.1. Definition of ∂_0 and $\partial_1 W$ of a submanifold.

- (2) N is a codimension zero submanifold with corners of M ;
- (3) the map $p: N \rightarrow X$ is a linear D^{n-k} -bundle such that $p(x) = x$ for all $x \in X$;
- (4) $\partial_1 N = p^{-1}(\partial X)$.

Here linear means that there exists an atlas of local trivialisations such that the transition maps take values in $O(n-k)$ instead of $\text{Homeo}(D^{n-k})$.

Remark 5.4. In the topological category, tubular neighbourhoods do not always exist. Indeed it is shown in [Hir68, Theorem 4] that there exists a 4-dimensional submanifold of S^7 that does not admit a tubular neighbourhood.

Fortunately, for submanifolds of 4-manifolds, tubular neighbourhoods exist and they are unique in the appropriate sense.

Theorem 5.5. (Tubular Neighbourhood Theorem) *Every compact proper submanifold X of a 4-manifold M admits a tubular neighbourhood.*

Theorem 5.6. (Uniqueness of tubular neighbourhoods) *Let M be a 4-manifold and let X be a compact proper k -dimensional submanifold. Furthermore let $p_i: N_i \rightarrow X$, $i = 1, 2$ be two tubular neighbourhoods of X , with inclusion maps $\iota_i: N_i \rightarrow M$. Then there exists an isomorphism $\Psi: N_1 \rightarrow N_2$ of linear disc bundles such that $\iota_2 \circ \Psi: N_1 \rightarrow M$ and $\iota_1: N_2 \rightarrow M$ are ambiently isotopic rel. X .*

The proofs of the above two theorems rely on the existence and uniqueness results for normal vector bundles in [FQ90, Section 9], which we discuss further in Section 5.2. Thus we postpone the proofs of the Theorems 5.5 and 5.6 to Section 5.3. Right now, let us first observe some nice consequences of the existence and uniqueness of tubular neighbourhoods.

Remark 5.7. Let X be a compact proper submanifold of a 4-manifold M . By Theorem 5.5 we can pick a tubular neighbourhood $p: N \rightarrow X$. We refer to $E_X := M \setminus \text{Int } N$ as the *exterior* of X . Note $E_X \subseteq M$ is a submanifold with corners; cf. Remark 5.2. By Theorem 5.6 the homeomorphism type of the exterior is well-defined.

Lemma 5.8. *Let X be a compact proper submanifold of a 4-manifold M . The exterior E_X of X is a deformation retract of the complement $M \setminus X$.*

PROOF. Let $p: N \rightarrow X$ be a tubular neighbourhood for X . Using the fact that p is a linear bundle, introduce compatible radial coordinates in the fibres and isotope $N \setminus X$

radially outwards. This implies that $\partial_0 N$ is a deformation retract of $N \setminus X$. But this also implies that the exterior $E_X = M \setminus N$ is a deformation retract of $M \setminus X$. \square

Corollary 5.9. *Let X be a proper submanifold of a compact 4-manifold M . If X is compact, then the fundamental group of each component of $M \setminus X$ is finitely generated, and $H_*(M \setminus X)$ is finitely generated.*

PROOF. It follows from Lemma 5.8 that $M \setminus X$ is homotopy equivalent to the exterior E_X of X . Each connected component of E_X is a compact 4-manifold since we assume that M is compact. The corollary is now a consequence of Corollary 3.18. \square

Proposition 5.10. *Let $X \subseteq M$ be a compact, proper 2-dimensional orientable submanifold of a compact, orientable 4-manifold M , such that each connected component of X has nonempty boundary. Then the tubular neighbourhood of Theorem 5.5 is homeomorphic to $X \times D^2$.*

PROOF. Connected surfaces with nonempty boundary are homotopy equivalent to wedges of circles. Every orientable linear D^n -bundle over a space X is classified up to isomorphism by a homotopy class of maps $X \rightarrow \text{BSO}(n)$. As $\text{SO}(n)$ is connected, so $\text{BSO}(n)$ is simply connected, and any orientable linear disc bundle over X is trivial. In particular, this is true when $n = 2$. \square

5.2. NORMAL VECTOR BUNDLES

The reader will be familiar with the definition of a normal vector bundle when working in the smooth category: if $X \subseteq M$ is a smooth submanifold of a smooth manifold, then the normal vector bundle is defined as the quotient of the vector bundle $TM|_X$ by the subbundle TX . This definition uses the smooth structure to ensure the existence of tangent vector bundles, and vector bundles are a strong enough bundle technology to ensure the existence of quotient bundles. While some (weaker) canonical tangential structures do exist in the topological category (see Chapter 6), the idea of a ‘quotient bundle’ no longer makes sense for them.

In the topological category, following [FQ90, Section 9], we will use a definition of normal vector bundle that is much closer to the geometry of tubular neighbourhoods. We begin with a definition that is almost what we need but suffers from a slight technical problem, which we then remedy.

Definition 5.11. Let M be an n -manifold and let X be a proper k -dimensional submanifold. An *internal linear bundle over X* is a pair $(E, p: E \rightarrow X)$ with the following properties.

- (1) E is a neighbourhood of X ;
- (2) E is a codimension zero submanifold of M ;
- (3) the map $p: E \rightarrow X$ is an $(n - k)$ -dimensional vector bundle such that $p(x) = x$ for all $x \in X$;
- (4) $\partial E = p^{-1}(\partial X)$.

An internal linear bundle $(E, p: E \rightarrow X)$ is intended to mirror the notion, from the smooth category, of an *open* tubular neighbourhood of X . As such, the definition as stands suffers from the potential technical problem that the closure of E in M , which should be a closed tubular neighbourhood, may no longer be a submanifold; see Figure 5.2. As in [FQ90, p. 137], we use the following additional idea to rule out this problem.

Definition 5.12. Let M be an n -manifold, let X be a proper k -dimensional submanifold, and let $(E, p: E \rightarrow X)$ be an internal linear bundle over X . Suppose that given any $(n - k)$ -dimensional vector bundle $(F, q: F \rightarrow X)$, any radial homeomorphism from an open convex disc bundle of F to E can be extended to a homeomorphism from the whole of F to a neighbourhood of E . Then we say $(E, p: E \rightarrow X)$ is *extendable*.




FIGURE 5.2. Non-extendable internal linear bundle.

Now we can define the notion of a normal vector bundle.

Definition 5.13. Let M be a n -manifold and let X be a proper k -dimensional submanifold. A *normal vector bundle* for X is an internal linear bundle that is extendable. (Note that the same concept is called a *normal bundle* on [FQ90, p. 137], we prefer the name *normal vector bundle*.)

Theorem 5.14. (Existence of normal vector bundles) *Every proper submanifold of a compact 4-manifold admits a normal vector bundle.*

Remark 5.15. Generally, in high dimensions, the existence of normal vector bundles is peculiar to when the submanifold has low dimension or low codimension. We refer the reader to [FQ90, Section 9.4] for a discussion of the other known situations where these objects always exist. Here is a summary of the known cases. A submanifold of dimension at most 3 in a closed manifold of dimension at least 5 has a normal vector bundle [FQ90, p. 150]. Codimension one submanifolds have normal vector bundles [Bro62, Theorem 3]. That every codimension two submanifold of a manifold of dimension not equal to four has a normal vector bundles was shown in [KS75], and this was extended to include dimension four in [FQ90, Section 9.3]. It is striking that, while among smooth manifolds dimension 4 exhibits worse than usual behaviour, in the topological category the existence of normal vector bundles seems to show that in this respect it is among the better behaved of the dimensions.

For the proof of Theorem 5.14 we will essentially appeal to results of [FQ90]. We reproduce these results here for the benefit of the reader.

Theorem 5.16. *Let N be a proper submanifold of a 4-manifold M , with a closed subset $\partial N \subseteq K \subseteq N$ and a normal vector bundle over some neighbourhood U of K in N . Then there is a normal vector bundle over N that agrees with the given one over some neighbourhood $V \subseteq U$ of K . Moreover this extension is unique up to ambient isotopy relative to some neighbourhood $W \subseteq V$ of K .*

PROOF OF THEOREM 5.16. The existence statement of the theorem follows immediately from [FQ90, Theorem 9.3A] and the uniqueness statement of the theorem follows immediately from [FQ90, Theorem 9.3D]. \square

PROOF OF THEOREM 5.14. Let X be a proper submanifold of a compact 4-manifold M . The case that X has no boundary follows immediately from the existence statement of Theorem 5.16 with $K = \emptyset$. The case that X has nonempty boundary follows also from Theorem 5.16 if we apply more care. We sketch the argument.

First it is well-known that given any pair (X, Σ) where X is a 3-manifold and Σ is a proper submanifold, then there exists a smooth structure on X such that Σ is a smooth submanifold. Even though this fact is well-known and often used, it is hard to give complete references. The existence of a smooth structure on X follows from [Moi77, p. 252 and 253] and [Mun60b, Theorems 6.2 and 6.3]. The extra complication of having a proper submanifold is partly taken care of by [Bin83, Theorem XVIII.4.B]. Thus we can view the submanifold $\partial X \subseteq \partial M$ as a smooth submanifold. Hence it has a smooth normal vector bundle, see e.g. [Kos93, Chapter III.2] or [Lan02b, Section IV.5].

Next use the Collar Neighbourhood Theorem 2.21 to obtain a collar $\partial M \times [0, 1] \subseteq M$ that restricts to a collar $\partial X \times [0, 1]$ for the boundary of X . Extend the smooth tubular neighbourhood of $\partial X \subseteq \partial M$ into the collar by taking a product with $[0, 1]$.

Finally, consider the 4-manifold without boundary $M' := M \setminus (\partial M \times [0, \frac{1}{2}])$. What remains of X is a submanifold $N := X \setminus (\partial X \times [0, \frac{1}{2}])$. The submanifold N already has a preferred normal vector bundle on the closed subset $K := \partial X \times (1/2, 1]$. Now apply Theorem 5.16 to the triple (M', N, K) to obtain a normal vector bundle $E \rightarrow N$ agreeing with the given one on K . The normal vector bundles over N and $\partial X \times [0, 1]$ agree on the overlap K . Thus they define a normal vector bundle on all of X . \square

Next we turn to the uniqueness of normal vector bundles.

Theorem 5.17. (Uniqueness of normal vector bundles) *Let M be a compact 4-manifold and let X be a proper submanifold of M . Suppose we are given two normal vector bundles $p_i: E_i \rightarrow X$, $i = 1, 2$ over X . For $i = 1, 2$ let $\iota_i: E_i \rightarrow M$ be the inclusion map. Then there exists a bundle isomorphism $f: E_1 \xrightarrow{\cong} E_2$ such that $\iota_2 \circ f$ and ι_1 are ambiently isotopic rel. X .*

PROOF. If X has no boundary, then the theorem is an immediate consequence of Theorem 5.16. Now suppose that X has nonempty boundary.




FIGURE 5.3. Illustration of the proof of Theorem 5.14.

First we claim that any normal vector bundle of X is obtained by the construction outlined in the proof of Theorem 5.14. To see this, let $p: E \rightarrow X$ be a normal vector bundle. Pick a collar neighbourhood $\partial X \times [0, 2] \subseteq X$. Since p is extendable, we can view p as the interior of a disc bundle $q: F \rightarrow X$ in M . Write $C := q^{-1}(\partial X) \subseteq \partial M$. The disc bundle $q: q^{-1}(\partial X \times [0, 2]) \rightarrow \partial X \times [0, 2]$ defines a collar neighbourhood $C \times [0, 2]$ for the compact submanifold C of ∂M . By the Collar Neighbourhood Theorem 2.16 we can extend the collar neighbourhood $C \times [0, 1]$ of C to a collar neighbourhood $\partial M \times [0, 1]$. With this choice of collar neighbourhood of ∂M , the construction in the proof of Theorem 5.14, with further appropriate choices, gives rise to the normal vector bundle $p: E \rightarrow X$. This completes the proof of the claim.

After this long preamble it suffices to prove the theorem for any two normal vector bundles obtained as in the proof of Theorem 5.14. Uniqueness follows by arguing that each step in the proof of existence of normal vector bundles was essentially unique. The proofs of uniqueness in the three steps make use of the following ingredients.

First, apply the uniqueness statement for normal vector bundles of submanifolds of smooth manifolds to $\partial X \subseteq \partial M$ e.g. [Kos93, Chapter III.2] or [Lan02b, Section IV.5].

Next use the uniqueness of collar neighbourhoods as formulated in Theorem 2.18, applied to the two collar neighbourhoods of ∂M subordinate to the given normal vector bundles of X .

Finally apply the full relative version of Theorem 5.16 to extend the normal vector bundle uniquely over the rest of X . \square

5.3. TUBULAR NEIGHBOURHOODS: PROOFS

Now we will use the results from the previous section to prove Theorems 5.5 and 5.6, i.e. we will prove the existence and uniqueness of tubular neighbourhoods. First we show how one can obtain tubular neighbourhoods from normal vector bundles.

Definition 5.18.

- (1) A *form* over a real vector space V is an \mathbb{R} -bilinear symmetric map $g: V \times V \rightarrow \mathbb{R}$. It is called *positive definite* if for every $v \in V \setminus \{0\}$ we have $g(v, v) > 0$.

- (2) Let $p: E \rightarrow X$ be a vector bundle over a topological space X . Given $x \in X$, write $E_x := p^{-1}(x)$. A *positive definite form* $g = \{g_x\}_{x \in X}$ consists of a positive definite form g_x over every E_x such that g_x changes continuously with x .

Lemma 5.19. *Let X be a compact manifold and let $p: E \rightarrow X$ be an n -dimensional vector bundle. Then the space of positive definite forms on E is nonempty and convex. Furthermore, let $g = \{g_x\}_{x \in X}$ be a positive definite form on E and consider the map*

$$p: E(g) := \bigcup_{x \in X} \{v \in E_x \mid g_x(v, v) \leq 1\} \rightarrow X.$$

This map has the following properties.

- (1) *The map $p|_{E(g)}: E(g) \rightarrow X$ is a linear D^n -bundle.*
- (2) *Given two different positive definite forms g and h on E there exists a continuous map $H: E \times [0, 1] \rightarrow E$ with the following properties.*
 - (a) *We have $H_0 = \text{Id}$.*
 - (b) *The map H_1 restricts to an isomorphism $H_1: E(g) \rightarrow E(h)$ of linear D^n -bundles.*

PROOF.

- (1) This statement can be proved easily using the observation that the set of positive definition \mathbb{R} -bilinear symmetric forms on a real vector space is nonempty and convex.
- (2) The proof of this statement follows from the same argument as [Wal16, Lemmas 2.5.2 and 2.5.4]. \square

Let X be a compact manifold and let $p: E \rightarrow X$ be a vector bundle. Given a positive definite form g on E we refer to

$$p: E(g) := \bigcup_{x \in X} \{v \in E_x \mid g_x(v, v) \leq 1\} \rightarrow X$$

as a *corresponding disc bundle*. It follows from Lemma 5.19 that for most purposes the precise choice of g is irrelevant.

We can now prove the existence of tubular neighbourhoods.

PROOF OF THE TUBULAR NEIGHBOURHOOD THEOREM 5.5. Let X be a compact proper submanifold of a 4-manifold M . By Theorem 5.14 there exists a normal vector bundle $p: N \rightarrow X$ for X . By Lemma 5.19 (1) there exists a corresponding disc bundle. Using the uniqueness statement of Lemma 5.19 (2) locally one can show that this disc bundle is a submanifold with corner and a tubular neighbourhood. \square

The uniqueness proof for tubular neighbourhoods also requires us to associate a normal vector bundle to a tubular neighbourhood.

Lemma 5.20. *Let M be a compact 4-manifold and let X be a compact proper k -dimensional submanifold. Let $p: N \rightarrow X$ be a tubular neighbourhood for X . There exists a normal vector bundle $q: E \rightarrow X$ and a positive definite form g such that $N = E(g)$ and $p: N \rightarrow X$ equals $q: E(g) \rightarrow X$.*

We call $q: E \rightarrow X$ a *corresponding normal vector bundle*.

PROOF. Let $p: N \rightarrow X$ be a tubular neighbourhood for X . Recall that we have $\text{Int } N = N \setminus \partial_0 N$. Consider $W := M \setminus \text{Int } N$. This is a compact 4-manifold. Pick a collar neighbourhood $\partial W \times [0, 1]$ and set $E := N \cup (\partial_0 N \times [0, \frac{1}{2}))$. We have an obvious projection map $q: E \rightarrow X$ turning q into a bundle map where the fibre is given by the open $(4-k)$ -ball of radius $\frac{3}{2}$. We leave it to the reader to turn $q: N \rightarrow X$ into an internal linear bundle, to show that it is in fact extendable (at this point one has to use that in the definition of E we only used “half” of the collar neighbourhood $\partial_0 N \times [0, 1]$), and to equip N with a positive definite form g such that $N = E(g)$. \square

We conclude the chapter with the proof of the uniqueness theorem for tubular neighbourhoods.

PROOF OF THEOREM 5.6. Let M be a 4-manifold and let X be a compact proper k -dimensional submanifold. Furthermore let $p_i: N_i \rightarrow X$, $i = 1, 2$ be two tubular neighbourhoods of X . For $i = 1, 2$, let $q_i: E_i \rightarrow X$ be two corresponding normal vector bundles and let g_i be the positive definite forms provided by Lemma 5.20. It follows from Theorem 5.17 that there exists a bundle isomorphism $f: E_1 \xrightarrow{\cong} E_2$ such that $\iota_2 \circ f$ and ι_1 are ambiently isotopic rel. X . It follows from the definitions that N_2 is equivalent to the disc bundle defined by f^*g_2 on E_1 . It follows from the Lemma 5.19 (2) together with the Isotopy Extension Theorem 2.20 that f^*g_2 and g_1 define equivalent tubular neighbourhoods. (Strictly speaking we did not formulate the Isotopy Extension Theorem 2.20 for submanifolds with corner, but it is not difficult to prove a generalization.) \square

CHAPTER 6

Background on bundle structures

In this chapter we recall the bundle technologies we will need to use in later chapters. The three standard manifold categories smooth (Diff), piecewise linear (PL) and topological (TOP) each have corresponding bundle types, with fibre \mathbb{R}^n . We first discuss the topological groups $\text{TOP}(n)$ and $\text{O}(n)$ which are structure groups for fibre bundles with fibre \mathbb{R}^n , corresponding to the TOP and Diff categories. There are natural topologies on $\text{TOP}(n)$ and $\text{O}(n)$, so that it is relatively straightforward to discuss the classification of these bundle types. We next discuss the \mathbb{R}^n -bundles that correspond to the PL category. In contrast to $\text{TOP}(n)$ and $\text{O}(n)$ there is no obvious appropriate topology on the group of piecewise-linear homeomorphisms of \mathbb{R}^n that fix the origin. For this reason, in the PL category it appears there is no choice but to delve into a more sophisticated approach to classify the bundles of interest, and we approach this via simplicial groups. We finish the section with a discussion of topological *microbundles*.

6.1. TOPOLOGICAL, SMOOTH, AND PIECEWISE LINEAR \mathbb{R}^n -BUNDLES

Before we turn to the different flavours of \mathbb{R}^n -bundles let us first recall some general facts about bundles.

Definition 6.1. Let G be a topological group G . A *universal principal G -bundle* is a principal G -bundle $p: EG \rightarrow BG$ such that the following two conditions are satisfied.

- (1) The space BG is a CW complex.
- (2) Given any principal G -bundle $q: F \rightarrow C$, where C is a topological space that is homotopy equivalent to a CW complex, there exists a map $f: C \rightarrow BG$, unique up to homotopy, such that q is isomorphic to the pullback bundle f^*EG .

The base space BG is called a *classifying space for G* .

Proposition 6.2. *Given a topological group G , there exists a universal principal G -bundle $p: EG \rightarrow BG$. This principal bundle is unique up to fibre homotopy equivalence.*

PROOF. The Milnor join construction [Mil56], [Hus93] or [Sta71], or alternatively the geometric bar construction [May75], gives an explicit principal G -bundle $p: E \rightarrow B$ which has the universal property for all numerable principal G -bundles. Since principal G -bundles over CW complexes are numerable, the Milnor bundle has the universal property for principal G -bundles over CW complexes.

The topological space B is not necessarily a CW complex. Thus we apply the CW approximation theorem to get a CW complex \tilde{B} and a weak homotopy equivalence $\varphi: \tilde{B} \rightarrow B$. The pullback bundle φ^*E over \tilde{B} has the desired properties.

The uniqueness of universal principal G -bundles over CW complexes follows from a standard argument about universal objects. \square

Definition 6.3. Given $n \in \mathbb{N}_0$ let $\text{TOP}(n)$ be the subgroup of homeomorphisms of \mathbb{R}^n that fix the origin, topologised using the compact open topology. A principal $\text{TOP}(n)$ -bundle has an associated fibre bundle with fibre \mathbb{R}^n and a preferred 0-section. Call such a bundle a *topological \mathbb{R}^n -bundle*. Let TOP be the colimit $\text{colim} \text{TOP}(n)$, in the category of topological groups, under the inclusions

$$\begin{aligned} \text{TOP}(n) &\rightarrow \text{TOP}(n+1) \\ (f: \mathbb{R}^n \rightarrow \mathbb{R}^n) &\mapsto (f \times \text{Id}_{\mathbb{R}}: \mathbb{R}^n \times \mathbb{R} \rightarrow \mathbb{R}^n \times \mathbb{R}). \end{aligned}$$

We obtain the corresponding classifying spaces $\text{BTOP}(n)$ and BTOP .

Definition 6.4. Given $n \in \mathbb{N}_0$ let $\text{O}(n)$ be the orthogonal homeomorphisms of \mathbb{R}^n that fix the origin, topologised in the standard way as a subspace of a vector space $\text{O}(n) \subseteq \text{M}(n \times n, \mathbb{R}) \cong \mathbb{R}^{n^2}$. A principal $\text{O}(n)$ -bundle has an associated fibre bundle with fibre \mathbb{R}^n and a preferred 0-section, and such a bundle is in particular a vector bundle. Define O , $\text{BO}(n)$ and BO analogously to Definition 6.3.

We also introduce the following structure group, for comparison, and use in Chapter 7.

Definition 6.5. Given $n \in \mathbb{N}_0$ let $\text{Diff}(n)$ be the subgroup of diffeomorphisms of \mathbb{R}^n that fix the origin, topologised using the weak C^∞ topology [Hir94, § 2.1] (also called the (weak) *Whitney* topology). A principal $\text{Diff}(n)$ -bundle has an associated fibre bundle with fibre \mathbb{R}^n and a preferred 0-section. Call such a bundle a *Diff \mathbb{R}^n -bundle*. Define Diff , $\text{BDiff}(n)$ and BDiff analogously to Definition 6.3.

Remark 6.6.

- (1) The pullback topology on $\text{Diff}(n)$ under the inclusion map $\text{Diff}(n) \rightarrow \text{TOP}(n)$ is by definition the coarsest topology such that $\text{Diff}(n) \rightarrow \text{TOP}(n)$ is continuous. It equals the compact open topology on the set $\text{Diff}(n)$ (equivalent to the weak C^0 topology), not the weak C^∞ topology. In other words, one does not topologise $\text{Diff}(n)$ as a subspace of $\text{TOP}(n)$.
- (2) It is nevertheless the case that $\text{Diff}(n) \rightarrow \text{TOP}(n)$ is a continuous map with respect to the weak C^∞ topology on $\text{Diff}(n)$ and the compact open topology on $\text{TOP}(n)$. There are induced maps $\text{BDiff}(n) \rightarrow \text{BTOP}(n)$ for each n , and $\text{BDiff} \rightarrow \text{BTOP}$.
- (3) On the other hand, we now argue that *both* inclusion maps $\text{GL}(n, \mathbb{R}) \rightarrow \text{TOP}(n)$ and $\text{GL}(n, \mathbb{R}) \rightarrow \text{Diff}(n)$ induce the standard topology on $\text{GL}(n, \mathbb{R})$ via pullback (which is a little surprising at first glance). In the former case, simply note that the standard topology on $\text{GL}(n, \mathbb{R})$ as a vector subspace of \mathbb{R}^{n^2} is equivalent to the compact open topology, or equivalently the weak C^0 topology. In the latter case, note that as the

maps in $\mathrm{GL}(n, \mathbb{R})$ are linear, and the weak C^r topologies for $0 \leq r < \infty$ are defined in terms of partial derivatives [Hir94, § 2.1], they are all equivalent topologies on $\mathrm{GL}(n, \mathbb{R})$. As the weak C^∞ topology is the limit of the C^r topologies, this proves the statement.

(4) The map

$$\begin{aligned} \mathrm{Diff}(n) \times [0, 1] &\rightarrow \mathrm{Diff}(n) \\ (f, t) &\mapsto \left(\begin{array}{l} \mathbb{R}^n \rightarrow \mathbb{R}^n \\ x \mapsto \begin{cases} \frac{1}{t} \cdot f(tx), & \text{if } t \neq 0, \\ \mathrm{D}f_0 \cdot x \end{cases} \end{array} \right) \end{aligned}$$

is a deformation retraction from $\mathrm{Diff}(n)$ to $\mathrm{GL}(n, \mathbb{R})$. Finally note that the Gram-Schmidt process can be used to determine a homotopy equivalence $\mathrm{GL}(n, \mathbb{R}) \simeq \mathrm{O}(n)$. Thus \mathbb{R}^n -bundles with structure group $\mathrm{Diff}(n)$ always admit a (unique) structure group reduction to ordinary vector bundles whose transition functions lie in $\mathrm{O}(n)$.

The following proposition shows that BTOP and BDiff classify CAT \mathbb{R}^n -bundles when CAT is TOP or Diff .

Proposition 6.7. *For $\mathrm{CAT} = \mathrm{TOP}$ or Diff , and for each $n \geq 0$, the space $\mathrm{BCAT}(n)$ is homotopy equivalent to a CW complex, and there exists a CAT \mathbb{R}^n -bundle γ_n^{CAT} over $\mathrm{BCAT}(n)$, such that for every CAT \mathbb{R}^n -bundle ξ over a CW complex, there is a CAT \mathbb{R}^n -bundle map $F: \xi \rightarrow \gamma_n^{\mathrm{CAT}}$, unique up to homotopy of CAT \mathbb{R}^n -bundle maps. In particular $F^*(\gamma_n^{\mathrm{CAT}}) \cong \xi$.*

PROOF. Let X be a space homotopy equivalent to a CW complex. There is a 1:1 correspondence between isomorphism classes of $\mathrm{CAT}(n)$ \mathbb{R}^n -bundles and isomorphism classes of principal $\mathrm{CAT}(n)$ -bundles over X . For more details, see e.g. [Swi02, Proposition 11.22].

Applying this to the universal principal $\mathrm{CAT}(n)$ -bundle of Proposition 6.2, we obtain a CAT \mathbb{R}^n -bundle γ_n^{CAT} over $\mathrm{BCAT}(n)$. The desired universal property for γ_n^{CAT} is inherited from the universal property of the universal principal $\mathrm{CAT}(n)$ -bundle over $\mathrm{BCAT}(n)$ \square

We finally move on to the piecewise linear category. For background on piecewise linear topology see [RS72].

Definition 6.8. A continuous map $f: K \rightarrow L$ between two simplicial complexes K and L is *piecewise linear* (PL) if there are subdivisions K' of K and L' of L such that $f: K' \rightarrow L'$ is a simplicial map.

Definition 6.9. A PL \mathbb{R}^n -bundle is a topological \mathbb{R}^n -bundle $p: E \rightarrow B$, where both E and B are simplicial complexes, where $p: E \rightarrow B$ and the 0-section are both PL maps, and where for every simplex $\Delta \subseteq B$ there exists a PL homeomorphism φ such that the composition $p^{-1}(\Delta) \xrightarrow{\varphi} \Delta \times \mathbb{R}^n \xrightarrow{\mathrm{pr}_1} \Delta$ is equal to the projection p .

The definition of the classifying space for PL \mathbb{R}^n -bundles is a little more involved than the constructions for $\mathrm{O}(n)$ and $\mathrm{TOP}(n)$, using the technology of semi-simplicial groups. For

a gentle introduction to simplicial sets, see [Fri12]. The canonical reference for classifying spaces constructed using simplicial groups is [May75].

Remark 6.10. When navigating the various references we use below, the reader should be aware that the terminology for simplicial groups and simplicial sets has changed over the years and there are some clashes between current usage and previous usage. What we are calling a *simplicial set* is, for example, what is defined in [Fri12, Definition 3.2]. In particular, note that this has both *face* and *degeneracy* maps. This object has historically been called a *complete semi-simplicial set (c.s.s.)*, which has unfortunately in the past been abbreviated to just “semi-simplicial set”. The modern terminology reserves *semi-simplicial set* for a simplicial set without degeneracy maps as part of the data. To add to the confusion, a simplicial set, but without degeneracies, has also historically been called a Δ -set, although in modern terminology this is usually reserved for the semi-simplicial version of a simplicial *complex*.

Let Δ^k be the standard k -simplex. Recall a *simplicial group* is a simplicial object in the category of groups, that is a contravariant functor from the simplicial category to the category of groups $\Delta \rightarrow \mathbf{Group}$.

Definition 6.11. Given $n \in \mathbb{N}_0$ let $\mathrm{PL}(n)_\bullet$ be the simplicial group defined as follows.

- (i) The group $\mathrm{PL}(n)_k$ assigned to the k -simplex is the group of PL \mathbb{R}^n -bundle isomorphisms of the trivial bundle $f: \mathbb{R}^n \times \Delta^k \rightarrow \mathbb{R}^n \times \Delta^k$. That is, f is a PL homeomorphism preserving the 0-section and commuting with the projection to Δ^k .
- (ii) A morphism $\lambda: \Delta^\ell \rightarrow \Delta^k$ is sent to the morphism $\lambda^\#$ which assigns to $f: \mathbb{R}^n \times \Delta^k \rightarrow \mathbb{R}^n \times \Delta^k$ the map such that the diagram below commutes

$$\begin{array}{ccc} \mathbb{R}^n \times \Delta^\ell & \xrightarrow{\lambda^\#(f)} & \mathbb{R}^n \times \Delta^\ell \\ \downarrow \lambda \times 1 & & \downarrow \lambda \times 1 \\ \mathbb{R}^n \times \Delta^k & \xrightarrow{f} & \mathbb{R}^n \times \Delta^k. \end{array}$$

Define $\mathrm{PL}(n)$ as the topological group [Mil57, Theorem 3] realising the simplicial group $\mathrm{PL}(n)_\bullet$. Then define $\mathrm{BPL}(n)$ using Proposition 6.2. Define PL and BPL as colimits, analogously to Definition 6.3.

Equivalently, one can define $\mathrm{BPL}(n)$ by first using the geometric bar construction level-wise on $\mathrm{PL}(n)_\bullet$ to obtain a simplicial space $\mathrm{BPL}(n)_\bullet$, and then geometrically realising to obtain a space $\mathrm{BPL}(n)$; see e.g. [ERW19, §1.2] for the equivalence to the previous definition.

Remark 6.12. Consider the subgroup of $\mathrm{TOP}(n)$ consisting of PL homeomorphisms (note this is *not* the definition of what we called $\mathrm{PL}(n)$ above. In particular $\mathrm{PL}(n)$ is a much larger set). One might think that this group, together with the subspace topology from $\mathrm{TOP}(n)$, is a realistic way to circumvent the construction above with simplicial groups. This would certainly produce a topological group that classifies *some* category of \mathbb{R}^n bundles, but

we do not know whether it classifies the PL \mathbb{R}^n -bundles defined above (it seems unlikely, cf. Remark 6.6 (1)).

The following proposition can now be viewed as an analogue of Proposition 6.2.

Proposition 6.13. *For each $n \geq 0$, the space $\text{BPL}(n)$ is homotopy equivalent to a CW complex, and there exists a PL \mathbb{R}^n -bundle γ_n^{PL} over $\text{BPL}(n)$, such that for every PL \mathbb{R}^n -bundle ξ over a CW complex, there is a PL \mathbb{R}^n -bundle map $F: \xi \rightarrow \gamma_n^{\text{PL}}$, unique up to homotopy of PL \mathbb{R}^n -bundle maps. In particular $F^*(\gamma_n^{\text{PL}}) \cong \xi$.*

PROOF. In [Mil61, p. 24], Milnor constructed a simplicial group $(\text{PL}_n)_\bullet$. It is defined similarly to $\text{PL}(n)_\bullet$, but differs from what we have done in Definition 6.11(i) by specifying that the maps f are rather PL microbundle isomorphisms of the trivial PL microbundle over the simplex (in particular are *germs* of the types of maps f we are using). The simplicial space $(\text{BPL}_n)_\bullet$ that results from the level-wise geometric bar construction is a classifying space for simplicial principal $(\text{PL}_n)_\bullet$ -bundles. In [Mil61, §5], Milnor showed that the geometric realisation $\text{BPL}_n := |(\text{BPL}_n)_\bullet|$ is homotopy equivalent to a locally finite simplicial complex (so in particular a CW complex), and that there is a universal rank n PL microbundle over BPL_n .

Kuiper-Lashof [KL66, Theorem 1] proved that each rank n PL microbundle is the underlying microbundle of a unique PL \mathbb{R}^n -bundle. Apply this to the universal rank n PL microbundle over BPL_n , to obtain a universal PL \mathbb{R}^n -bundle over BPL_n . The map $g: \text{PL}(n)_\bullet \rightarrow (\text{PL}_n)_\bullet$, given by simplex-wise taking the germs of maps f as in Definition 6.11(i), is a homotopy equivalence of simplicial sets [KL66, Lemma 1.6(f)]. This induces a homotopy equivalence $\text{BPL}(n) \xrightarrow{\cong} \text{BPL}_n$. Use this latter homotopy equivalence to pull back the universal rank n PL microbundle over BPL_n to the desired base space $\text{BPL}(n)$. \square

Remark 6.14.

- (1) The simplicial method used above for PL \mathbb{R}^n -bundles can be used in the smooth and topological categories as well, giving a uniform treatment. The resulting classifying spaces for the smooth and topological categories are homotopy equivalent to the spaces $\text{BO}(n)$ and $\text{BTOP}(n)$, defined earlier, by the universal property.
- (2) An alternative uniform proof of Propositions 6.7 and 6.13, for all three categories TOP, PL and Diff simultaneously, was given by Kirby-Siebenmann [KS77, Essay IV, Proposition 8.1, p. 181]. Instead of a simplicial approach, they use E. H. Brown's theory of representability to obtain universal CAT *microbundles* (cf. Section 6.2) over classifying spaces which are locally finite simplicial complexes. The theorems of Kister (Theorem 6.17) and Kuiper-Lashof [KL66, Theorem 1], together with the analogue for smooth microbundles, means that the Kirby-Siebenmann universal CAT microbundles can then be upgraded to universal CAT \mathbb{R}^n -bundles with the same base space, similarly to how we worked at the end of the proof of Proposition 6.13. The Kirby-Siebenmann proof also includes a statement relative to a closed subspace of the base.

6.2. MICROBUNDLES

All smooth manifolds have tangent vector bundles and all smooth submanifolds have normal vector bundles. This is one reason that vector bundles, corresponding to the structure group $O(n)$ are the de facto bundle technology in the smooth category. A general difficulty we will face when talking about manifold transversality in Chapter 10 is that we will need to use some well-defined notion of normal structure for a submanifold and, outside of the smooth category, submanifolds do not necessarily admit normal vector bundles. However, various weaker bundle technologies have been developed, which replace this crucial concept in the topological category.

This subsection is devoted to a discussion of *microbundles*, which were introduced by Milnor in [Mil64]. The existence and uniqueness of tangent and (stable) normal microbundles leads to the existence and uniqueness of tangent and (stable) normal TOP \mathbb{R}^n -bundles, via Kister's Theorem (Theorem 6.17). Source material on microbundles is not hard to find in the literature, but has been included here for the convenience of the reader, in order for this survey to be more self-contained.

The interaction between the weaker structure groups $PL(n)$ and $TOP(n)$ for tangent and (stable) normal \mathbb{R}^n -bundles, and the topological/PL/smooth structures on the manifold itself are the topic of *smoothing theory*, to which we turn in Chapter 9.

Definition 6.15. An n -dimensional *microbundle* ξ consists of a base space B and a total space E sitting in a diagram

$$B \xrightarrow{i} E \xrightarrow{p} B,$$

such that $p \circ i = \text{Id}_B$, and that is *locally trivial* in the following sense: for every point $b \in B$, there exists an open neighbourhood U of B , an open neighbourhood V of $i(b)$ and a homeomorphism $\phi_b: V \rightarrow U \times \mathbb{R}^n$ such that

$$\begin{array}{ccc} & V & \\ i \nearrow & & \searrow p \\ U & & U \\ u \mapsto (u,0) \searrow & \downarrow \phi_b & \nearrow (u,v) \mapsto u \\ & U \times \mathbb{R}^n & \end{array}$$

commutes. We refer to i as the *inclusion map* of the microbundle and we refer to p as the *projection*.

Note that we only require neighbourhoods of the points $i(b)$ to be trivial, and not all of the fibre $p^{-1}(b)$. In fact, we only care about neighbourhoods $i(B) \subseteq E$, and declare two microbundles $B \xrightarrow{i} E \xrightarrow{p} B$ and $B \xrightarrow{i'} E' \xrightarrow{p'} B$ to be *equivalent*, if $i(B)$ and $i'(B)$ have homeomorphic neighbourhoods such that the homeomorphism commutes with both the inclusion map and the restriction of the projection map.

Definition 6.16. Let $B \xrightarrow{i} E \xrightarrow{p} B$ be a microbundle ξ and let $f: A \rightarrow B$ be a map. The *pullback* of ξ under f is the microbundle $f^*\xi$ with total space

$$f^*E = \{(a, e) \in A \times E \mid f(a) = p(e)\},$$

projection $(f^*p)(a, e) = a$, and injection $(f^*i)(a) = (a, i(f(a)))$. In the case that f is an inclusion, also consider the microbundle $\xi|_A$, which has total space $p^{-1}(A) \subseteq E$, and projection $p|_{p^{-1}(A)}: p^{-1}(A) \rightarrow A$ and injection $i_A: A \rightarrow p^{-1}(A)$ are both the restrictions of p, i . In this case, the map of total spaces $(a, e) \mapsto e$ gives a preferred isomorphism $f^*\xi$ to $\xi|_A$.

A topological \mathbb{R}^n -bundle clearly has an underlying microbundle. Kister proved the surprising result that *every* microbundle over a manifold is equivalent to such an underlying microbundle [Kis64, Theorem 2 and Corollary 1].

Theorem 6.17. (Kister's Theorem) *Let B be a manifold and $B \xrightarrow{i} E \xrightarrow{p} B$ be an n -dimensional microbundle ξ . Then there exists an open set $F \subseteq E$ containing $i(B)$ such that $p|_F: F \rightarrow B$ is the projection map of a topological \mathbb{R}^n -bundle, whose 0-section is i and whose underlying microbundle is ξ . Moreover, if F_1 and F_2 are any two topological \mathbb{R}^n -bundles over B such that the underlying microbundles are equivalent, then F_1 and F_2 are isomorphic as topological \mathbb{R}^n -bundles.*

Every manifold admits a tangent microbundle.

Definition 6.18. The *tangent microbundle* of an n -dimensional manifold M is the microbundle $M \xrightarrow{\Delta} M \times M \xrightarrow{(x,y) \mapsto x} M$ where Δ is the diagonal map. Kister's theorem implies this corresponds to a unique *topological tangent bundle* $\tau_M: M \rightarrow \text{BTOP}(n)$, with corresponding *stable topological tangent bundle* $\tau_M: M \rightarrow \text{BTOP}$.

More subtle is the concept of a normal microbundle.

Definition 6.19. A *normal microbundle* of a submanifold S of a manifold M is a microbundle $S \rightarrow E \rightarrow S$ such that E is a neighbourhood of S in M and such that $S \rightarrow E$ is the inclusion.

It is immediate from the definition of normal microbundle that the local flatness in the definition of a submanifold S is a necessary condition for the existence of a normal microbundle. For example wild knots and the Alexander horned sphere do not admit normal microbundles. Indeed, it is generally far from straightforward to prove the existence of normal microbundles at all. Here is an existence and uniqueness result due to Stern [Ste75, Theorem 4.5]. See also [Hir66], [Hir68, p. 65], and [KS77, Essay IV, Appendix A, p. 203].

Theorem 6.20. *Let M^{n+q} be a manifold, and let $N^n \subseteq M^{n+q}$ be a proper submanifold of codimension q . Suppose that $n \leq q + 1 + j$ and $q \geq 5 + j$ for some $j = 0, 1, 2$. Then N admits a normal microbundle restricting to a normal microbundle of $\partial N \subseteq \partial M$.*

If in addition $n \leq q + j$, then this normal microbundle is unique up to isotopy.

Remark 6.21 (Unique up to isotopy). For a submanifold $N \subseteq M$ we say a normal microbundle $N \xrightarrow{i} \nu(N) \xrightarrow{p} N$ is *unique up to isotopy* if whenever there is another normal microbundle $N \xrightarrow{i'} \nu'(N) \xrightarrow{p'} N$, there exists a microbundle equivalence f between $\nu(N)$ and $\nu'(N)$ such that $p' \circ f$ is isotopic to p relative to N .

We exploit these theorems to define a stable topological normal structure on any closed manifold, that will play an important role in Chapter 9. Every closed smooth n -manifold M can be embedded in \mathbb{R}^k , for some k . The stable class of a normal vector bundle gives rise to *stable normal vector bundle of M* , denoted $\mu_M: M \rightarrow \text{BO}$. For k large enough, the embedding of M in \mathbb{R}^k is unique up to isotopy, and using this one can show that the stable normal bundle is uniquely determined up to isomorphism. We seek to adapt this construction to the topological category.

Consider that any closed n -manifold M can be embedded as a submanifold $M \subseteq \mathbb{R}^m$ for large m (this follows e.g. from [Hat02, Corollary A.9] together with [Mil72, Theorem 5]). For large enough m , any two such embeddings are isotopic. For large enough m , Theorem 6.20 implies there is a normal microbundle ξ . After possibly increasing m further, the last sentence of Theorem 6.20 implies this normal microbundle ξ is unique. By Kister's Theorem this defines a unique topological \mathbb{R}^{m-n} -bundle. We remove the dependence on m by passing to the stable bundle $\text{TOP}(m-n) \subseteq \text{TOP}$. Thus the process described gives a well-defined classifying map $\nu_M: M \rightarrow \text{BTOP}$. Summarising, we have the following.

Definition 6.22. Given any closed n -manifold, the topological \mathbb{R}^∞ -bundle $\nu_M: M \rightarrow \text{BTOP}$, described above, is called the *stable topological normal bundle*. It is well-defined and unique.

The next example shows that outside the hypotheses of Theorem 6.20, we should expect that normal microbundles can be very badly behaved.

Example 6.23. Normal microbundles do not necessarily exist. Rourke and Sanderson [RS67, Example 2] construct S^{19} as a submanifold of a certain 28-dimensional PL manifold M in such a way that it does not admit a topological normal microbundle. The embedding is even piecewise linear.

Hirsch's example of a 4-submanifold of S^7 from [Hir68, Theorem 4] mentioned in Remark 5.4 also does not admit a normal microbundle. Such a normal microbundle would contain a $\text{TOP}(3)$ bundle, and every $\text{TOP}(3)$ bundle can be improved to an $\text{O}(3)$ bundle, which in turn contains a tubular neighbourhood. So it follows that there can be no normal microbundle in Hirsch's example. Note that historically this deduction was not possible until Hatcher [Hat83] proved that $\text{BO}(3) \rightarrow \text{BTOP}(3)$ is a homotopy equivalence.

Even when topological normal microbundles exist, they are not always unique: Rourke and Sanderson consider the smooth standard embedding $S^{18} \subseteq S^{27}$ [RS68, Theorem 3.12] and construct a certain normal microbundle ξ of $S^{18} \subseteq S^{27}$. The construction of ξ is such that if ξ were concordant to the trivial normal microbundle, this concordance would induce a normal microbundle structure back on the embedding $S^{19} \subseteq M^{28}$ of the previous

paragraph. As this is not possible, ξ is nontrivial. Note that the normal vector bundle νS^{18} of the standard embedding is trivial, so $S^{18} \subseteq S^{27}$ admits at least two different normal microbundles.

The following theorem ensures the issues of the previous example are not seen in dimension 4.

Theorem 6.24. *Let X be a proper submanifold of a 4-manifold M . Then X admits a normal microbundle. Moreover, if ξ is a normal microbundle of X , it is the underlying microbundle to a normal vector bundle.*

PROOF. The existence of normal microbundles in ambient dimension 4 is an immediate consequence of the existence of normal vector bundles (Theorem 5.14), and this is the only proof of which we are aware for this fact. (It would be interesting to know of a more elementary proof.)

We denote by n the codimension of X in M . Given a normal microbundle ξ , we apply Kister's theorem 6.17 to obtain an embedded \mathbb{R}^n -bundle with underlying microbundle ξ . For $n \leq 3$, the homotopy fibre $\text{TOP}(n)/\text{O}(n)$ for the forgetful map $\text{BO}(n) \rightarrow \text{BTOP}(n)$ is contractible; see Proposition 7.5 for the relevant citations. Using these facts, and checking the obstructions in each of the cases $n = 0, 1, 2, 3, 4$, we see in each case the embedded topological \mathbb{R}^n -bundle can be upgraded to an embedded vector bundle. Choose such a vector bundle refinement. By restricting to an open disc bundle and rescaling we can ensure this internal linear bundle is extendable and thus is a normal vector bundle in the sense of Definition 5.12. \square

We will make use of our discussion of normal microbundles in Chapter 10 on topological transversality.

CHAPTER 7

Stiefel-Whitney classes

The well-known treatment of Stiefel-Whitney classes in Milnor-Stasheff [MS74] is for vector bundles, that is bundles over a space B with linear transition functions. In this chapter we discuss the analogous characteristic classes for TOP and PL \mathbb{R}^n -bundles. These arise, in particular, for tangent bundles of PL and topological manifolds respectively. Throughout this chapter we will use the terminology *CAT \mathbb{R}^n -bundle*, where

$$\text{CAT} \in \{\text{Diff}, \text{PL}, \text{TOP}\}.$$

When $\text{CAT} = \text{TOP}$, this will mean an \mathbb{R}^n -bundle with structure group $\text{TOP}(n)$. When $\text{CAT} = \text{PL}$, this will mean PL \mathbb{R}^n -bundles in the sense of Definition 6.9. When $\text{CAT} = \text{Diff}$, we will mean an \mathbb{R}^n -bundle with structure group $\text{O}(n)$, taking advantage of the structure group reduction described in Remark 6.6.

Generalising work of Thom [Tho52] in the case of orthogonal structure group, Fadell [Fad65, Definition 6.1] (see also Stern [Ste75]) defined Stiefel-Whitney classes of CAT \mathbb{R}^n -bundles, where $\text{CAT} \in \{\text{Diff}, \text{PL}, \text{TOP}\}$. As we shall see, the definitions of Stiefel-Whitney classes in the PL and TOP cases are analogous to the classical definition for vector bundles.

Remark 7.1. Even more generally, one can also develop Stiefel-Whitney classes for spherical fibrations, using an analogous definition. Passing to the underlying S^{n-1} -bundle of a CAT \mathbb{R}^n -bundle, the spherical fibration definition will recover all definitions we give below. However, we have chosen not work in this level of generality.

Let $\xi = (p: E \rightarrow B)$ be a CAT \mathbb{R}^n -bundle over some topological space B . Let E_0 denote the complement in the total space E of the zero section. Recall that the Thom isomorphism [Tho52] is an isomorphism

$$\Phi: H^i(B; \mathbb{Z}/2) \xrightarrow{\cong} H^{n+i}(E, E_0; \mathbb{Z}/2).$$

Fadell [Fad65, Theorem 5.2] checked that the Thom isomorphism holds in the present context, with $\text{PL}(n)$ or $\text{TOP}(n)$ structure group. The Steenrod squares are homomorphisms

$$\text{Sq}^i: H^n(E, E_0; \mathbb{Z}/2) \rightarrow H^{n+i}(E, E_0; \mathbb{Z}/2).$$

Here, let us recall that given a pair of spaces (X, A) , for each $i \geq 0$ and for each $n \geq 0$ the Steenrod square Sq^i is a homomorphism $\text{Sq}^i: H^n(X, A; \mathbb{Z}/2) \rightarrow H^{n+i}(X, A; \mathbb{Z}/2)$. The Steenrod squares have the following properties, which we will use; see e.g. [Ste62].

- (1) The Sq^i are natural with respect to maps of pairs $f: (X, A) \rightarrow (Y, B)$, for all $i \geq 0$.
- (2) $\text{Sq}^0 = \text{Id}$;

- (3) $\text{Sq}^i(x) = 0$ for $x \in H^n(X, A; \mathbb{Z}/2)$ with $n < i$;
- (4) $\text{Sq}^i(x) = x \cup x$ for $x \in H^i(X, A; \mathbb{Z}/2)$.

Let $1 \in H^0(B; \mathbb{Z}/2)$ denote the unit of the cohomology ring $H^*(B; \mathbb{Z}/2)$.

Definition 7.2 (Stiefel-Whitney classes [Fad65, Definition 6.1], [Ste75, p. 262]). The i th Stiefel-Whitney class of ξ is

$$w_i(\xi) := \Phi^{-1} \circ \text{Sq}^i \circ \Phi(1) \in H^i(B; \mathbb{Z}/2).$$

The definition uses the sequence of maps

$$H^0(B; \mathbb{Z}/2) \xrightarrow{\Phi, \cong} H^n(E, E_0; \mathbb{Z}/2) \xrightarrow{\text{Sq}^i} H^{n+i}(E, E_0; \mathbb{Z}/2) \xrightarrow{\Phi^{-1}, \cong} H^i(B; \mathbb{Z}/2).$$

Since $\text{Sq}^j: H^n(E, E_0; \mathbb{Z}/2) \rightarrow H^{n+j}(E, E_0; \mathbb{Z}/2)$ is the zero map for $j > n$, by the third property of Steenrod squares, it follows that $w_j(\xi) = 0$ for $j > n$.

We now restrict ourselves to bundles over spaces that are homotopy equivalent to CW complexes. In this context it was shown by [Fad65] and Stern [Ste75, Theorem 2.0] that the Stiefel-Whitney classes $w_i(\xi)$ for $0 \leq i \leq n$ satisfy the following properties:

Proposition 7.3. *Let B be a space homotopy equivalent to a CW complex and let $\xi = (p: E \rightarrow B)$ be a CAT \mathbb{R}^n -bundle over B . Define the total Stiefel-Whitney class*

$$w(\xi) := \sum_{i=0}^n w_i(\xi) \in H^*(B; \mathbb{Z}/2).$$

- (1) *For a CAT \mathbb{R}^n -bundle map $f = (f_E, f_B): \xi \rightarrow \eta$, which consists of maps*

$$\begin{array}{ccc} E(\xi) & \xrightarrow{f_E} & E(\eta) \\ \downarrow p(\xi) & & \downarrow p(\eta) \\ B(\xi) & \xrightarrow{f_B} & B(\eta), \end{array}$$

we have that $f_B^(w(\eta)) = w(\xi)$.*

- (2) *If $\xi = \eta^q \oplus \varepsilon^{n-q}$, where η is a CAT \mathbb{R}^q -bundle over B and ε^{n-q} is a trivial CAT \mathbb{R}^{n-q} -bundle over B , then $w(\xi) = w(\eta)$.*
- (3) *For each n there is a CAT \mathbb{R}^n -bundle such that $w_n(\xi) \neq 0$.*

Under the assumption that either $\text{CAT} \in \{\text{PL}, \text{Diff}\}$ or $\text{CAT} = \text{TOP}$ and $q \neq 4, 5$, Stern also proved that the properties in Proposition 7.3 characterise the Stiefel-Whitney classes. We are not sure whether the assumption that $q \neq 4, 5$ can now be removed in the TOP case using Quinn's work.

Recall the universal CAT \mathbb{R}^n -bundle γ_n^{CAT} over $\text{BCAT}(n)$ from Propositions 6.7 and 6.13. Here we are abusing notation, and using the deformation retract $\text{Diff}(n) \simeq \text{O}(n)$ (Remark 6.6) to conflate these structure groups. We denote the *universal Stiefel-Whitney classes* by

$$\overline{w}_k^{\text{CAT}} := w_k(\gamma_n^{\text{CAT}}) \in H^k(\text{BCAT}(n); \mathbb{Z}/2),$$

for some $n \geq k$. We also write $\bar{w}_k^{\text{CAT}}: \text{BCAT}(n) \rightarrow K(\mathbb{Z}/2, k)$ for the corresponding map to the Eilenberg-MacLane space. For $F: B \rightarrow \text{BCAT}(n)$, classifying a CAT \mathbb{R}^n -bundle ξ , we have by Proposition 7.3 that

$$w_k(\xi) = F^*(\bar{w}_k^{\text{CAT}}) \in H^k(B; \mathbb{Z}/2).$$

The definitions for each value of CAT are compatible in the following sense.

Proposition 7.4. *Let ξ be a CAT \mathbb{R}^n -bundle over B , for some $\text{CAT} \in \{\text{Diff}, \text{PL}\}$. Let ξ_{TOP} be the underlying TOP \mathbb{R}^n -bundle. Then $w_k(\xi) = w_k(\xi_{\text{TOP}})$ for every $k \geq 0$. In fact, the diagram*

$$\begin{array}{ccc} B & \xrightarrow{\xi} & \text{BCAT} \\ \xi_{\text{TOP}} \downarrow & \swarrow & \downarrow \bar{w}_k^{\text{CAT}} \\ \text{BTOP} & \xrightarrow{\bar{w}_k^{\text{TOP}}} & K(\mathbb{Z}/2, k). \end{array}$$

commutes up to homotopy.

PROOF. This is a consequence of the fact that the definitions in all three cases are directly analogous; cf. [Tho52, Théorème III.8] and [Fad65, Theorem 6.10], which consider the case of tangent bundles. \square

From now on we will often take the previous proposition to heart, and omit the CAT superscript, writing \bar{w}_k instead of \bar{w}_k^{CAT} . For the rest of this chapter this will in any case only be used with $\text{CAT} = \text{TOP}$.

Recall that $\text{TOP}(n)/\text{O}(n)$ is by definition the homotopy fibre of $\text{BO}(n) \rightarrow \text{BTOP}(n)$.

Proposition 7.5. *$\text{TOP}(n)/\text{O}(n)$ is contractible for $n \leq 3$, while for $n \geq 4$ there is a 5-connected map $\text{TOP}(n)/\text{O}(n) \rightarrow K(\mathbb{Z}/2, 3)$.*

PROOF. According to [KS77, Essay V, Section 5.0, p. 246], the homotopy fibre $\text{TOP}(2)/\text{O}(2)$ is contractible, $\pi_i(\text{TOP}(3)/\text{O}(3)) = 0$ for $i \geq 4$, and $\pi_i(\text{TOP}(3)/\text{O}(3)) \cong \pi_i(\text{Diff}(D^3, \partial D^3))$ for $i \geq 4$. The latter group is trivial for all i , by Hatcher's theorem [Hat83]. Thus all the homotopy groups of $\text{TOP}(3)/\text{O}(3)$ vanish, and therefore $\text{BO}(3) \rightarrow \text{BTOP}(3)$ induces an isomorphism on homotopy groups $\pi_i(\text{BO}(3)) \rightarrow \pi_i(\text{BTOP}(3))$ for all i . Since $\text{BO}(3) \rightarrow \text{BTOP}(3)$ is a map between spaces homotopy equivalent to a CW complex, we deduce that this map is a homotopy equivalence by Whitehead's theorem. Thus $\text{TOP}(3)/\text{O}(3)$ is the homotopy fibre of a homotopy equivalence and so is contractible.

The reference [KS77, Essay V, Section 5.0, p. 246] also includes the statement that for $n \geq 5$ and $i \leq 7$ we have $\pi_i(\text{TOP}/\text{O}, \text{TOP}(n)/\text{O}(n)) = 0$. In addition $\pi_i(\text{TOP}/\text{O}) \cong \pi_i(K(\mathbb{Z}/2, 3))$ for $i \leq 6$, with the isomorphism induced by the map $\text{TOP}/\text{O} \rightarrow \text{TOP}/\text{PL} \simeq K(\mathbb{Z}/2, 3)$, which is therefore a 5-connected map. The homotopy equivalence $\text{TOP}/\text{PL} \simeq K(\mathbb{Z}/2, 3)$ is from [KS77, Essay IV, Section 10.12, p. 200]. Since the composition of two 5-connected maps is 5-connected, it follows that there is a 5-connected map $\text{TOP}(n)/\text{O}(n) \rightarrow K(\mathbb{Z}/2, 3)$ for $n \geq 5$, as claimed.

It remains to consider $n = 4$. For this we appeal to [FQ90, Theorem 8.7A], which states that $\text{TOP}(4)/\text{O}(4) \rightarrow \text{TOP}/\text{O}$ is 5-connected. Combined with the fact already discussed that there is a 5-connected map $\text{TOP}/\text{O} \rightarrow K(\mathbb{Z}/2, 3)$, we obtain the sought-for 5-connected map $\text{TOP}(4)/\text{O}(4) \rightarrow K(\mathbb{Z}/2, 3)$. \square

Proposition 7.6.

- (1) For $n \geq 2$, $\pi_1(\text{BTOP}(n)) = \mathbb{Z}/2$.
- (2) The corresponding unique homotopically nontrivial map $\text{BTOP}(n) \rightarrow K(\mathbb{Z}/2, 1)$ is the universal first Stiefel-Whitney class \bar{w}_1 .
- (3) We have that $\pi_2(\text{BTOP}(2)) \cong \mathbb{Z}$.
- (4) For $n \geq 3$, $\pi_2(\text{BTOP}(n)) \cong \mathbb{Z}/2$.
- (5) The corresponding unique homotopically nontrivial map $\text{BTOP}(n) \rightarrow K(\mathbb{Z}/2, 2)$ is the universal second Stiefel-Whitney class \bar{w}_2 .

PROOF. For $n \leq 3$ we have $\text{TOP}(n) \simeq \text{O}(n)$ by Proposition 7.5, and so the homotopy groups of $\text{TOP}(n)$ are isomorphic to those of $\text{O}(n)$. For $n \geq 4$ we have the long exact sequence in homotopy groups:

$$\begin{aligned} \pi_2(\text{TOP}(n)/\text{O}(n)) &\rightarrow \pi_2(\text{BO}(n)) \rightarrow \pi_2(\text{BTOP}(n)) \rightarrow \\ \pi_1(\text{TOP}(n)/\text{O}(n)) &\rightarrow \pi_1(\text{BO}(n)) \rightarrow \pi_1(\text{BTOP}(n)) \rightarrow \{*\}. \end{aligned}$$

Since $\pi_i(\text{TOP}(n)/\text{O}(n)) = 0$ for $i = 1, 2$ by Proposition 7.5, we deduce that $\pi_i(\text{BO}(n)) \cong \pi_i(\text{BTOP}(n))$ for $i = 1, 2$, with the map induced by the canonical forgetful map. Since $\pi_i(\text{BO}(n)) \cong \pi_{i-1}(\text{O}(n)) \cong \mathbb{Z}/2$ for $i = 1, 2$, the result follows from this and Proposition 7.4. \square

We see that $\text{TOP}(n)$ has two connected components, which are homotopy equivalent because $\text{TOP}(n)$ is a topological group, and $\pi_1(\text{TOP}(n), \text{Id}) \cong \mathbb{Z}/2$ for $n \geq 3$.

Definition 7.7. We define $\text{STOP}(n)$ to be the subgroup of $\text{TOP}(n)$ consisting of orientation preserving homeomorphisms. We define $\text{TOPSpin}(n)$ to be the universal cover of $\text{STOP}(n)$. Define STOP and TOPSpin as corresponding colimits.

This definition is analogous to the definition of $\text{SO}(n)$ as the subgroup of $\text{O}(n)$ of orientation preserving orthogonal matrices, and of $\text{Spin}(n)$ as the connected double cover of $\text{SO}(n)$; this is the universal cover for $n \geq 3$.

Theorem 7.8. *The topological group $\text{STOP}(n)$ is the connected component of $\text{TOP}(n)$ containing the identity.*

PROOF. We saw above that $\text{TOP}(n)$ has two connected components. For a homeomorphism $f: \mathbb{R}^n \rightarrow \mathbb{R}^n$, and another such homeomorphism g , if f and g are isotopic, then f and g are either both orientation preserving (o.p.) or both orientation reversing (o.r.). The map $\pi_0(\text{TOP}(n)) \rightarrow \{\text{o.p.}, \text{o.r.}\}$ is a surjective map from a set with two elements to another set with two elements, hence is a bijection.

(Alternatively, the theorem can be seen as a consequence of the Stable Homeomorphism Theorem 4.3, which says that every orientation-preserving homeomorphism of \mathbb{R}^n is stable. Using that every homeomorphism of \mathbb{R}^n that is the identity on some subset is isotopic to the identity, via the (inverted) Alexander trick, as in Corollary 4.5, we deduce the result. The computations of the homotopy type of $\text{TOP}(n)/\text{O}(n)$ also used the Stable Homeomorphism Theorem, so this alternative proof is not independent of the first.) \square

We have the following commutative diagram of classifying spaces, together with the universal Stiefel-Whitney classes.

$$\begin{array}{ccccc}
 \text{BSpin}(n) & \longrightarrow & \text{BTOPSpin}(n) & & \\
 \downarrow & & \downarrow & & \\
 \text{BSO}(n) & \longrightarrow & \text{BSTOP}(n) & \xrightarrow{\bar{w}_2} & K(\mathbb{Z}/2, 2) \\
 \downarrow & & \downarrow & & \\
 \text{BO}(n) & \longrightarrow & \text{BTOP}(n) & \xrightarrow{\bar{w}_1} & K(\mathbb{Z}/2, 1)
 \end{array}$$

The horizontal maps in the bottom row induce isomorphisms on π_1 , while the horizontal maps in the middle row induce isomorphisms on π_2 . Thus up to homotopy equivalence, $\text{BSTOP}(n)$ is the 1-connected cover of $\text{BTOP}(n)$, and $\text{BTOPSpin}(n)$ is the 2-connected cover. In other words, $\text{BSTOP}(n)$ is the homotopy fibre of \bar{w}_1 and $\text{BTOPSpin}(n)$ is the homotopy fibre of \bar{w}_2 .

The analogous statements hold in the case of $\text{O}(n)$.

Definition 7.9. Let M be a space homotopy equivalent to a CW complex and let ξ be a TOP \mathbb{R}^n -bundle over M classified by a map which we also denote $\xi: M \rightarrow \text{BTOP}(n)$. An *orientation* on ξ is a lift $M \rightarrow \text{BSTOP}(n)$ of ξ , and two orientations are equivalent if the lifts are homotopic over $\text{BSTOP}(n) \rightarrow \text{BTOP}(n)$. If an orientation exists then we say ξ is *orientable*.

Proposition 7.10. *An orientation for an n -manifold M is equivalent to an orientation on the topological tangent bundle $\tau_M: M \rightarrow \text{BTOP}(n)$. In particular M is orientable if and only if τ_M is orientable.*

PROOF. An orientation for M is equivalent to a homology orientation for M , i.e. a coherent choice of generators of $H_n(M, M \setminus \{x\}; \mathbb{Z})$, for $x \in M$. In turn, a homology orientation for M is equivalent to a coherent system of orientations of the fibre of the tangent microbundle $M \xrightarrow{\Delta} M \times M \xrightarrow{\text{pr}_1} M$, rel. the zero section $\Delta(M)$. This is because both are, by definition, a coherent system of generators of $H_n(M, M \setminus \{x\}; \mathbb{Z})$, for each $x \in M$. See [MS74, Lemma 11.6] for details.

Next, the latter notion is equivalent, via Kister's theorem [Kis64] and excision, to a coherent choice of generators of $H_n(F_x, F_x \setminus \{0\}; \mathbb{Z}) \cong \mathbb{Z}$. Here, F_x is the fibre over x in the topological tangent \mathbb{R}^n -bundle of M ; see [MS74, Lemma 11.7], but replace the exponential map with the embedding from Kister's theorem.

To see that a coherent choice of isomorphisms $H_n(F_x, F_x \setminus \{0\}; \mathbb{Z}) \cong \mathbb{Z}$ is equivalent to a choice of lift $M \rightarrow \text{BSTOP}(n)$, recall that the two connected components of $\text{TOP}(n)$ correspond to whether a homeomorphism preserves or changes a fixed generator of $H_n(\mathbb{R}^n, \mathbb{R}^n \setminus \{0\}; \mathbb{Z})$, so the structure group reduces to $\text{STOP}(n)$ if and only if there is a coherent choice of generators for the homology groups $H_n(F_x, F_x \setminus \{0\}; \mathbb{Z})$, for $x \in M$. \square

Definition 7.11. For an n -manifold M , with topological tangent bundle τ_M , we define $w_i(M) := w_i(\tau_M)$.

Proposition 7.12. *Let M be a space homotopy equivalent to a CW complex and let ξ be a TOP \mathbb{R}^n -bundle over M . The bundle ξ is orientable if and only if $w_1(\xi) = 0$. In particular, by Proposition 7.10, a manifold M is orientable if and only if $w_1(M) = 0$.*

PROOF. Since $\text{BSTOP}(n) \rightarrow \text{BTOP}(n) \xrightarrow{\bar{w}_1} K(\mathbb{Z}/2, 1)$ is a fibration sequence, we have an exact sequence of pointed sets

$$[M, \text{BSTOP}(n)] \rightarrow [M, \text{BTOP}(n)] \rightarrow [M, K(\mathbb{Z}/2, 1)] \cong H^1(M; \mathbb{Z}/2).$$

The bundle ξ is orientable if and only if the classifying map $\xi \in [M, \text{BTOP}(n)]$ is homotopic to a map in the image of $[M, \text{BSTOP}(n)]$. The latter is equivalent to $\bar{w}_1 \circ \xi \in [M, K(\mathbb{Z}/2, 1)]$ being null-homotopic, using the sequence. Translating to cohomology groups this is equivalent to $w_1(\xi) = \xi^*(\bar{w}_1) = 0$. Here we used that $w_1(\xi)$ is equal to the pullback of the universal bundle along the classifying map for ξ . \square

Definition 7.13. Let M be a space homotopy equivalent to a CW complex and let ξ be a TOP \mathbb{R}^n -bundle over M classified by a map which we also denote $\xi: M \rightarrow \text{BTOP}(n)$. Suppose that $w_1(M) = 0$, so ξ is orientable. A *spin structure* on ξ is a lift $M \rightarrow \text{BTOPSpin}(n)$ of ξ , and two spin structures are equivalent if the lifts are homotopy equivalent. If a spin structure exists then we say ξ is *spin*.

Now suppose that M is a topological n -manifold with topological tangent bundle classified by $\tau_M: M \rightarrow \text{BTOP}(n)$. Suppose that $w_1(M) = 0$, so M is orientable. A *spin structure* on M is a spin structure on τ_M . If a spin structure on τ_M exists then we say M is *spin*.

Proposition 7.14. *Let M be a space homotopy equivalent to a CW complex and let ξ be a TOP \mathbb{R}^n -bundle over M . Suppose that ξ is orientable. Then ξ is spin if and only if $w_2(\xi) = 0$. In particular, an orientable manifold M is spin if and only if $w_2(M) = 0$.*

PROOF. This holds because $\text{BTOPSpin}(n) \rightarrow \text{BSTOP}(n) \xrightarrow{\bar{w}_2} K(\mathbb{Z}/2, 2)$ is a fibration sequence. Thus a lift of $\xi: M \rightarrow \text{BSTOP}(n)$ to $\text{BTOPSpin}(n)$ exists if and only if $\bar{w}_2 \circ \xi$ is null-homotopic, i.e. if and only if $w_2(\xi) = 0$. Here we are again using that a fibration sequence gives rise to an exact sequence of sets

$$[M, \text{BTOPSpin}(n)] \rightarrow [M, \text{BSTOP}(n)] \rightarrow [M, K(\mathbb{Z}/2, 2)] \cong H^2(M; \mathbb{Z}/2),$$

and hence a lift to $\text{BTOPSpin}(n)$ exists if and only if $w_2(\xi) = \xi^*(\bar{w}_2) = 0$. \square

We now recall the Wu classes and the Wu formulae for the Stiefel-Whitney classes of compact manifolds. The key point here is that while the treatment in [MS74, Chapter 11] is

presented for smooth manifolds, in fact it uses only Poincaré duality and algebraic topology, so works just as well for topological manifolds. This has been observed by Thom [Tho52] and Fadell [Fad65]. We proceed to summarise the treatment in Milnor-Stasheff.

Let M be a compact n -manifold with $\mathbb{Z}/2$ -fundamental class $[M, \partial M] \in H_n(M, \partial M; \mathbb{Z}/2)$. Consider the homomorphism

$$\begin{aligned} \theta: H^{n-k}(M, \partial M; \mathbb{Z}/2) &\rightarrow \mathbb{Z}/2 \\ x &\mapsto \langle \text{Sq}^k(x), [M, \partial M] \rangle. \end{aligned}$$

By Poincaré duality, there is a unique class $v_k(M) \in H^k(M; \mathbb{Z}/2)$ with

$$\langle v_k(M) \cup x, [M, \partial M] \rangle = \theta(x)$$

for all $x \in H^{n-k}(M, \partial M; \mathbb{Z}/2)$. In fact $v_k(M) \cup x = \text{Sq}^k(x) \in H^n(M, \partial M; \mathbb{Z}/2)$ for every $x \in H^{n-k}(M, \partial M; \mathbb{Z}/2)$.

Definition 7.15. The class $v_k(M)$ is the k th Wu class of M .

Proposition 7.16. Let M be a compact n -manifold. Then the Wu formulae

$$w_k(M) = \sum_{i+j=k} \text{Sq}^i(v_j)$$

hold for $k = 0, \dots, n$.

PROOF. The proof in [MS74, Theorem 11.11, Lemma 11.13, and Theorem 11.14], which relies on [MS74, Theorem 11.11 and Lemma 11.13], uses only Poincaré duality and products from algebraic topology, and so proceeds exactly as in Milnor-Stasheff. \square

Here is a sample, and often used, application for the second Stiefel-Whitney class.

Definition 7.17. Let M be a compact, oriented n -manifold and $\Sigma^{n-2} \subseteq M$ a proper submanifold. We say Σ is *characteristic* if $\text{PD}(w_2(M)) = j_*([\Sigma]) \in H_{n-2}(M, \partial M; \mathbb{Z}/2)$, where $j_*: H_{n-2}(\Sigma, \partial\Sigma; \mathbb{Z}/2) \rightarrow H_{n-2}(M, \partial M; \mathbb{Z}/2)$ is the inclusion-induced map.

Proposition 7.18. Let M be a compact n -manifold and $\Sigma \subseteq M$ a proper submanifold. Then $w_2(M \setminus \Sigma) = 0$ if and only if Σ is characteristic.

PROOF. Throughout the proof, $\mathbb{Z}/2$ -coefficients are understood. By the Collar Neighbourhood Theorem (2.16), we may take a boundary collar on M that restricts on Σ to a boundary collar on Σ . Set $K := M \setminus (\partial M \times [0, 1))$, the complement of the open collar on ∂M ; note K is compact. Write $\nu\Sigma$ for an open tubular neighbourhood of Σ . Tubular neighbourhoods were proved to exist in codimension 2 when $n \neq 4$ by Kirby-Siebenmann [KS75] and for $n = 4$ by Freedman-Quinn (Theorem 5.16). Next we draw a diagram, then define the maps and justify that it has exact rows, is commutative, and that the variously claimed

isomorphisms indeed are so.

$$\begin{array}{ccccc}
H^2(M, M \setminus \nu\Sigma) & \xrightarrow{\beta} & H^2(M) & \xrightarrow{\alpha} & H^2(M \setminus \nu\Sigma) \\
\cong \downarrow & & \cong \downarrow & & \cong \downarrow \\
H^2(K, K \setminus \nu\Sigma) & \xrightarrow{\beta'} & H^2(K) & \xrightarrow{\alpha'} & H^2(K \setminus \nu\Sigma) \\
\cong \downarrow -\cap \mathcal{O} & & \cong \downarrow -\cap \mathcal{O} & & \\
H_{n-2}(\partial M \cup \nu\Sigma, \partial M) & \xrightarrow{\text{incl}} & H_{n-2}(M, \partial M) & & \\
\cong \uparrow \text{incl} & & \uparrow j_* & & \\
H_{n-2}(\nu\Sigma, \partial M \cap \nu\Sigma) & \xleftarrow[\text{incl}]{\cong} & H_{n-2}(\Sigma, \partial\Sigma) & &
\end{array}$$

The top row is a section of the long exact sequence of the pair $(M, M \setminus \nu\Sigma)$, and the central row is a section of the long exact sequence of the pair $(K, K \setminus \nu\Sigma)$, thus both are exact. The downwards maps from the top row to the middle row are by definition the maps induced by inclusion, which is a homotopy equivalence, justifying these isomorphisms and the fact that the subdiagram consisting of the top two rows commutes.

The class \mathcal{O} is the $\mathbb{Z}/2$ -orientation class; see [Bre97, §VI.8]. Replacing M by $M \setminus \partial M$, we obtain a (non-compact) 4-manifold M' with empty boundary. By [Bre97, §VI Theorem 8.3], capping with \mathcal{O} induces isomorphisms

$$H^2(K, L) \xrightarrow{\cong} H_{n-2}(M' \setminus L, M' \setminus K),$$

both for $L := K \setminus \nu\Sigma$ and for $L := \emptyset$. Thus, we obtain isomorphisms

$$H^2(K, K \setminus \nu\Sigma) \xrightarrow{\cong} H_{n-2}(\partial M \times (0, 1) \cup \nu\Sigma, \partial M \times (0, 1)) \cong H_{n-2}(\partial M \cup \nu\Sigma, \partial M)$$

and

$$H^2(K) \xrightarrow{\cong} H_{n-2}(M', \partial M \times (0, 1)) \cong H_{n-2}(M, \partial M).$$

These define the middle vertical maps labelled $-\cap \mathcal{O}$. By naturality of the cap product, the left middle square commutes. The commutativity of the bottom left square is clear, as is the isomorphism of the left-bottom arrow. We also note that the composite of the top two central downwards arrows is the Poincaré-Lefschetz duality isomorphism $PD: H^2(M) \xrightarrow{\cong} H_{n-2}(M, \partial M)$. Having established the relevant properties of the diagram, we now prove the lemma.

First note that by naturality of Stiefel-Whitney classes, we have $\alpha(w_2(M)) = w_2(M \setminus \nu\Sigma)$. Note as well that $H_{n-2}(\partial M \cup \nu\Sigma, \partial M; \mathbb{Z}/2)$ is isomorphic to $\mathbb{Z}/2$ and generated by the image of $[\Sigma]$.

For one direction of the lemma, assume that $w_2(M \setminus \Sigma) = 0$. By exactness of the top row, this implies $w_2(M)$ lies in the image of β . But as $H^2(M, M \setminus \nu\Sigma) \cong \mathbb{Z}/2$ and $w_2(M) \neq 0$, this implies the generator of $\mathbb{Z}/2$ is sent to $w_2(M)$ by β . By commutativity of the diagram, we get $PD(w_2(M)) \equiv j_*([\Sigma]) \in H_{n-2}(M, \partial M; \mathbb{Z}/2)$.

Conversely, assume that $\text{PD}(w_2(M)) \equiv j_*([\Sigma]) \in H_{n-2}(M, \partial M; \mathbb{Z}/2)$. By commutativity of the diagram, this implies $w_2(M)$ is in the image of β . Thus $w_2(M \setminus \nu\Sigma) = \alpha(w_2(M)) = 0$ by exactness of the top row. \square

Remark 7.19. We note that the previous proposition is valid when none of the manifolds involved is orientable and also Σ is closed. We also note that the proof given above is fairly robust and could be easily adapted to other Stiefel-Whitney classes, provided the tubular neighbourhood conditions are met. For example, the same idea shows that the complement of a codimension 1 submanifold is orientable if and only if that submanifold is Poincaré dual to the first Stiefel-Whitney class.

CHAPTER 8

Intersection forms and smooth 4-manifolds

In this chapter we introduce and study one of the most interesting invariants of 4-manifolds, namely the intersection form. Later, in Theorem 12.2 we will see that any unimodular symmetric form over \mathbb{Z} occurs as the intersection form of a closed oriented 4-manifold. In contrast we will see in this chapter that not all unimodular symmetric form over \mathbb{Z} can be realised as the intersection forms of closed oriented *smooth* 4-manifolds.

8.1. INTERSECTION FORMS

We start out with the definition of the intersection form.

Definition 8.1.

- (1) Given a finitely generated abelian group H we write $FH := H/\text{torsion}$ subgroup.
- (2) Given a compact oriented n -manifold M we denote by $[M] \in H_n(M, \partial M; \mathbb{Z})$ its fundamental class. Given a decomposition $\partial M = A \cup B$ where A and B are compact codimension-zero submanifolds of ∂M with $A \cap B = \partial A = \partial B$ we denote the Poincaré duality isomorphism by

$$\begin{aligned} \text{PD}: H^l(M, A; \mathbb{Z}) &\rightarrow H_{n-l}(M, B; \mathbb{Z}) \\ \phi &\mapsto \varphi \cap [M]. \end{aligned}$$

- (3) Given a compact oriented 4-manifold M we refer to the map

$$\begin{aligned} Q_M: FH_2(M; \mathbb{Z}) \times FH_2(M; \mathbb{Z}) &\rightarrow \mathbb{Z} \\ (a, b) &\mapsto Q_M(a, b) := \langle \text{PD}_M^{-1}(a) \cup \text{PD}_M^{-1}(b), [M] \rangle \end{aligned}$$

as the *intersection form*. (Here $\langle -, - \rangle$ denotes the Kronecker pairing.) Using Poincaré Duality one can easily show that if M is closed, then Q_M is nonsingular.

Let E_8 denote the even 8×8 Cartan matrix of the eponymous exceptional Lie algebra; that is,

$$E_8 = \begin{pmatrix} 2 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 2 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 2 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 2 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 2 & 1 & 0 & 1 \\ 0 & 0 & 0 & 0 & 1 & 2 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 2 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 2 \end{pmatrix}.$$

Note that this is a symmetric integral matrix with determinant one.

Example 8.2. Here are some important closed, smooth 4-manifolds.

- (1) The 4-sphere S^4 . This is simply connected and has $H_2(S^4; \mathbb{Z}) = \{0\}$.
- (2) The complex projective plane $\mathbb{C}P^2$, which comes with a canonical orientation. The same underlying manifold with the opposite orientation is $\overline{\mathbb{C}P^2}$. They are simply-connected manifolds with $H_2(\mathbb{C}P^2; \mathbb{Z}) \cong \mathbb{Z}$. The intersection form of $\mathbb{C}P^2$ is represented by the 1×1 -matrix (1) and the intersection form of $\overline{\mathbb{C}P^2}$ is represented by the 1×1 -matrix (-1) .
- (3) The manifold $S^2 \times S^2$ is simply-connected and $H_2(S^2 \times S^2; \mathbb{Z}) \cong \mathbb{Z} \oplus \mathbb{Z}$. The intersection form of $S^2 \times S^2$ is represented by the standard hyperbolic form $H := \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$.
- (4) The $K3$ surface or Kummer surface

$$K3 := \{[z_1 : z_2 : z_3 : z_4] \in \mathbb{C}P^3 \mid z_1^4 + z_2^4 + z_3^4 + z_4^4 = 0\}$$

This is a simply connected, smooth, spin, closed 4-manifold with $H_2(K3; \mathbb{Z}) \cong \mathbb{Z}^{22}$. As is shown in [GS99, Theorem 1.3.8] or alternatively [MS17, p. 176], the intersection form of $K3$ is isometric to $E_8 \oplus E_8 \oplus H \oplus H \oplus H$.

The next proposition shows that the intersection form is well behaved under the connected sum operation.

Proposition 8.3. *Let M and N be two oriented compact 4-manifolds. Then there is an isomorphism $H_2(M) \oplus H_2(N) \rightarrow H_2(M \# N)$ that induces an isometry of $Q_M \oplus Q_N$ and $Q_{M \# N}$.*

PROOF. The usual tools of algebraic topology, namely a Mayer-Vietoris argument and the excision theorem, show that there exists an isomorphism $\Theta: H_2(M) \oplus H_2(N) \rightarrow H_2(M \# N)$.

The statement that this isomorphism Θ induces an isometry between $Q_M \oplus Q_N$ and $Q_{M \# N}$ can be deduced from the functoriality of the cup and cap products [Bre97, Theorem VI.5.2.(4)] for maps between pairs of topological spaces. Full details are provided in [Fri23, Proposition 153.12].

In the smooth case the statement that the isomorphism Θ induces an isometry of forms follows immediately from the fact that any class in second homology can be represented by an embedded oriented submanifold [GS99, Proposition 1.2.3] and the fact that one can calculate the intersection form in terms of algebraic intersection numbers of embedded oriented surfaces [Bre97, Theorem VI.11.9]. To apply this approach to general manifolds, one needs to use topological transversality, which holds, as discussed in Chapter 10. \square

8.2. INTERSECTION FORMS OF SPIN MANIFOLDS

Using the results from the previous chapter we can prove the following proposition.

Proposition 8.4. *Let M be a compact, connected, oriented 4-manifold. If M is spin then the intersection form of M is even.*

PROOF. In this proof we use the properties of the Steenrod squares introduced just before Definition 7.2.

By definition the k th Wu class $v_k \in H^k(M; \mathbb{Z}/2)$ satisfies $\text{Sq}^k(a) = v_k \cup a$, for every class $a \in H^{4-k}(M, \partial M; \mathbb{Z}/2)$. Hence if $a \in H^2(M, \partial M; \mathbb{Z}/2)$ then $v_2 \cup a = \text{Sq}^2(a) = a \cup a$. By Proposition 7.16, the n th Stiefel-Whitney class of M is given by $w_n = \sum_i \text{Sq}^i(v_{n-i})$. Since M is oriented, we have

$$0 = w_1 = \text{Sq}^0(v_1) + \text{Sq}^1(v_0) = v_1.$$

Since M is spin, we have that

$$0 = w_2 = \text{Sq}^0(v_2) + \text{Sq}^1(v_1) + \text{Sq}^2(v_0) = v_2.$$

So for any $a \in H^2(M, \partial M; \mathbb{Z}/2)$, we have $a \cup a = 0 \cup a = 0 \in H^4(M, \partial M; \mathbb{Z}/2) = \mathbb{Z}/2$. But this implies that for any $x \in FH_2(M; \mathbb{Z})$ we have that $Q_M(x, x) = \langle \text{PD}^{-1}(x) \cup \text{PD}^{-1}(x), [M, \partial M] \rangle \equiv 0 \pmod{2}$. In other words Q_M is an even form. \square

Proposition 8.5. *Let M be a closed, oriented, connected, spin 4-manifold. Then the signature $\text{sign}(M)$ is divisible by 8.*

PROOF. In Proposition 8.4 we just proved that the intersection form Q_M is even. Since M is closed we know that that Q_M is nonsingular. Finally note that it is an algebraic fact, shown for example in [MH73, Theorem 5.1], that for any symmetric nonsingular, bilinear, even form Q , the signature of Q is divisible by 8. \square

8.3. TWISTED INTERSECTION FORMS AND TWISTED SIGNATURES

In this section we introduce twisted intersection forms for topological manifolds and discuss some properties of the corresponding twisted signatures.

Let M be a compact, orientable, connected $4m$ -dimensional manifold. We write $\pi := \pi_1(M)$. Let $\alpha: \pi \rightarrow U(k)$ be a unitary representation. We view the elements of \mathbb{C}^k as row vectors. Given $g \in \pi$ and $v \in \mathbb{C}^k$, define $v \cdot g := v \cdot \alpha(g)$. Thus we can view \mathbb{C}^k as a right $\mathbb{Z}[\pi]$ -module. Denote this module by \mathbb{C}_α^k . Define the *twisted intersection form* of (M, α) to be the form

$$\begin{aligned} Q_M: H_{2m}(M; \mathbb{C}_\alpha^k) \times H_{2m}(M; \mathbb{C}_\alpha^k) &\xrightarrow{\text{PD}^{-1} \times \text{PD}^{-1}} H^{2m}(M, \partial M; \mathbb{C}_\alpha^k) \times H^{2m}(M, \partial M; \mathbb{C}_\alpha^k) \\ &\downarrow \cup \\ &H^{4m}(M, \partial M; \mathbb{C}_\alpha^k \otimes \mathbb{C}_\alpha^k) \\ &\downarrow \langle, \rangle \\ &H^{4m}(M, \partial M; \mathbb{C}) \\ &\downarrow \text{PD} \\ &H_0(M; \mathbb{C}) = \mathbb{C}. \end{aligned}$$

Here the first and the last map are given by the isomorphisms from the Poincaré Duality Theorem A.15 and the second map is given by Lemma A.11. Note that in the bottom we view \mathbb{C} as a trivial $\mathbb{Z}[\pi]$ -module. The third map is induced by the following homomorphism of right $\mathbb{Z}[\pi]$ -modules:

$$\begin{aligned} \mathbb{C}_\alpha^k \otimes \mathbb{C}_\alpha^k &\rightarrow \mathbb{C} \\ (v, w) &\mapsto \langle v, w \rangle = \bar{v}w^T. \end{aligned}$$

It follows easily from the definitions that Q_M is sesquilinear, namely \mathbb{C} -conjugate linear in the first entry and \mathbb{C} -linear in the second entry. The usual proof for the (anti-) symmetry of the cup product e.g. [Hat02, Theorem 3.14], can be modified to show that Q_M is hermitian, that is for every $v, w \in H^{2m}(M; \mathbb{C}_\alpha^k)$ we have $Q_M(v, w) = \overline{Q_M(w, v)}$. Since Q_M is hermitian, its signature is defined as the difference in the number of positive and negative eigenvalues. We refer to the signature of Q_M as the *twisted signature* $\sigma(M, \alpha)$.

For a group homomorphism $\gamma: \pi_1(M) \rightarrow \Gamma$, denote the corresponding L^2 -signature by $\sigma^{(2)}(M, \gamma)$, as defined in say [Ati76, Lüc02] and [COT03, Chapter 5].

Theorem 8.6. *Let M be a closed, oriented, connected 4-manifold.*

- (1) *For every finite cover $p: \widetilde{M} \rightarrow M$ we have $\sigma(\widetilde{M}) = [\widetilde{M} : M] \cdot \sigma(M)$.*
- (2) *For every unitary representation $\alpha: \pi_1(M) \rightarrow U(k)$ we have $\sigma(M, \alpha) = k \cdot \sigma(M)$.*
- (3) *For every group homomorphism $\gamma: \pi_1(M) \rightarrow \Gamma$ we have $\sigma^{(2)}(M, \gamma) = \sigma(M)$.*

Remark 8.7.

- (1) The same statement does not hold for 4-dimensional Poincaré complexes in general. More precisely, Wall [Wal67, Corollary 5.4.1] gave examples of 4-dimensional Poincaré complexes for which the signature is not multiplicative under finite covers.
- (2) Alternative proofs for the first and the third statement are provided by Schafer [Sch70, Theorem 8] and Lück-Schick [LS03, Theorem 0.2]. The approach taken in Lück-Schick [LS03] and Teleman [Tel84] should also provide a proof of the second statement. In fact these papers are also valid for manifolds of any dimension $4m$.

PROOF. First we give references for these three statements for smooth manifolds.

- (1) This statement is a consequence of the Hirzebruch Signature Theorem (see e.g. [MS74]).
- (2) This statement was proven in [APS75] (in fact the second statement contains the first statement as a special case).
- (3) This statement was proven in [Ati76, p. 44].

We now turn to manifolds that are not necessarily smooth. We will prove the second statement of the theorem. The other statements can be proved in a similar fashion. We refer to [COT03, Lemma 5.9] for a proof of (3).

So let M be a closed oriented connected 4-manifold and let $\alpha: \pi_1(M) \rightarrow U(k)$ be a unitary representation. By Theorem 9.9, there exists a closed orientable simply-connected 4-manifold N such that $M \# N$ is smooth. We have $\pi_1(M \# N) = \pi_1(M) * \pi_1(N) \cong \pi_1(M)$ since $\pi_1(N) = \{1\}$. Let $\beta: \pi_1(N) \rightarrow U(k)$ be the trivial representation. We also write

$\alpha * \beta: \pi_1(M \# N) = \pi_1(M) \rightarrow U(k)$ for the representation uniquely determined by α on $\pi_1(M)$.

By Proposition 8.3, we have $\sigma(M \# N) = \sigma(M) + \sigma(N)$. Furthermore a slight generalisation of Proposition 8.3 shows that $\sigma(M \# N, \alpha * \beta) = \sigma(M, \alpha) + \sigma(N, \beta)$. Finally, we have $\sigma(N, \beta) = k \cdot \sigma(N)$. The desired statement follows from these equalities and from the formula for twisted signatures of the closed smooth manifold $M \# N$. \square

8.4. INTERSECTION FORMS OF SMOOTH 4-MANIFOLDS

In Theorem 12.2 we will see that any unimodular symmetric form occurs as the intersection form of a closed oriented 4-manifold. In the following we survey results on intersection forms of closed oriented smooth 4-manifolds. As we will see, the results in the smooth setting differ dramatically from the results in the topological setting.

In Proposition 8.5 we saw that the signature of any closed, oriented, connected, spin 4-manifolds is divisible by 8. The Rochlin Theorem [Roh52] gives an extra restriction on the signatures of intersection forms of spin 4-manifolds that admit a smooth structure.

Theorem 8.8. (Rochlin Theorem) *Let M be a closed, oriented, connected, spin, smooth 4-manifold. Then the signature $\text{sign}(M)$ is divisible by 16.*

Remark 8.9. Let M be a closed oriented 4-manifold with an even intersection form and such that $H_1(M; \mathbb{Z})$ has no 2-torsion. This implies $H^2(M; \mathbb{Z}/2) \cong \text{Hom}(H_2(M; \mathbb{Z}), \mathbb{Z}/2)$ and that the mod 2 reduction of Q_M is isomorphic to the pairing $(a, b) = \langle a \cup b, [M] \rangle$ on $H^2(M; \mathbb{Z}/2)$. As Q_M is even, this implies that $(a, a) = 0 \in \mathbb{Z}/2$ for any $a \in H^2(M; \mathbb{Z}/2)$. But we saw in the proof of Proposition 8.5 that $a \cup a = v_2 \cup a$, so we must have that $v_2 = 0$ as this pairing is nondegenerate. We also saw in the proof of Proposition 8.5 that $v_2 = w_2$ when M is oriented, so in fact $w_2 = 0$ and M admits a spin structure.

It is not true that simply having an even intersection form implies M is spin. Indeed, it is possible to construct a closed oriented 4-manifold M that has $Q_M = 0$ (which is in particular an even form), but has nonvanishing w_2 [GS99, Exercise 5.7.7(a)]. In a similar spirit, by [Hab82, FS84] there exists a closed oriented 4-dimensional smooth manifold M with an even intersection form Q_M that satisfies $\text{sign}(M) = 8$. Hence this must also fail to be spin, now by the Rochlin Theorem 8.8.

Theorem 8.10 (Freedman [Fre82]). *There exists a closed orientable connected 4-manifold that does not admit a smooth structure.*

PROOF. By Theorem 12.2 there exists a simply connected closed oriented 4-manifold M with $Q_M \cong E_8$. By the Rochlin Theorem 8.8 this manifold does not admit a smooth structure. \square

In a remarkable twist, shortly after Freedman proved Theorem 12.2, Donaldson [Don83, Theorem A] [Don87, Theorem 1], proved the following result regarding intersection forms of smooth 4-manifolds.

Theorem 8.11. (Donaldson’s Theorem) *Let M be a closed oriented connected smooth 4-manifold. If Q_M is positive-definite, then Q_M can be represented by the identity matrix.*

To understand the significance of Donaldson’s Theorem it is helpful to consider the following table from [MH73, p. 28], which basically says that there are lots of isometry types of nonsingular positive definite forms.

Dimension:	8	16	24	32	40
Number of isometry types of nonsingular positive definite even symmetric forms:	1	2	24	$\geq 10^7$	$\geq 10^{51}$

Remark 8.12. Note that if M be a closed oriented smooth 4-manifold such that Q_M is negative-definite, then $Q_{-M} = -Q_M$ is positive-definite. Thus we see that Donaldson’s Theorem implies that Q_M is represented by $-\text{Id}$.

It follows from [MH73, Theorem II.5.3] that every nonsingular indefinite *odd* symmetric form is isometric to $k \cdot (1) \oplus \ell \cdot (-1)$. These are realised by $k \cdot \mathbb{C}P^2 \# \ell \cdot \overline{\mathbb{C}P}^2$. Therefore we only need to discuss the realisability of nonsingular indefinite *even* symmetric forms. Again by [MH73, Theorem II.5.3], every nonsingular *even* indefinite symmetric form is isometric to $n \cdot E_8 \oplus m \cdot H$ for some $(m, n) \in \mathbb{N}_0 \times \mathbb{Z} \setminus \{(0, 0)\}$. The following theorem, proven by Furuta [Fur01], gives some restrictions on the possible values of m and n .

Theorem 8.13. (Furuta’s 10/8 Theorem) *If M is a closed oriented connected smooth 4-manifold with indefinite even intersection form, then*

$$b_2(M) \geq \frac{10}{8} \cdot |\text{sign}(M)| + 2.$$

In particular $Q_M \cong n \cdot E_8 \oplus m \cdot H$ for some $n \in 2\mathbb{Z}$ and $m \in \mathbb{N}$ with $m \geq |n| + 1$.

Furuta’s 10/8 Theorem does not quite close the gap between the forms we can realise by smooth manifolds and the forms we can exclude. More precisely, it follows from the calculation of the intersection form of the K3 surface and of $S^2 \times S^2$ that for any $n = 2p \in \mathbb{Z}$ and every $m \geq 3|p|$ there exists a closed oriented simply connected 4-dimensional smooth manifold with intersection form isometric to $n \cdot E_8 \oplus m \cdot H$. In other words, we have

$$\text{intersection form of } p \cdot \text{K3} \# (m - 3|p|) \cdot (S^2 \times S^2) \cong 2p \cdot E_8 \oplus m \cdot H.$$

The following conjecture predicts that this result is optimal.

Conjecture 8.14. (11/8-Conjecture) *If M is a closed oriented connected smooth 4-manifold with indefinite even intersection form, then*

$$b_2(M) \geq \frac{11}{8} \cdot |\text{sign}(M)|.$$

Equivalently, if $Q_M \cong 2p \cdot E_8 \oplus m \cdot H$ with $p \neq 0$, then $m \geq 3|p|$.

Remark 8.15.

- (1) A proof of the 11/8-Conjecture would imply, by Freedman's Theorem 12.2, that any closed oriented simply connected smooth 4-manifold is *homeomorphic* to either a connected sum of the form $k \cdot \mathbb{C}P^2 \# \ell \cdot \overline{\mathbb{C}P}^2$ or to a connected sum of the form $n \cdot K3 \# m \cdot (S^2 \times S^2)$.
- (2) Currently the best known result in the direction of the 11/8-Conjecture is [HLSX22, Corollary 1.13], which says that if M is a closed oriented simply-connected 4-manifold that is not homeomorphic to S^4 , $S^2 \times S^2$ or the K3 surface and whose intersection form is indefinite and even, then $b_2(M) \geq \frac{10}{8} \cdot |\text{sign}(M)| + 4$.

CHAPTER 9

Smoothing 4-manifolds

In this chapter we present three theorems which associate a smooth manifold to a given 4-manifold. Often these theorems can be used to reduce proofs about 4-manifolds to the case of smooth 4-manifolds, where the standard tools of differential topology are available.

9.1. SMOOTHING NONCOMPACT 4-MANIFOLDS

The first of our smoothing theorems [Qui82, Corollary 2.2.3], [FQ90, p. 116], which is due to Freedman and Quinn, says that noncompact connected 4-manifolds admit a smooth structure.

Theorem 9.1. *Every connected, noncompact 4-manifold is smoothable. Thus every 4-manifold M has a smooth structure in the complement of any closed set that has at least one point in each compact component of M .*

There are some related statements in the literature on smoothing 4-manifolds in the complement of a point, that appeared prior to Freedman's work [Fre82] and prior to [Qui82]. We discuss them briefly here. For the case of PL structures on noncompact 4-manifolds, given a lift of the (unstable) tangent microbundle classifying map $M \rightarrow \text{BTOP}(4)$ to $M \rightarrow \text{BPL}(4)$ (see Chapter 6), the result can be found in [Las70, p. 54] and [KS77, Essay V, Addendum 1.4.1, p. 222]. The analogous result for smooth bundle structures and smooth structures on noncompact manifolds was stated in [Las71, p. 156]. Alternatively, [HM74], [FQ90, Theorem 8.3B] apply to improve a PL structure to a smooth structure, unique up to isotopy, for any manifold of dimension at most six. Again, in [Las71] Lashof assumes a lift of the (unstable) tangent microbundle classifying map $M \rightarrow \text{BTOP}(4)$ to a map $M \rightarrow \text{BO}(4)$. For noncompact connected 4-manifolds, such a lift always exists, as was later shown by Quinn [Qui82, Qui84], [FQ90, p. 116] using the full disc embedding theorem [Fre82], and giving rise to Theorem 9.1.

Due to the seminal nature of Freedman's Fields medal winning paper [Fre82], it is well worth clarifying the details of some citations therein. In the proof of Corollary 1.2, in the proof of Theorem 1.5 on page 369, in the proof of Theorem 1.6, and at the start of Section 10, Freedman uses that smoothing theory is available for noncompact 4-manifolds. In particular, smoothing for noncompact contractible 4-manifolds plays a vital rôle in Freedman's proof of the topological 4-dimensional Poincaré conjecture [Fre82, Theorem 1.6]. Freedman cites [KS77] for this fact, however [KS77, Essay V, Remarks 1.6 (A), p. 230] specifically excludes smooth structures (but for a stronger result). Nevertheless, as mentioned above,

Lashof [Las71, p. 156] proved the smooth version of [KS77, Essay V, Addendum 1.4.1, p. 222], or one can use PL smoothing theory [HM74], [FQ90, Theorem 8.3B] to improve a PL structure from [KS77, Essay V, Addendum 1.4.1, p. 222] to a smooth structure, essentially uniquely.

Freedman only applies smoothing theory in cases, such as for contractible M , that he can ensure the existence of a lift of $\tau_M: M \rightarrow \text{BTOP}(4)$ to $\text{BO}(4)$. Later, Quinn [Qui82, Corollary 2.2.3] showed that such a lift always exists for connected noncompact 4-manifolds. In fact, he showed that the map $\text{TOP}(4)/\text{O}(4) \rightarrow \text{TOP}/\text{O}$ is 5-connected [FQ90, Theorem 8.7A], where only 3-connected is needed for Theorem 9.1. In other words, it was shown prior to Freedman's work that homotopy 4-spheres admit a smooth structure in the complement of a point, so the results that Freedman required were indeed known. However, smoothing in the complement of a point was not known for general connected, compact 4-manifolds until after the work of Quinn in 1982. Further discussion can also be found in Quinn [Qui84] and Lashof-Taylor [LT84].

Below we will give applications of Theorem 9.1; see e.g. the proof of Theorem 10.17.

9.2. THE KIRBY-SIEBENMANN INVARIANT AND STABLE SMOOTHING OF 4-MANIFOLDS

The formulation of the other two statements on smoothing 4-manifolds that we will give (Theorems 9.1 and 9.9) make use of the Kirby-Siebenmann invariant. The Kirby-Siebenmann invariant $\text{ks}(M) \in \mathbb{Z}/2$ of a compact 4-manifold is defined in [FQ90, Section 10.2B], or alternatively by [KS77, p. 318] or [Rud16, Definition 3.4.2], and we describe the construction now.

The homotopy fibre TOP/PL of the forgetful map $\text{BPL} \rightarrow \text{BTOP}$ has the homotopy type of a $K(\mathbb{Z}/2, 3)$ [KS77, Essay IV, §10, p. 194] and has the structure of a loop space, permitting the construction of the delooping $\text{B}(\text{TOP}/\text{PL})$ [BV68, Theorem C], [BV73], which is an Eilenberg-MacLane space of type $K(\mathbb{Z}/2, 4)$. A connected topological 4-manifold has a unique smooth structure on its boundary. Using the homotopy fibre sequence

$$\text{TOP}/\text{PL} \rightarrow \text{BPL} \rightarrow \text{BTOP} \rightarrow \text{B}(\text{TOP}/\text{PL}),$$

the unique obstruction to a lift of the classifying map $\tau_M: M \rightarrow \text{BTOP}$ of the stable topological tangent bundle to BPL is therefore a homotopy class in

$$[(M, \partial M), \underbrace{(\text{B}(\text{TOP}/\text{PL}), *)}_{=K(\mathbb{Z}/2, 4)}] \cong H^4(M, \partial M; \mathbb{Z}/2) = \mathbb{Z}/2.$$

Here, we used again that 4-manifolds have the homotopy type of a CW complex (Theorem 3.16). We refer to the corresponding element of $\mathbb{Z}/2$ as the *Kirby-Siebenmann invariant*, $\text{ks}(M)$, of the compact, connected manifold M . For disconnected compact 4-manifolds, $M = \bigsqcup_{i=1}^n M_i$, define

$$\text{ks}(M) := \sum_{i=1}^n \text{ks}(M_i) \in \mathbb{Z}/2.$$

For comparison, we note that in every dimension, a compact n -manifold with a PL structure on its boundary has a Kirby-Siebenmann invariant, which is a homotopy class of maps in $[(M, \partial M), (\mathbf{B}(\text{TOP}/\text{PL}), *)]$ determined by its topological tangent bundle. In the following theorem we summarise some key properties of the Kirby-Siebenmann invariant.

Theorem 9.2. *Let M and N be compact 4-manifolds.*

- (1) *If $M \times \mathbb{R}$ admits a smooth structure (e.g. if M admits a smooth structure), then $\text{ks}(M) = 0$.*
- (2) *The Kirby-Siebenmann invariant gives rise to a surjective homomorphism $\Omega_4^{\text{TOP}} \rightarrow \mathbb{Z}/2$. In particular for M a closed 4-manifold that bounds a compact 5-manifold, $\text{ks}(M) = 0$.*
- (3) *The Kirby-Siebenmann invariant is additive under the connected sum operation.*
- (4) *There is a short exact sequence*

$$0 \rightarrow \Omega_4^{\text{Spin}} \rightarrow \Omega_4^{\text{TOPSpin}} \xrightarrow{\sigma/8} \mathbb{Z}/2 \rightarrow 0,$$

with the first map the forgetful map and last map given by the signature divided by 8, modulo 2. This sequence does not split, so $\Omega_4^{\text{TOPSpin}} \cong \mathbb{Z}$. Moreover, for a closed spin manifold, the signature divided by 8, modulo 2, is equal to the Kirby-Siebenmann invariant, i.e. the map which takes $\sigma/8$ modulo 2 equals the composition $\Omega_4^{\text{TOPSpin}} \rightarrow \Omega_4^{\text{STOP}} \xrightarrow{\text{ks}} \mathbb{Z}/2$.

- (5) *If $S \subseteq \partial M$ and $T \subseteq \partial N$ are compact codimension zero submanifolds with a homeomorphism $S \cong T$, then*

$$\text{ks}(M \cup_{S \cong T} N) = \text{ks}(M) + \text{ks}(N).$$

- (6) *If there exists a compact 5-manifold with $\partial W = M \cup_{\partial M \cong \partial N} N$, for some homeomorphism $\partial M \cong \partial N$, then $\text{ks}(M) = \text{ks}(N)$.*

While they are certainly well-known to the experts, and frequently used, we could not find explicit proofs of these facts in the literature, so we give some details.

PROOF OF THEOREM 9.2. Let us prove (1). The topological tangent bundle of $M \times \mathbb{R}$ is isomorphic to $\tau_M \oplus \varepsilon$, where τ_M is the tangent microbundle of M and ε denotes a rank one trivial bundle over M . If $M \times \mathbb{R}$ admits a smooth structure, then there is a lift $\tau_{M \times \mathbb{R}}^{\text{Diff}} : M \times \mathbb{R} \rightarrow \text{BO}(5)$, the smooth tangent bundle to $M \times \mathbb{R}$. Let $p : \text{BO}(5) \rightarrow \text{BTOP}(5)$ be the canonical map. Then $\tau_M \oplus \varepsilon = p \circ \tau_{M \times \mathbb{R}}^{\text{Diff}}$. Passing to the stable classifying spaces, we obtain a lift $M \rightarrow \text{BO}$ whose composition with the canonical map $\text{BO} \rightarrow \text{BTOP}$ agrees with $\tau_M \oplus \varepsilon^\infty$, the stable tangent microbundle of M . Since the map $\text{BO} \rightarrow \text{BTOP}$ factors through $\text{BPL} \rightarrow \text{BTOP}$, we have a stable lift of τ_M and so $\text{ks}(M) = 0$. This completes the proof of (1).

Now to prove (2), suppose that a closed 4-manifold $M = \bigsqcup_{i=1}^k M_i$ bounds a compact 5-manifold W' . Perform 0 and 1-surgeries on W' to obtain a path connected, simply

connected, compact 5-manifold W with $\partial W = M$. We prove that

$$\text{ks}(M) := \sum_{i=1}^k \text{ks}(M_i) = 0.$$

Consider the diagram

$$\begin{array}{ccccccc} M_i & \longrightarrow & M & \longrightarrow & W & \xrightarrow{\tau_W} & \text{BTOP} \\ & & & \searrow^{\text{ks}(M)} & & & \downarrow \\ & & & & & & \text{B(TOP / PL)} \xrightarrow{\cong} K(\mathbb{Z}/2, 4) \\ & \searrow^{\text{ks}(M_i)} & & & & & \\ & & & & & & \end{array}$$

The restriction $M_i \rightarrow W \rightarrow \text{BTOP}$ equals the stable tangent microbundle of M , since M has a collar $M \times [0, 1] \subseteq W$ by Theorem 2.16. Therefore the diagram commutes. It follows that the top left horizontal map in the next diagram sends $\text{ks}(W)$ to $(\text{ks}(M_1), \dots, \text{ks}(M_k))$, so the map $H^4(W; \mathbb{Z}/2) \rightarrow \mathbb{Z}/2$ sends $\text{ks}(W)$ to $\sum_{i=1}^k \text{ks}(M_i) = \text{ks}(M)$.

$$\begin{array}{ccccccc} H^4(W; \mathbb{Z}/2) & \longrightarrow & H^4(M; \mathbb{Z}/2) & \xrightarrow{\cong} & \bigoplus_{i=1}^k H^4(M_i; \mathbb{Z}/2) & & \\ \text{PD} \cong \downarrow & & \text{PD} \cong \downarrow & & \text{PD} \cong \downarrow & \searrow^{\cong} & \bigoplus_{i=1}^k \mathbb{Z}/2 \xrightarrow{(1, \dots, 1)} \mathbb{Z}/2 \\ H_1(W, M; \mathbb{Z}/2) & \longrightarrow & H_0(M; \mathbb{Z}/2) & \xrightarrow{\cong} & \bigoplus_{i=1}^k H_0(M_i; \mathbb{Z}/2) & \xrightarrow{=} & \end{array}$$

The left square of this diagram commutes by Poincaré-Lefschetz duality. The middle square and the triangle commute trivially. But since W is connected and simply connected, every element of $H_1(W, M; \mathbb{Z}/2)$ can be represented by a (possibly empty) union of arcs with boundary on M . Thus the image of $\text{ks}(W)$ in $\bigoplus_{i=1}^k H_0(M_i; \mathbb{Z}/2)$ is nonzero in evenly many summands, and therefore its image in $\mathbb{Z}/2$ on the far right is zero. By commutativity of the diagram it follows that $\text{ks}(M) = 0$, as desired.

Now (2) follows. First note that the addition on Ω_4^{TOP} is by disjoint union, so ks is additive by definition. We have just shown that the map $\text{ks}: \Omega_4^{\text{TOP}} \rightarrow \mathbb{Z}/2$ is well-defined, since for M a closed 4-manifold that bounds a compact 5-manifold, $\text{ks}(M) = 0$. Therefore $\text{ks}: \Omega_4^{\text{TOP}} \rightarrow \mathbb{Z}/2$ is a homomorphism as desired.

Freedman [Fre82, Theorem 1.7] showed that there exists a closed 4-manifold with intersection pairing that is isometric to E_8 . This manifold is called the E_8 manifold. Since the E_8 form has signature 8, $\sigma(E_8) = 8$, so $\sigma(E_8)/8 = 1$. Since the E_8 form is even, and $H_1(E_8; \mathbb{Z}) = 0$, E_8 is spin. The construction of the E_8 manifold was a key step in

the proof of the Classification Theorem 12.2. We will now show that $\text{ks}(E_8) = 1$. To see this note that E_8 cannot be smoothed, even after adding copies of $S^2 \times S^2$, by the Rochlin Theorem 8.8 that every closed spin smooth 4-manifold has signature divisible by 16. Whereas if $\text{ks}(E_8) = 0$, then E_8 would be stably smoothable by Theorem 9.9. Therefore $\text{ks}: \Omega_4^{\text{TOP}} \rightarrow \mathbb{Z}/2$ is surjective.

Now we can prove (3) easily. Observe that a disjoint union $M \sqcup N$ is cobordant to $M \# N$ via the cobordism

$$(M \times I \sqcup N \times I) \cup_{S^0 \times D^4} (D^1 \times D^4),$$

with $\{-1\} \times D^4$ embedded in the interior of $M \times \{1\}$, and $\{1\} \times D^4$ embedded in the interior of $N \times \{1\}$. Then we have just shown that the Kirby-Siebenmann invariant vanishes on $M \# N \sqcup M \sqcup N$ and therefore $\text{ks}(M \# N) = \text{ks}(M) + \text{ks}(N) \in \mathbb{Z}/2$.

To prove (4), we consider the following diagram. The maps between bordism groups are structure forgetting maps, so the diagram commutes.

$$\begin{array}{ccccccc} 0 & \longrightarrow & \Omega_4^{\text{Spin}} & \longrightarrow & \Omega_4^{\text{TOPSpin}} & \xrightarrow{\text{ks}} & \mathbb{Z}/2 \longrightarrow 0 \\ & & \downarrow \cdot 16 & & \downarrow & & \downarrow = \\ 0 & \longrightarrow & \Omega_4^{\text{SO}} & \longrightarrow & \Omega_4^{\text{STOP}} & \xrightarrow{\text{ks}} & \mathbb{Z}/2 \longrightarrow 0 \end{array}$$

Recall that $\Omega_4^{\text{SO}} \cong \mathbb{Z}$ given by the signature and generated by $\mathbb{C}\mathbb{P}^2$. The signature provides a splitting homomorphism, so $\Omega_4^{\text{STOP}} \cong \mathbb{Z} \oplus \mathbb{Z}/2$. Also $\Omega_4^{\text{Spin}} \cong \mathbb{Z}$ given by the signature divided by 16 and generated by the $K3$ surface, so the forgetful map $\Omega_4^{\text{Spin}} \rightarrow \Omega_4^{\text{SO}}$ becomes, on identifying domain and codomain with \mathbb{Z} , multiplication by 16.

Both sequences are exact: a smooth manifold has vanishing ks invariant, and vanishing $\text{ks}(M)$ implies smoothable after adding copies of $S^2 \times S^2$ by Theorem 9.9 below. Since $M \# (S^2 \times S^2)$ is (spin) bordant to M , the sequences are exact at their middle terms. The maps labelled ks are surjective because the E_8 manifold is spin and has $\text{ks}(E_8) = 1$, as discussed in the proof of (2). Finally, after surgery to make it 1-connected, a topological null bordism of a compact smooth 4-manifold can be smoothed by high dimensional smoothing theory, so the left hand maps are injective.

We claim that the sequence in the upper row does not split. Consider the $K3$ surface generating $\Omega_4^{\text{Spin}} \cong \mathbb{Z}$. By the down-then-left route, $[K3]$ maps to $(16, 0) \in \mathbb{Z} \oplus \mathbb{Z}/2 \cong \Omega_4^{\text{STOP}}$. On the other hand the E_8 manifold represents a class in $\Omega_4^{\text{TOPSpin}}$ and maps to $(8, 1) \in \mathbb{Z} \oplus \mathbb{Z}/2 \cong \Omega_4^{\text{STOP}}$.

Since ks is a homomorphism by (2), we see that $2 \cdot [E_8]$, which equals $[E_8 \# E_8]$ by (3), maps to $0 \in \mathbb{Z}/2$ and so has trivial ks invariant. By exactness of the top row it lies in the image of Ω_4^{Spin} . Let N be a closed spin smooth 4-manifold TOPSpin-bordant to $E_8 \# E_8$. Since $\sigma(E_8 \# E_8) = 16 = \sigma(K3)$, we have $[N] = [K3] \in \Omega_4^{\text{Spin}}$. It follows that $K3$, the generator of $\Omega_4^{\text{Spin}} \cong \mathbb{Z}$, maps to $2 \cdot [E_8] \in \Omega_4^{\text{TOPSpin}}$. Thus we have a diagram with exact

rows:

$$\begin{array}{ccccccc}
0 & \longrightarrow & \mathbb{Z} & \xrightarrow{\cdot 2} & \mathbb{Z} & \longrightarrow & \mathbb{Z}/2 \longrightarrow 0 \\
& & \downarrow = & & \downarrow 1 \mapsto [E_8] & & \downarrow = \\
0 & \longrightarrow & \mathbb{Z} & \xrightarrow{1 \mapsto 2 \cdot [E_8]} & \Omega_4^{\text{TOPSpin}} & \xrightarrow{\text{ks}} & \mathbb{Z}/2 \longrightarrow 0.
\end{array}$$

Since $\text{ks}(E_8) = 1$, the diagram commutes. Then by the five lemma, $\Omega_4^{\text{TOPSpin}} \cong \mathbb{Z}$, generated by E_8 , and the sequence does not split, as claimed. For a topological spin, compact 4-manifold, $\sigma/8$ is an integer, by Proposition 8.4. By Proposition 8.3 we know that the signature is additive. It follows from this observation and the fact that $\Omega_4^{\text{TOPSpin}}$ is generated by E_8 that $M \mapsto \sigma(M)/8$ gives rise to the isomorphism $\Omega_4^{\text{TOPSpin}} \cong \mathbb{Z}$.

The diagram

$$\begin{array}{ccccc}
\Omega_4^{\text{TOPSpin}} & \xrightarrow[\cong]{\sigma/8} & \mathbb{Z} & & \\
\downarrow & & \downarrow 1 \mapsto (8,1) & \searrow & \\
\Omega_4^{\text{STOP}} & \xrightarrow[\cong]{} & \mathbb{Z} \oplus \mathbb{Z}/2 & \xrightarrow{\text{pr}_2} & \mathbb{Z}/2,
\end{array}$$

which commutes by computing on the generator E_8 of $\Omega_4^{\text{TOPSpin}} \cong \mathbb{Z}$, shows that $\text{ks}(M) = \sigma(M)/8 \in \mathbb{Z}/2$ for TOPSpin manifolds M . This completes the proof of (4).

To prove (5), it was suggested by Jim Davis to consider the exact sequence

$$\Omega_4^{\text{O}} \rightarrow \Omega_4^{\text{TOP}} \rightarrow \Omega_4^{\{\text{O} \rightarrow \text{TOP}\}} \rightarrow \Omega_3^{\text{O}} = 0.$$

Here elements of $\Omega_4^{\{\text{O} \rightarrow \text{TOP}\}}$ are represented by compact topological 4-manifolds with smooth boundary, considered up to 5-dimensional cobordism relative to a smooth cobordism on the boundary. That is, 4-manifolds with boundary $(M, \partial M)$ and $(N, \partial N)$ are equivalent if there is a compact 5-manifold W with boundary

$$\partial W = M \cup_{\partial M} \partial_{\text{vert}} W \cup_{\partial N} N,$$

for some smooth 4-dimensional cobordism $\partial_{\text{vert}} W$ with boundary $\partial M \sqcup \partial N$.

By the exact sequence, $\Omega_4^{\{\text{O} \rightarrow \text{TOP}\}}$ is isomorphic to the cokernel of $\Omega_4^{\text{O}} \rightarrow \Omega_4^{\text{TOP}}$. We claim that this cokernel is isomorphic to $\mathbb{Z}/2$ via the Kirby-Siebenmann invariant. To see this, by (2) there is a surjective homomorphism $\text{ks}: \Omega_4^{\text{TOP}} \rightarrow \mathbb{Z}/2$. If $\text{ks}(M) = 0$ then M is stably smoothable by Theorem 9.9, so M is bordant to a smooth manifold and therefore lies in the image of Ω_4^{O} . If M is smooth, then $\text{ks}(M)$ is zero, so the sequence $\Omega_4^{\text{O}} \rightarrow \Omega_4^{\text{TOP}} \xrightarrow{\text{ks}} \mathbb{Z}/2 \rightarrow 0$ is exact, and we may identify this sequence with the given sequence.

To prove (5), we therefore need that the disjoint union $M \sqcup N$ is bordant to $M \cup_{S \cong T} N$, where $S \subseteq \partial M$ and $T \subseteq \partial N$ are compact codimension zero submanifolds with a choice of homeomorphism $S \cong T$. Here is a construction of such a bordism. For $I = [0, 1]$, take

$$(M \times I) \sqcup (S \times I \times [1/2, 1]) \sqcup (N \times I),$$

identify

$$S \times \{0\} \times [1/2, 1] \sim S \times [1/2, 1] \subseteq (M \times [1/2, 1]),$$

and, using the identification $S \cong T$, identify

$$S \times \{1\} \times [1/2, 1] \sim T \times [1/2, 1] \subseteq N \times [1/2, 1].$$

Let W be the result of this gluing and some rounding of corners. The boundary of W is

$$(M \sqcup N) \cup_{\partial M \sqcup \partial N} \partial_{\text{vert}} W \cup_{\partial(M \cup_{S \cong T} N)} M \cup_{S \cong T} N,$$

where

$$\begin{aligned} \partial_{\text{vert}} W &= (\partial M \times [0, 1/2]) \cup (\overline{\partial M \setminus S} \times [1/2, 1]) \\ &\cup (S \times I \times \{1/2\}) \cup (\partial S \times I \times [1/2, 1]) \\ &(\partial N \times [0, 1/2]) \cup (\overline{\partial N \setminus T} \times [1/2, 1]). \end{aligned}$$

This shows that $M \sqcup N$ and $M \cup_{S \cong T} N$ are equal in $\Omega_4^{\{0 \rightarrow \text{TOP}\}}$, and therefore have the same Kirby-Siebenmann invariants. Since $\text{ks}(M \sqcup N) = \text{ks}(M) + \text{ks}(N)$, this completes the proof of (5).

Finally we prove (6). If $M \cup_{\partial M = \partial N} N$ bounds a compact 5-manifold, then by (2) we have that $\text{ks}(M \cup_{\partial} N) = 0$. By (5), $\text{ks}(M) + \text{ks}(N) = \text{ks}(M \cup_{\partial} N) \in \mathbb{Z}/2$. Therefore $\text{ks}(M) = \text{ks}(N)$ as required. This proves (6) and therefore completes the proof of Theorem 9.2. \square

The following theorem says that the converse to Theorem 9.2 (1) holds for M connected.

Theorem 9.3. *If M is a compact, connected 4-manifold with vanishing Kirby-Siebenmann invariant, then $M \times \mathbb{R}$ admits a smooth structure.*

PROOF. The vanishing of the Kirby-Siebenmann invariant implies that there is a lift of $\tau_M: M \rightarrow \text{BTOP}$ to a map $M \rightarrow \text{BPL}$. Since PL/O is 6-connected [FQ90, Theorem 8.3B], [HM74, Proof of 4.13], there is in fact a lift $\tilde{\tau}_M: M \rightarrow \text{BO}$. This corresponds to a lift $\tilde{\tau}_M \oplus \varepsilon^n: M \rightarrow \text{BO}(4+n)$, for some n . This in turn corresponds to a lift

$$\tilde{\tau}_{M \times \mathbb{R}^n}: M \times \mathbb{R}^n \rightarrow \text{BO}(4+n)$$

of the tangent microbundle $\tau_{M \times \mathbb{R}^n}: M \times \mathbb{R}^n \rightarrow \text{BTOP}$. By [KS77, Essay V, Theorem 1.4 p. 222], there exists a corresponding smooth structure on $M \times \mathbb{R}^n$. Then apply the Product Structure Theorem 4.16 [KS77, Essay I, Theorem 5.1, p. 31], to deduce the existence of a smooth structure on $M \times \mathbb{R}$, using that the dimension of $M \times \mathbb{R}$ is at least five. \square

Example 9.4. Here is an application of Theorem 9.3. By the classification of simply connected, closed 4-manifolds [FQ90, Section 10.1] (see also our Theorem 12.2), there is a simply connected, closed 4-manifold N with intersection form $E_8 \oplus E_8$. As the form is even, the manifold is spin; see Remark 8.9. Since this form is not diagonalisable over \mathbb{Z} , by Donaldson's Theorem [Don83] (Theorem 8.11) this 4-manifold does not admit a smooth structure. However the Kirby-Siebenmann invariant of N vanishes, since for a closed simply connected 4-manifold M with even intersection form, the Kirby-Siebenmann

invariant $\text{ks}(M)$ coincides with $\sigma(M)/8 \pmod{2}$ (Theorem 9.2(4)), and $E_8 \oplus E_8$ is rank 16 and positive definite, with signature 16. Therefore $N \times \mathbb{R}$ admits a smooth structure by Theorem 9.3, even though N does not.

We give a straightforward consequence of Theorem 9.2 in the following. We first recall a definition. Let N be an integral homology 3-sphere. As the 3-dimensional spin bordism group is trivial $\Omega_3^{\text{Spin}} = 0$ [GS99, Theorem 5.7.14], we may pick a smooth, compact, orientable, spin 4-manifold M with boundary N . The *Rochlin invariant* $\mu(N) \in \mathbb{Z}/2$ is defined as the quantity $\sigma(M)/8 \pmod{2}$. This is well-defined, as a consequence of Novikov Additivity [Kir89, Theorem 5.3] and Theorem 8.8.

Theorem 9.5. *Let M be a compact, oriented, spin manifold with boundary N an integral homology 3-sphere. Then $\text{ks}(M) = \sigma(M)/8 + \mu(N) \in \mathbb{Z}/2$.*

PROOF. As $\Omega_3^{\text{Spin}} = 0$ [GS99, Theorem 5.7.14], we may pick a smooth, compact, orientable, spin 4-manifold X with boundary N . Form the closed, spin 4-manifold $Z = -X \cup_N M$. By Theorem 9.2 (5), and using that X is smooth, we have $\text{ks}(Z) = \text{ks}(X) + \text{ks}(M) = \text{ks}(M) \in \mathbb{Z}/2$. By Theorem 9.2 (4), we have $\text{ks}(Z) = \sigma(Z)/8 \in \mathbb{Z}/2$. Using Novikov additivity and the facts so far, we have

$$\text{ks}(M) = \text{ks}(Z) = \sigma(Z)/8 = \sigma(X)/8 + \sigma(M)/8 = \mu(N) + \sigma(M)/8 \in \mathbb{Z}/2. \quad \square$$

Lemma 9.6. *There exists a unique closed 4-manifold that is homotopy equivalent to $\mathbb{C}P^2$ and with nontrivial Kirby-Siebenmann invariant, and so which in particular is not homeomorphic to $\mathbb{C}P^2$.*

PROOF. Let $K \subseteq \partial D^4$ be any knot with $\text{Arf}(K) = 1 \in \mathbb{Z}/2$ (for example, the trefoil). Attach a 2-handle $D^2 \times D^2$ to D^4 by identifying $S^1 \times D^2$ with a tubular neighbourhood of K , via the +1 framing of K , to obtain the 4-manifold with boundary $X(K)$. The boundary $\partial X(K)$ is an integral homology sphere. Freedman proved that every integral homology sphere bounds a compact, contractible 4-manifold [Fre82, Theorem 1.4'], [FQ90, Corollary 9.3C]. Write Y for this contractible 4-manifold with $\partial Y \cong \partial X(K)$ and form the closed 4-manifold $Z = -Y \cup X(K)$.

The manifold Z is homotopy equivalent to $\mathbb{C}P^2$. Indeed, $H^2(Z; \mathbb{Z}) \cong \mathbb{Z}$ so we obtain a map $Z \rightarrow \mathbb{C}P^\infty = K(\mathbb{Z}, 2)$ representing a generator. This map can now be homotoped to have domain $\mathbb{C}P^2 \subseteq \mathbb{C}P^\infty$. This map will be a homology equivalence (this is clear on H_1, H_2 , and H_3 , and can be deduced on H_4 by considering that the cohomology ring of Z agrees with that of $\mathbb{C}P^2$ by Poincaré duality). As Z is simply connected, Whitehead's theorem now implies we have a homotopy equivalence.

We show that $\text{ks}(Z) = 1 \in \mathbb{Z}/2$. The Rochlin invariant of +1 surgery on a knot in S^3 is equal to its Arf invariant (see e.g. [Sav02, Example 2.5]), and thus $\mu(\partial X(K)) = 1 \in \mathbb{Z}/2$. Applying Theorem 9.5 to the contractible manifold Y , we obtain $\text{ks}(Y) = \mu(\partial X(K)) = 1 \in \mathbb{Z}/2$. On the other hand, $\text{ks}(X(K)) = 0$ because this manifold is smooth. By Theorem 9.2 (5), $\text{ks}(Z) = \text{ks}(Y) + \text{ks}(X(K)) = 1 + 0 = 1$.

By Freedman's classification (Theorem 12.2), Z is the unique closed manifold that is homotopy equivalent to $\mathbb{C}P^2$ with $\text{ks}(Z) = 1$. \square

Definition 9.7. The unique closed 4-manifold that is homotopy equivalent to $\mathbb{C}P^2$ but not homeomorphic to $\mathbb{C}P^2$ (constructed above) is called the *Chern manifold* and denoted $*\mathbb{C}P^2$.

Remark 9.8. The Chern manifold was first constructed in [Fre82, p. 370]. It is not smoothable because $\text{ks}(*\mathbb{C}P^2) = 1$. For further discussion of star partners, see [FQ90, Section 10.4], [Sto94], and [Tei97].

The following theorem says in particular that given any compact 4-manifold M there exists a closed, orientable, simply-connected 4-manifold N such $M\#N$ is smoothable.

Theorem 9.9. *Let M be compact, connected 4-manifold. There exists a closed, orientable, simply connected 4-manifold N such $M\#N$ admits a smooth structure. If moreover the Kirby-Siebenmann invariant of M is zero, then there exists a $k \in \mathbb{N}_0$ such that $M\#^k S^2 \times S^2$ admits a smooth structure.*

PROOF. Let M be compact 4-manifold. Perform the connected sum with an appropriate number of copies of $*\mathbb{C}P^2$, in order to obtain a manifold with every connected component having zero Kirby-Siebenmann invariant. It follows from the discussion on [FQ90, p. 164] and the Sum-Stable Smoothing Theorem [FQ90, p. 125], that performing the connected sum with enough copies of $S^2 \times S^2$ produces a manifold that admits a smooth structure. \square

Remark 9.10. Given a lift of the classifying map of the (unstable) tangent microbundle of M to $\text{BO}(4)$, Lashof-Shaneson [LS71] showed that there exists a $k \in \mathbb{N}_0$ such that $M\#^k S^2 \times S^2$ admits a smooth structure. The result quoted in the previous proof extended this to require only a lift of the corresponding *stable* maps to deduce the same result. The existence of a stable lift is significantly easier to verify.

CHAPTER 10

Topological transversality


We turn to the subject of transversality in the topological category. Some discussion of this concept is in order. There are two important contexts for transversality: submanifold transversality and map transversality. In this book, map transversality will be deduced from submanifold transversality. Submanifold transversality when none of the manifolds involved has dimension 4 is due to Marin [Mar77]; cf. [KS77, Essay III, Section 1, p. 83]. Transversality in the remaining cases is due to Quinn [Qui82, Qui88]; see also [FQ90, Section 9.5].

A naive definition of submanifold transversality in the topological category is that manifolds are *locally transverse* if around any intersection point there is a chart in which the submanifolds appear as perpendicular planes. On the other hand, there are examples (in the relative setting, in high dimensions) of submanifolds which cannot be made locally transverse via ambient isotopy; see Remark 10.4. Thus one cannot generally use this definition.

In light of this, in order to make general statements, one passes to some notion of *global transversality*. Global transversality means that transversality statements are made with respect to a given choice of normal structure on one of the submanifolds involved. Of course, this forces one to engage with the question of existence and uniqueness of whatever normal structure is used, and the ‘correct’ choice of normal structure is still not fully settled in the topological category. We refer the reader to [FQ90, Sections 9.4, 9.6C] for a brief discussion of the competitors.

The most general statement of transversality [Qui88, Theorem] uses microbundles to describe normal structure, and this is the technology we will use. As discussed in Chapter 6, for general manifolds, tangent microbundles always exist but normal microbundles do not (see Example 6.23).

The case of dimension 4 is special, since here the normal vector bundles of Section 5.2, which are a stronger notion than normal microbundles, always exist. In fact, the results obtained for these normal vector bundles in dimension 4 are strong enough to ensure that submanifold transversality holds in ambient dimension 4 with the naive, local transversality definition discussed above. The reader may therefore wonder why we even introduce normal microbundles into a discussion primarily focused on 4-manifold transversality. The answer is that the ‘submanifold transversality implies map transversality’ argument of Section 10.2 requires a bundle technology that works in all dimensions, and microbundles appear to be the most convenient.

FIGURE 10.1. Sketch of a transverse intersection of Y to νX .

10.1. TRANSVERSALITY FOR SUBMANIFOLDS

Definition 10.1. Consider proper submanifolds X, Y of an ambient manifold and a normal microbundle νX for X , with projection $r_X: E(\nu X) \rightarrow X$. The proper submanifold Y is *transverse to νX* if there exists a neighbourhood $U \subseteq E(\nu X)$ of X such that $Y \cap U = r_X^{-1}(X \cap Y) \cap U$.

Lemma 10.2. *Let X, Y be submanifolds of a manifold M . Let Y be transverse to a normal microbundle νX of X . Then $X \cap Y$ is a submanifold of Y with normal microbundle $(\nu X)|_{X \cap Y}$.*

PROOF. Let $X \xrightarrow{i} \nu X \xrightarrow{p} X$ be a normal microbundle of X . Once we have established that $(\nu X)|_{X \cap Y}$ is a normal microbundle of $X \cap Y$ in Y , the subspace $X \cap Y$ will automatically be a submanifold since the trivialisation of the microbundle $(\nu X)|_{X \cap Y}$ gives the required charts for $X \cap Y$.

At least after shrinking the total space of $E(\nu X)$, each fibre $p^{-1}(x)$ for $x \in X \cap Y$ will be contained in Y by the definition of transversality. That is $E((\nu X)|_{X \cap Y})$ is a subset of Y and neighbourhood of $X \cap Y$. This shows that $(\nu X)|_{X \cap Y}$ is a normal microbundle of $X \cap Y \subseteq Y$. \square

Transversality in high dimensions is due to Marin [Mar77], cf. [KS77, Essay III, Section 1, p. 83]. The formulation below is from Quinn [Qui88]. Recall Definition 2.5 of a proper submanifold. Note that in the next theorem there is no restriction on dimensions. The manifold M is allowed to be noncompact and have nonempty boundary.

Theorem 10.3. (Transversality Theorem for submanifolds) *Let X and Y be proper submanifolds of a compact manifold M . Let νY be a normal microbundle for Y . Let $C \subseteq M$ be a closed subset such that X is transverse to νY in a neighbourhood of C . Let U be a neighbourhood of the set $(M \setminus C) \cap X \cap Y$. Then there exists an isotopy of X supported in U to a proper submanifold X' such that X' is transverse to νY .*

PROOF. See Quinn [Qui88] for all cases but $\dim M = 4$, $\dim X = 2$ and $\dim Y = 2$. For the remaining case, first establish local transversality using [FQ90, Section 9.5]. Note

that $X \cap Y$ is a discrete collection of points. Therefore, the coordinate chart, witnessing local transversality, defines a normal neighbourhood of Y near $X \cap Y$. This normal vector bundle can be extended to a normal vector bundle $\nu Y'$ on all of Y by [FQ90, Theorem 9.3A]. The submanifold X is now transverse to $\nu Y'$, but (possibly) not to νY . By Theorem 6.24, our microbundle νY comes from a normal vector bundle. By uniqueness of normal vector bundles (Theorem 5.17), there is an isotopy from $\nu Y'$ to νY . Apply this isotopy to X . Now X is transverse to νY . \square

Remark 10.4. The analogous statement to Theorem 10.3 is false for local transversality. Examples of this failure even exist in the PL category: Hudson [Hud69] constructs, for certain large n , closed PL submanifolds $X, Y \subseteq \mathbb{R}^n$, that are topologically unknotted Euclidean spaces of codimension ≥ 3 , in such a way that X and Y are PL locally transverse near a closed neighbourhood K of infinity but also so that it is impossible to move X and Y by isotopy relative to K to make them locally transverse everywhere.

Although transversality for submanifolds (Theorem 10.3) is only stated for a pair of submanifolds, it can be used to make collections of submanifolds transverse.

Lemma 10.5. *Let M be an $2m$ -dimensional manifold for $m \geq 1$, and let X_1, \dots, X_n be m -dimensional compact submanifolds with normal microbundles νX_i . Then the submanifolds X_i can be isotoped such that there are no triple intersection points and the submanifolds intersect (pairwise) transversely.*

PROOF. We give a proof by induction. When $n = 1$, there is nothing to show, since every submanifold is embedded. For the inductive step, denote X_n by Y . The inductive hypothesis states that we can isotope any $n - 1$ submanifolds X_1, \dots, X_{n-1} so there are no triple points and that they intersect pairwise transversely. We will prove that the submanifolds X_1, \dots, X_{n-1} can be further isotoped so that they are transverse to νY and Y is free of triple points. Note that having no triple points on Y implies that there exists an open set U_Y such that: $X_i \cap X_j \subseteq U_Y$ for $1 \leq i < j \leq n - 1$, and $M \setminus U_Y$ is a neighbourhood of Y . To obtain the lemma apply the inductive hypothesis, picking all further isotopies to be supported in U_Y .

We proceed by showing the inductive step: we can isotope every X_i to be transverse to νY such that no triple points lie on Y . For each $i = 1, \dots, n - 1$, apply Theorem 10.3 to arrange that Y and X_i intersect transversely. By compactness of the submanifolds, the subset

$$T_Y = Y \cap (X_1 \cup \dots \cup X_{n-1})$$

is compact. Pick disjoint open neighbourhoods $V_y \subseteq Y$ around each point $y \in T_Y$. Pick a chart ϕ of Y around $\phi(0) = y$ contained in V_y , and a microbundle chart around $y \in Y$. In the local model, Y corresponds to $\mathbb{R}^m \times \{0\}$ and the X_i that intersect Y in y will be mapped to $0 \times \mathbb{R}^m$. For those X_i , pick disjoint points $u_i \in \mathbb{R}^m$ (here we use $m > 0$), and pick a continuous function $\eta: \mathbb{R}_{\geq 0} \rightarrow [0, 1]$ with $\eta(t) = 1$ for $0 \leq t \leq 1$ and $\eta(t) = 0$ for $t \geq 2$.




FIGURE 10.2. Displacing a triple point y in a microbundle chart.

Replace X_i in the chart with the image of

$$\begin{aligned} \mathbb{R}^m &\rightarrow \mathbb{R}^m \times \mathbb{R}^m \\ v &\mapsto (\eta(\|v\|)u_i, v). \end{aligned}$$

Call this new submanifold X'_i . It agrees with X_i outside the ball of radius 2, and is isotopic to X_i . In V_y , the submanifold X'_i intersects Y only in $\phi(u_i)$ and there it intersects Y transversely with respect to νY . The collection $\{X'_i\}$ has no triple intersection points in the set V_y anymore. \square

Here is another result on submanifold transversality. It might often happen that one can find a continuous map of, for example, a disc D^2 into a 4-manifold M , perhaps if fundamental group computations yield a null homotopy of a circle. Then this disc can be isotoped to a generic immersion. If M were smooth, this would be a consequence of standard differential topology, an observation that we leverage.

Theorem 10.6. *Let Σ be a connected, compact 1 or 2 dimensional manifold and let $f: (\Sigma, \partial\Sigma) \rightarrow (M, \partial M)$ be a continuous map of Σ into a connected 4-manifold M , such that f is a smooth embedding near $\partial\Sigma$. Then there is an homotopy of f rel. a (possibly smaller) neighbourhood of $\partial\Sigma$ to a generic immersion $f': (\Sigma, \partial\Sigma) \rightarrow (M, \partial M)$.*

PROOF. We start out with the following claim.

Claim. The map $f: \Sigma \rightarrow M$ is homotopic to a map that misses a point $P \in M$.

We pick $P \in M \setminus \partial M$. Since $\dim(\Sigma) \leq 2$ we can equip Σ with a smooth structure. Using a chart we can equip an open neighbourhood V of P with a smooth structure. We pick another open neighbourhood U of P with $\bar{U} \subseteq V$. Since $f^{-1}(V \setminus \bar{U})$ is an open subset of Σ we can find a compact submanifold $F \subseteq \Sigma$ with $f^{-1}(\bar{U}) \subseteq \text{Int}(F) \subseteq F \subseteq f^{-1}(V)$. The map $f: F \rightarrow V$ is now a map between smooth manifolds. Thus, using Whitney

approximation, we can find a homotopy rel. ∂F from f to a smooth map $g: F \rightarrow V$. Since $\dim(F) < \dim(U)$ we see that this map misses a point in U . Since $f|_{\partial F} = g|_{\partial F}$, and ∂F is fixed throughout the homotopy, we can extend, by a constant homotopy, the homotopy from $f: F \rightarrow V$ to $g: F \rightarrow V$ to a homotopy from $f: \Sigma \rightarrow M$ to a map $g: \Sigma \rightarrow M$ such that f and g agree outside of F . Thus g also misses the point P . This concludes the proof of the claim.

Using Theorem 9.1 we smooth M in the complement of that point P . Now by [Hir94, Theorem 2.2.6 and Theorem 2.2.12] we can isotope f rel. $\partial\Sigma$ to a smooth immersion, which we can then isotope rel. $\partial\Sigma$ to a smooth generic immersion f' , i.e. a map that is self-transverse by [Hir94, Theorem 4.2.1], [Wal16, Theorem 4.6.6], with no triple points by general position [Wal16, Theorem 4.7.7] or [GG73, Chapter III, Corollary 3.3]. We have now shown that $f: \Sigma \rightarrow M$ is homotopic to a map $f': \Sigma \rightarrow M \setminus \{P\}$ which is a smooth generic immersion. But this implies that $f': \Sigma \rightarrow M \setminus \{P\}$ is in particular a generic immersion. But then $f': \Sigma \rightarrow M$ is also an generic immersion. \square

No purely topological proof of this is known. When manipulating generic immersions of surfaces, it is helpful to have control on the homotopies between them.

Definition 10.7. A *generic homotopy* between generic immersions $F_0, F_1: \Sigma \rightarrow M$ as in Definition 2.13 is sequence of ambient isotopies, finger moves, Whitney moves, and cusp homotopies.

We finish by quoting the following theorem, which was stated in [FQ90], and proven in [PRT19].

Theorem 10.8. *Every homotopy $H: \Sigma \times [0, 1] \rightarrow M$ between generic immersions $F_0, F_1: \Sigma \rightarrow M$ is homotopic rel. $\Sigma \times \{0, 1\}$ to a generic homotopy.*

10.2. TRANSVERSALITY FOR MAPS

Definition 10.9. Let $f: M \rightarrow N$ be a continuous map between two manifolds and let X be a submanifold of N with normal microbundle νX . The map f is said to be *transverse* to νX if $f^{-1}(X)$ is a submanifold admitting a normal microbundle $\nu f^{-1}(X)$ and

$$\begin{aligned} f: \nu f^{-1}(X) &\rightarrow f^* \nu X \\ m &\mapsto (r(m), f(m)) \end{aligned}$$

is an isomorphism of microbundles.

In the next theorem, we show how to reduce transversality for maps to transversality for submanifolds. Again, there are restrictions neither on dimensions nor codimensions.

Theorem 10.10. *Let M and N be a manifolds, let $Y \subseteq N$ be a proper submanifold with normal microbundle νY , let $f: M \rightarrow N$ be a map such that $f^{-1}(Y)$ is a submanifold of M , and let U be a neighbourhood of the set*

$$\text{graph } f \cap (M \times Y) \subseteq M \times N.$$




FIGURE 10.3. Transversality for maps from transversality for submanifolds.

Then there exists a homotopy $F: M \times I \rightarrow N$ such that

- (1) $F(m, 0) = f(m)$ for all $m \in M$;
- (2) $F_1: m \mapsto F(m, 1)$ is transverse to νY ; and
- (3) for $m \in M$ either
 - (a) $(m, f(m)) \notin U$, in which case $F(m, t) = f(m)$ for all $t \in I$, or
 - (b) $(m, f(m)) \in U$, in which case $(m, F(m, t)) \in U$ for all $t \in I$.

PROOF. Note that $M \times Y \subseteq M \times N$ is a proper submanifold with normal microbundle $M \times \nu Y = \text{pr}_Y^* \nu Y$. Also $\text{graph } f$ is a proper submanifold of $M \times N$. By Theorem 10.3, there exists an isotopy

$$G: \text{graph } f \times I \rightarrow M \times N,$$

supported in U , of the submanifold $\text{graph } f$ to a submanifold $T \subseteq M \times N$ such that T is transverse to $M \times \nu Y$ over $M \times Y$.

Define the map F as the composition

$$F: M \times I \rightarrow \text{graph } f \times I \xrightarrow{G} M \times N \xrightarrow{\text{pr}_N} N.$$

Since the isotopy G is supported in U , statement (3) holds. By construction, $F(x, 0) = \text{pr}_N(x, f(x)) = f(x)$, which proves statement (1).

Now we prove statement (2). Let $F_1: M \rightarrow N$ be the map that sends $x \mapsto F(x, 1)$. We keep track of the preimages through the maps of the composition that defines F_1 ; see Figure 10.3. By transversality of T to $M \times \nu Y$, we see that $Z = T \cap (M \times Y) = \text{pr}_N^{-1}(Y)$ is a submanifold of T with normal microbundle $M \times \nu Y|_Z$, and that the projection to N induces a microbundle isomorphism $M \times \nu Y|_Z \xrightarrow{\sim} \text{pr}_N^* \nu Y$. By definition, $\text{pr}_N: T \rightarrow N$ is transverse to νY .

We transport the submanifold Z back to M . Consider the commutative diagram

$$\begin{array}{ccc}
 & & N \\
 & \xrightarrow{F_1} & \uparrow \text{pr}_N \\
 M & \xrightarrow{\cong} \text{graph } f \times \{1\} \xrightarrow{\cong} & T \\
 & \searrow q & \\
 & &
 \end{array}
 ,$$

where q is the composition, which is a homeomorphism. Now $F_1^{-1}(Y) = q^{-1}(Z)$ is a submanifold with normal microbundle

$$q^*(M \times \nu Y|_Z) = q^* \text{pr}_N^* \nu Y = F_1^* \nu Y.$$

That is $F_1: M \rightarrow N$ is transverse to νY . □

10.3. REPRESENTING HOMOLOGY CLASSES BY SUBMANIFOLDS

Our first goal in this section is to prove the following theorem.

Theorem 10.11. *Let X be a compact orientable n -manifold. Let $k = n-2$ or $k = n-1$ and let $\sigma \in H_k(X, \partial X; \mathbb{Z})$. Then the class σ can be represented by a k -dimensional submanifold Y with $\partial Y \subseteq \partial X$.*

We will prove the theorem using purely topological methods, in particular the topological transversality arguments from Chapter 10, in particular Sections 10.1 and 10.2. Then we will prove a refined version in dimension four that uses the trick of smoothing away from a point. We will need the notion of a Thom class.

Definition 10.12. Let $\xi = S \xrightarrow{i} E \xrightarrow{p} S$ be a k -dimensional microbundle over S . For each $x \in S$ we write $E_x := p^{-1}(\{x\})$. A *Thom class* of ξ is a class $\tau(\xi) \in H^k(E, E \setminus i(S); \mathbb{Z})$ that restricts to a generator $H^k(E_x, E_x \setminus i(x); \mathbb{Z}) \cong H^k(\mathbb{R}^k; \mathbb{R}^k \setminus \{0\}; \mathbb{Z}) \cong \mathbb{Z}$ for all $x \in S$. The microbundle ξ together with a Thom class is called an *oriented microbundle*.

A Thom class of a topological \mathbb{R}^n bundle over S is by definition a Thom class of the underlying microbundle.

Remark 10.13. As in the smooth case, consider the orientation bundle $\pi: \text{Or}(\xi) \rightarrow S$ with fibre over $x \in S$ the discrete set $\text{Or}(\xi)_x = \{\text{primitive classes of } H_k(E_x, E_x \setminus i(x); \mathbb{Z})\}$. This is a $\mathbb{Z}/2$ -principal bundle, and a Thom class $\tau(\xi)$ determines a global section $s \in \Gamma(\text{Or}(\xi))$ by enforcing $\langle \tau(\xi), s(x) \rangle = 1$ for every $x \in S$. By the same equation, a global section $\Gamma(\text{Or}(\xi))$ determines a Thom class.

Remark 10.14. Let X be an oriented manifold. Let S be a submanifold with normal microbundle νS . A standard argument, similar to the construction of the fundamental class of an oriented manifold, shows that an orientation of S determines a unique Thom class of νS compatible with the ambient orientation and vice versa.

To prove Theorem 10.11, we will consider a map $f: X \rightarrow Y$ that is transverse to an oriented submanifold S of Y . By Remark 10.14, νS is an oriented microbundle carrying the Thom class τ . Note as $f: \nu f^{-1}(S) \rightarrow f^*\nu S$ is an isomorphism, also $f^*\tau$ is a Thom class of $\nu f^{-1}(S)$ and we orient $f^{-1}(S)$ accordingly. Before we proceed with the proof, we recall the following compatibility between Thom classes and Poincaré duality [Bre97, Definition VI.11.1, Corollary VI.11.6], interpreted for microbundles.

Lemma 10.15. *Let X be a compact oriented n -manifold with fundamental class $[X] \in H_n(X, \partial X)$, and let $i: S \rightarrow X$ be an oriented proper k -dimensional submanifold of X with normal microbundle νS . The composition*

$$H^{n-k}(\nu S, \nu S \setminus i(S)) \xleftarrow[\cong]{\text{Exc.}} H^{n-k}(X, X \setminus S) \rightarrow H^{n-k}(X) \xrightarrow{\text{PD}_X} H_k(X, \partial X)$$

maps the Thom class τ of νS , that is determined by the orientations of X and S , to the fundamental class $i_[S]$.*

PROOF. We start out with a general piece of notation. Given an oriented n -dimensional topological manifold W and given a compact subset $K \subseteq W \subseteq W$ we write $[W] \in H_n(W, (W \setminus K) \cup \partial W)$ for the unique element which, for each $x \in W \setminus (K \setminus \partial W)$ is sent to the generator of $H_n(W, W \setminus \{x\})$ given by the orientation of W . Recall that if W is compact, then Poincaré duality map PD_W is $\cap[W]$. (We refer to [Dol95, Section VII.12] for the definition and precise nature of the cap product on relative (co-) homology.)

We set $d := n - k$ and make a few preliminary observations.

- (1) It suffices to prove the lemma for connected S .
- (2) We identify S with $i(S)$.
- (3) By Kister's Theorem 6.17 there exists an open subset $E \subseteq \nu S$ containing S such that $p: E \rightarrow B$ is a projection map of a topological \mathbb{R}^d -bundle whose 0-section is i .
- (4) Pick an $x \in S \setminus \partial S$ and a closed k -disc $U \subseteq S \setminus \partial S$ containing x such that there is a local trivialisation $\Phi: p^{-1}(U) \rightarrow U \times \mathbb{R}^d$ for the \mathbb{R}^d -bundle in a neighbourhood of x . We can and will choose Φ such that it preserves the orientation of the fibres.
- (5) We let $j: U \rightarrow U \times \{0\}$ and $k: U \times \mathbb{R}^d \rightarrow \mathbb{R}^d$ denote the obvious maps.

Next we consider the following diagram.

$$\begin{array}{ccccc}
H^d(X, X \setminus S) & \xrightarrow{\cap [X]} & H_k(X, \partial X) & & \\
\cong \downarrow \text{Exc.} & & \uparrow & & \\
H^d(\nu S, \nu S \setminus S) & \xrightarrow{\cap [\nu S]} & H_k(\nu S, \partial \nu S) & & \\
\cong \downarrow \text{Exc.} & & \uparrow & & \\
H^d(E, E \setminus S) & \xrightarrow{\cap [E]} & H_k(E, \partial E) & \xleftarrow[\cong]{i_*} & H_k(S, \partial S) \\
\cong \downarrow \text{Id} & & \downarrow & & \downarrow \cong \\
H^d(E, E \setminus S) & \xrightarrow{\cap [E]} & H_k(E, E \setminus p^{-1}(x)) & \xleftarrow[\cong]{i_*} & H_k(S, S \setminus x) \\
\cong \downarrow \text{Exc.} & & \text{Exc.} \uparrow \cong & & \text{Exc.} \uparrow \cong \\
H^d(p^{-1}(U), p^{-1}(U) \setminus S) & \xrightarrow{\cap [p^{-1}(U)]} & H_k(p^{-1}(U), p^{-1}(U) \setminus x) & \xleftarrow[\cong]{i_*} & H_k(U, U \setminus x) \\
\Phi^* \uparrow \cong & & \cong \downarrow \Phi_* & \swarrow j_* & \\
H^d(U \times \mathbb{R}^d, U \times (\mathbb{R}^d \setminus \{0\})) & \xrightarrow[\cong]{\cap [U \times \mathbb{R}^d]} & H_k(U \times \mathbb{R}^d, (U \setminus \{x\}) \times \mathbb{R}^d) & & \\
k^* \uparrow \cong & & & & \\
H^d(\mathbb{R}^d, \mathbb{R}^d \setminus \{0\}) & & & &
\end{array}$$

The bottom vertical maps are given by the local trivialisation Φ . All other vertical maps are the obvious maps of pairs of topological spaces. The maps decorated with “Exc.” are isomorphisms by the excision theorem. The horizontal maps i_* to the right are isomorphisms by the Serre spectral sequence (note that a priori we do not know whether $i: S \rightarrow E$ is a homotopy equivalence.) Note that bottom vertical map is an isomorphism since $\{x\}$ is a deformation retract of the disc U .

We make the following observations.

- (1) By definition of the Thom class, it is sent, under the left vertical maps, to the standard generator $[\mathbb{R}^d]^* \in H^d(\mathbb{R}^d, \mathbb{R}^d \setminus \{0\})$.
- (2) It follows from the compatibility of the cross product with the cap product [Dol195, Section VII.12.17] (see also [Fri23, Proposition 138.2]) that

$$k^*([\mathbb{R}^d]^*) \cap [U \times \mathbb{R}^d] = j_*([S]).$$

Hence the Thom class in $H^d(\nu S, \nu S \setminus S)$ and the fundamental class $[S, \partial S] \in H_k(S, \partial S)$ have the same image in $H_k(U \times \mathbb{R}^d, (U \setminus \{x\}) \times \mathbb{R}^d)$.

Standard facts about the relative cap product show that the diagram commutes. In particular note that the map $H_k(E, \partial E) \rightarrow H_k(E, E \setminus p^{-1}(x))$ is an isomorphism. It is then not too hard to chase the diagram to deduce that the Thom class and $[S, \partial S]$ also have the same image in $H_k(X, \partial X)$, and so the lemma holds. \square

PROOF OF THEOREM 10.11. First we let $k = n - 1$. Let $\alpha \in H^1(X; \mathbb{Z})$ be the Poincaré dual to $\sigma \in H_{n-1}(X, \partial X)$. Recall that in Theorem 3.16 we showed that X is homotopy

equivalent to a CW complex. Therefore we have the following correspondence between homotopy classes of maps to Eilenberg-MacLane spaces and cohomology classes of X :

$$\begin{aligned} [X, S^1] &= [X, K(\mathbb{Z}, 1)] \xrightarrow{\cong} H^1(X; \mathbb{Z}) \\ &f \mapsto f^*\theta, \end{aligned}$$

where θ is the Hom dual of the fundamental class of S^1 . Note that we used here that X is homotopy equivalent to a CW complex. Pick an arbitrary point $\text{pt} \in S^1$ and denote a tubular neighbourhood by $\nu(\text{pt})$. Note that the Thom class τ_{pt} for $\nu(\text{pt})$ is mapped under $H^1(S^1, S^1 \setminus \text{pt}) \rightarrow H^1(S^1)$ to $\theta = \text{PD}_{S^1}^{-1}[\text{pt}]$. Let $f: X \rightarrow S^1$ be a map corresponding to α , so $f^*\theta = \alpha$. Make f transverse to a tubular neighbourhood of $\text{pt} \in S^1$ using Theorem 10.10. Consequently, $S := f^{-1}(\text{pt})$ is an $(n-1)$ -dimensional submanifold of X . By definition, f induces a bundle isomorphism $f: \nu S \rightarrow f^*\nu(\text{pt})$. We have, as elements in $H^1(X; \mathbb{Z})$, that

$$\alpha = f^*\theta = f^* \text{PD}_{S^1}^{-1}[\text{pt}] = f^*\tau_{\text{pt}}.$$

Note that $f^*\tau_{\text{pt}}$ is the image of the Thom class of νS under the first two maps in the composition displayed in the statement of Lemma 10.15. Lemma 10.15 thus tells us the last equality of:

$$\alpha = f^*\tau_{\text{pt}} = \tau_S = \text{PD}_X^{-1}[S] \in H^1(X; \mathbb{Z}).$$

We have thus shown that $[S] = \text{PD}_X(\alpha) = \sigma$.

A similar proof works for the codimension two case, i.e. $k = n - 2$. Let $\alpha \in H^2(X; \mathbb{Z})$ be Poincaré dual to $\sigma \in H_{n-2}(X, \partial X)$. Recall that

$$\begin{aligned} [X, \mathbb{C}P^\infty] &= [X, K(\mathbb{Z}, 2)] \xrightarrow{\cong} H^2(X; \mathbb{Z}) \\ &f \mapsto f^*\theta, \end{aligned}$$

where $\theta \in H^2(\mathbb{C}P^\infty; \mathbb{Z})$ is the Hom dual of the fundamental class of $\mathbb{C}P^1 \subseteq \mathbb{C}P^\infty$. Since X is homotopy equivalent to an n -dimensional CW complex we can homotope a given representing map f to have image in $\mathbb{C}P^m \subseteq \mathbb{C}P^\infty$, where $m = \lfloor n/2 \rfloor$. We abuse notation and denote the image of θ under the restriction map $H^2(\mathbb{C}P^\infty; \mathbb{Z}) \rightarrow H^2(\mathbb{C}P^m; \mathbb{Z})$ also by θ .

Let $f: X \rightarrow \mathbb{C}P^m$ be a map corresponding to α , so $f^*\theta = \alpha$. Make f transverse to a normal microbundle of $\mathbb{C}P^{m-1} \subseteq \mathbb{C}P^m$ using Theorem 10.10. The inverse image $S := f^{-1}(\mathbb{C}P^{m-1})$ is an $(n-2)$ -dimensional submanifold of X .

Let $\tau_{\mathbb{C}P^{m-1}} \in H^2(\mathbb{C}P^m, \mathbb{C}P^m \setminus \mathbb{C}P^{m-1})$ denote a Thom class for the normal bundle $\nu\mathbb{C}P^{m-1}$. By Lemma 10.15, the map $H^2(\mathbb{C}P^m, \mathbb{C}P^m \setminus \mathbb{C}P^{m-1}; \mathbb{Z}) \rightarrow H^2(\mathbb{C}P^m; \mathbb{Z})$ sends $\tau_{\mathbb{C}P^{m-1}}$ to $\theta = \text{PD}_{\mathbb{C}P^m}^{-1}[\mathbb{C}P^{m-1}]$.

By definition of S and νS , the map $f: X \rightarrow \mathbb{C}P^m$ induces a bundle isomorphism $f: \nu S \rightarrow f^*\nu\mathbb{C}P^{m-1}$. We have, as elements in $H^2(X; \mathbb{Z})$, that

$$\alpha = f^*\theta = f^* \text{PD}_{\mathbb{C}P^m}^{-1}[\mathbb{C}P^{m-1}] = f^*\tau_{\mathbb{C}P^{m-1}}.$$

Note that $f^*\tau_{\mathbb{C}P^{m-1}}$ is the image of the Thom class of νS under the first two maps in the composition displayed in the statement of Lemma 10.15. Thus Lemma 10.15 gives the last

equality of:

$$\alpha = f^* \tau_{\mathbb{C}P^{m-1}} = \tau_S = \text{PD}_X^{-1}[S] \in H^2(X; \mathbb{Z}).$$

We have now shown that $[S] = \text{PD}_X(\alpha) = \sigma$. \square

Remark 10.16. We have seen that $f^![\text{pt}] = \text{PD}_X \circ f^* \circ \text{PD}_{S^1}^{-1}[\text{pt}] = [f^{-1}(\text{pt})]$, when f is transverse to pt , and similarly that $f^![\mathbb{C}P^{m-1}] = f^{-1}(\mathbb{C}P^{m-1})$.

Next we offer the following promised refinement of Theorem 10.11 in the 4-dimensional case, together with an alternative proof that uses smoothing away from a point.

Theorem 10.17. *Let X be a compact orientable 4-manifold and let A be a union of components of ∂X . Let $k = 2$ or $k = 3$ and let $\sigma \in H_k(X, A; \mathbb{Z})$.*

- (1) *The class σ can be represented by a k -dimensional submanifold Y with $\partial Y \subseteq A$.*
- (2) *In the case $k = 3$, the boundary of Y can be specified: if $B \subseteq A$ is an oriented closed 2-dimensional smooth submanifold contained in A such that $\partial(\sigma) = [B] \in H_2(A; \mathbb{Z})$, then σ can be represented by an oriented compact 3-dimensional submanifold Y with $\partial Y = B$.*

The submanifold B can be assumed to be smooth, since ∂X is a 3-manifold and so has a unique smooth structure by [Moi52], [Moi77, p. 252–253].

Note that Theorem 10.17 also holds for $k = 0$ and $k = 1$. This is trivial for $k = 0$. To see this for $k = 1$, remove a point from each connected component to get a smooth 4-manifold by Theorem 9.1. Note that $H_1(X \setminus \{\text{pt}\}, A; \mathbb{Z}) \cong H_1(X, A; \mathbb{Z})$. Then by smooth approximation and general position, every 1-dimensional homology class can be represented by a 1-dimensional submanifold of X .

Example 10.18.

- (1) If we apply the theorem to $A = \partial X$, we see that any homology class in $H_2(X, \partial X; \mathbb{Z})$ and $H_3(X, \partial X; \mathbb{Z})$ can be represented by a properly embedded submanifold. For $A = \emptyset$ we obtain the analogous statement for absolute homology groups.
- (2) Let F be a properly embedded 2-dimensional submanifold of D^4 and let S be a surface in $\partial D^4 = S^3$ with $\partial S = \partial F$. Consider the 4-manifold $X := D^4 \setminus \nu F$. In the boundary of X we have the surface $B = S \cup F \times \{1\}$. It follows from the long exact sequence of the pair $(X, \partial X)$ and Poincaré duality that the map $H_3(X, \partial X; \mathbb{Z}) \rightarrow H_2(\partial X; \mathbb{Z})$ is an epimorphism. It follows from Theorem 10.17, applied to $A = \partial X$, that there exists a 3-dimensional submanifold Y of $D^4 \setminus \nu F$ with $\partial Y = B$. This statement is folklore, and a proof using topological transversality for maps was written down by Lewark-McCoy [LM19].

For $n = 4$ the statement of the following theorem is precisely the statement of Theorem 10.17 in the smooth category.

Proposition 10.19. *Let X be a compact, orientable, smooth n -manifold and let A be a union of components of ∂X . Let $\ell = 1$ or $\ell = 2$ and let $\sigma \in H_{n-\ell}(X, A; \mathbb{Z})$. Then the following holds.*

- (1) The class σ is represented by an $(n - \ell)$ -dimensional smooth orientable submanifold Y with $\partial Y \subseteq A$.
- (2) Suppose $\ell = 1$ and suppose we are given a closed, oriented $(n - 2)$ -dimensional smooth submanifold B of A such that $\partial(\sigma) = [B] \in H_{n-2}(A)$. Then σ is represented by an oriented compact $(n - 1)$ -dimensional smooth submanifold Y with $\partial Y = B$.

Example 10.20. Let $K \subseteq S^3$ be a knot. We write $X = S^3 \setminus \nu K$. Let $\lambda \subseteq \partial X$ be a longitude of K , i.e. λ is a curve that represents a generator of $\ker(H_1(\partial X; \mathbb{Z}) \rightarrow H_1(X; \mathbb{Z}))$. There exists a homology class $\sigma \in H_2(X, \partial X; \mathbb{Z})$ with $\partial(\sigma) = [\lambda] \in H_1(\partial X; \mathbb{Z})$. It follows from Proposition 10.19 that there exists an orientable surface F in X with $\partial F = K$.

PROOF. Let X be a compact orientable smooth n -manifold and let A be a union of components of ∂X . First we prove statements (1) and (2) for the case $\ell = 1$. Let $\sigma \in H_{n-1}(X, A; \mathbb{Z})$.

- (1) Write $\tilde{A} = \partial X \setminus A$. Let $\text{PD}: H^1(X, \tilde{A}; \mathbb{Z}) \rightarrow H_{n-1}(X, A; \mathbb{Z})$ be the Poincaré duality isomorphism. We have $H^1(X, \tilde{A}; \mathbb{Z}) \cong [X/\tilde{A}, S^1]$ and any such class can be represented by a continuous map $\varphi: X \rightarrow S^1$ that is constant on \tilde{A} , and uniquely determined up to homotopy rel. \tilde{A} . We can and shall homotope φ to a smooth map. Furthermore, arrange that $-1 \in S^1$ is a regular value of φ . Then $Y := \varphi^{-1}(-1)$ is an $(n - 1)$ -dimensional submanifold whose boundary lies on $\partial X \setminus \tilde{A}$, that is the boundary lies on A . The manifold $Y = \varphi^{-1}(-1)$ satisfies $[Y] = \sigma$ (this follows from Lemma 10.15, as explained in the proof of Theorem 10.11).
- (2) Now suppose that we are given an oriented closed $(n - 2)$ -dimensional submanifold B of A such that $\partial(\sigma) = [B] \in H_{n-1}(A)$. Pick a collar neighbourhood $\partial X \times [0, 1]$ and choose a continuous map $\varphi: X \setminus (\partial X \times [0, 1]) \rightarrow S^1$ as above. Also choose a tubular neighbourhood $B \times [-1/2, 1/2]$ of B in A . Consider the map sending (b, t) to $e^{\pi i(t-1)}$, $b \in B, t \in [-1/2, 1/2]$ and extend it to a smooth map $\psi: A \rightarrow S^1$ by sending all other points into $\{e^{\pi i t} \in S^1 \mid t \in [-2/3, 2/3]\}$. Since $\partial(\sigma) = [B] \in H_{n-2}(A) \cong H^1(A; \mathbb{Z})$, we see that the restriction of φ to $A \times \{1\} = A$ is homotopic to $\psi: A \rightarrow S^1$. Therefore, using this homotopy in the interval $[\frac{1}{2}, 1]$, we can extend φ to a function on X that restricts to ψ on each $A \times \{s\}$ with $s \in [0, \frac{1}{2}]$. Finally, smoothen φ without changing it on $A \times [0, \frac{1}{4}]$ to obtain a smooth map $X \rightarrow S^1$ in the same homotopy class. This is possible since the original φ was already smooth on $A \times [0, \frac{1}{2}]$. Put differently, the new smooth map $\varphi: X \rightarrow S^1$ restricts to ψ on $A = A \times \{0\}$.

Note that -1 is a regular value of ψ , and by changing φ outside $A \times [0, \frac{1}{4}]$, we can also arrange -1 to be a regular value of φ . The manifold $Y = \varphi^{-1}(-1)$ satisfies $[Y] = \sigma$ (as in (1) this follows from Lemma 10.15 below) and $\partial Y = B \times \{0\} = B$.

For $\ell = 2$ the argument is similar: we have to replace the argument using S^1 by the argument of [GS99, Proposition 1.2.3]. Recall from Theorem 3.16 that X is homotopy equivalent to a finite CW complex. Therefore, we represent a codimension 2 homology class $\sigma \in H_{n-2}(X, \partial X; \mathbb{Z}) \cong H^2(X; \mathbb{Z})$ by a map $X \rightarrow \mathbb{C}\mathbb{P}^\infty$, and homotope into the k -skeleton

to a map $f: X \rightarrow \mathbb{C}\mathbb{P}^k$ for $k \geq 2$. Now arrange f to be transverse to the codimension 2 submanifold $\mathbb{C}\mathbb{P}^{k-1} \subseteq \mathbb{C}\mathbb{P}^k$. The desired submanifold is the preimage $Y = f^{-1}(\mathbb{C}\mathbb{P}^{k-1})$. We leave further details to the reader. Again the argument is similar to that in the proof of Theorem 10.11. \square

Lemma 10.21. *Let W be a smooth n -manifold and let C be a compact subset. There exists a compact smooth n -dimensional submanifold X of W that contains C .*

PROOF. By the Whitney Embedding Theorem (see e.g. [Lee11, Theorem 6.15]), there exists a proper embedding $f: W \rightarrow \mathbb{R}^{2n+1}$. Recall that in this context proper means that the preimage of a compact set is compact. Pick a point $P \in \mathbb{R}^{2n+1}$ that does not lie in the image of f . Denote the Euclidean distance to the point P by $d: \mathbb{R}^{2n+1} \rightarrow \mathbb{R}_{\geq 0}$. This map is smooth outside P , so in particular $d \circ f: W \rightarrow \mathbb{R}_{\geq 0}$ is smooth. Since C is compact, there exists an $r \in \mathbb{R}_{\geq 0}$ such that $(d \circ f)(C) \subseteq [0, r]$. By Sard's Theorem, there exists a regular value $x > r$. Then $X := (d \circ f)^{-1}([0, x])$ has the desired properties. \square

PROOF OF THEOREM 10.17. Let M be a compact orientable connected 4-manifold and let A be a union of components of ∂M . Let $k = 2$ or $k = 3$ and let $\sigma \in H_k(X, A; \mathbb{Z})$.

Pick a point $P \in M \setminus \partial M$ and pick an open ball $B \subseteq M \setminus \partial M$ containing P . It follows from a Mayer-Vietoris argument applied to $M = (M \setminus \{P\}) \cup B$ that the inclusion induced map $H_k(M \setminus \{P\}, A) \rightarrow H_k(M, A)$ is an isomorphism for $k = 2, 3$.

Now let $\sigma \in H_k(M, A)$. By the previous paragraph we can view σ as an element in $H_k(M \setminus \{P\}, A)$. By Theorem 9.1 the manifold $M \setminus \{P\}$ is smooth. There exists a compact subset K of $M \setminus \{P\}$ such that σ lies in the image of $H_k(K, A) \rightarrow H_k(M \setminus \{P\}, A)$, since one can take the union of the images of the singular simplices in a singular chain representing σ . By Lemma 10.21, there exists a compact 4-dimensional smooth submanifold X of $M \setminus \{P\}$ that contains the compact set $K \cup \partial M$. Note that A is again a union of components of ∂X . The desired statement of Theorem 10.17 is now an immediate consequence of Proposition 10.19 (1), with σ the image of $\sigma \in H_k(K, A)$ under the inclusion induced map to $H_k(X, A)$. \square

Tubing of surfaces

As an example of the use of the technology we have discussed thus far, we show that one can tube together two locally flat embedded surfaces in a 4-manifold, to obtain an embedding of the connected sum. This operation is standard in the smooth category, but as ever in the topological category one should take some care.

The following situation is by no means the most general such result possible. We wish to illustrate two things. First, that operations on surfaces that can be performed in the smooth category can usually also be performed in general 4-manifolds with locally flat surfaces (although performing these operations in a parametrised way seems to be beyond current knowledge). Second, we want to show the level of detail required to demonstrate that such operations work.

Proposition 11.1. (Tubing Theorem) *Let S and T be 2-dimensional proper submanifolds of a connected 4-manifold M , that is S and T are locally flat embedded surfaces. Pick a point $P \in S \setminus \partial S$ and $Q \in T \setminus \partial T$. Let $[\gamma] \in H_1(M, \{P, Q\}; \mathbb{Z})$ be a relative homology class. There is a locally flat embedded arc C joining P and Q , satisfying the following.*

- (i) *We have $[C] = [\gamma] \in H_1(M, \{P, Q\}; \mathbb{Z})$.*
- (ii) *The interior of C is disjoint from $S \cup T$.*
- (iii) *The arc C extends to a neighbourhood $C \times D^2$ embedded in M such that $E_S := \{P\} \times D^2 \subseteq S$ and $E_T := \{Q\} \times D^2 \subseteq T$.*
- (iv) *We have $(C \times D^2) \setminus (E_S \cup E_T) \subseteq M \setminus (S \cup T)$.*
- (v) *The intersection of $C \times D^2$ with a normal disc bundle $D(S)$ of S is such that for every d , $(C \times \{d\}) \cap D(S)$ is a ray in a single fibre of $D(S)$, and similarly for T . Moreover there is a trivialisation of the normal vector bundle over E_S as $E_S \times D^2$ such that for every $c \in C$ with $(\{c\} \times D^2) \cap D(S) \neq \emptyset$, we have that $\{c\} \times D^2 = E_S \times \{e\}$ for some $e \in D^2$, and all such e that arise this way lie on a fixed ray from the origin of D^2 .*

These data allow us to perform tubing of surfaces ambiently.

Proposition 11.2. *Given data $S, T, C \times D^2, E_S$ and E_T as in Proposition 11.1, the subset*

$$(S \setminus \frac{1}{2}E_S) \cup (T \setminus \frac{1}{2}E_T) \cup C \times \frac{1}{2}S^1$$

is a 2-dimensional submanifold abstractly homeomorphic to $S \# T$.

PROOF. The surfaces and the tube are locally flat by assumption, or by construction from Proposition 11.1. The circles where the tube is glued to the surface are locally flat

points. To see this observe that we have arranged a coordinate system in which this gluing is a completely standard attachment at angle $\pi/2$. \square




FIGURE 11.1. Illustration of Proposition 11.1.

PROOF OF PROPOSITION 11.1. Since S and T are proper submanifolds, they have normal vector bundles by Theorem 5.14. Pick normal disc bundles $D(S)$ and $D(T)$, and remove the interiors of $\frac{1}{2}D(S)$ and $\frac{1}{2}D(T)$ i.e. smaller disc bundles inside the normal disc bundles. We obtain a manifold with boundary

$$X := M \setminus (\text{Int } \frac{1}{2}D(S) \cup \text{Int } \frac{1}{2}D(T))$$

together with a collar neighbourhood of the boundary arising from $D(S) \setminus \text{Int } \frac{1}{2}D(S)$, and the same with T replacing S , extended using Theorem 2.16 to a collar neighbourhood for all of ∂X . Choose a closed disc neighbourhood E_S of P in S . We write $\partial_S X$ for the fibrewise boundary of $\frac{1}{2}D(S)$, $\partial_T X$ for the fibrewise boundary of $\frac{1}{2}D(T)$, and $\partial_1 X$ for $\partial_S X \cup \partial_T X = \overline{\partial X} \setminus \partial M$.

Choose a trivialisation of the normal vector bundle νS in a neighbourhood $N(E_S)$ of E_S , as $N(E_S) \times D^2$. A ray in D^2 from the origin to the boundary determines an embedding $E_S \times [0, 1] \subseteq \frac{1}{2}D(S)$. We obtain in particular a disc $E_S \times \{1\} \in N(E_S) \times \{\text{pt}\} \subseteq N(E_S) \times S^1$. Choose a smooth structure on ∂X (which we may do since ∂X is a 3-manifold), and choose a smoothly embedded neighbourhood $F_S \cong D^3$ in $\partial_S X$ that contains $E_S \times \{1\}$ in its interior.

Make the analogous set of choices and constructions for T , to obtain E_T , $N(E_T)$, $E_T \times [0, 1] \subseteq \frac{1}{2}D(T)$, and $F_T \cong D^3$ in $\partial_T X$ that contains $E_T \times \{1\}$ in its interior.

Remove a point r from X , and using Theorem 9.1 choose a smooth structure on $X \setminus \{r\}$ extending the chosen smooth structure on ∂X . Choose a smoothly embedded path $C_X \subseteq X$ between the centres of $E_S \times \{1\}$ and $E_T \times \{1\}$, such that C_X extends along the previously chosen rays inside the normal vector bundles to a path C between P and Q such that $[C] = [\gamma] \in H_1(M, \{P, Q\}; \mathbb{Z})$. Extend C_X to a codimension zero submanifold $N(C_X)$ homeomorphic to $I \times D^3$, with $I \times \{0\} \subseteq I \times D^3$ mapping to C_X , and such that $\{0\} \times D^3$ maps to $F_S \subseteq \partial_S X$ and $\{1\} \times D^3$ maps to $F_T \subseteq \partial_T X$.

Now, for small ε , $[0, \varepsilon] \times D^3$ and $[1 - \varepsilon, 1] \times D^3$ give rise to collar neighbourhoods of the closed subsets F_S and F_T of $\partial_1 X$. Use Theorem 2.16 to extend this collar neighbourhood to a collar neighbourhood over all of ∂X .




FIGURE 11.2. Illustration for the proof of Proposition 11.1.

We now have two collar neighbourhoods of ∂X , the collar $\Psi_1: \partial X \times [0, 1] \hookrightarrow X$ we have just constructed which is compatible with $N(C_X)$, and the collar neighbourhood $\Psi_2: \partial X \times [0, 1] \hookrightarrow X$ constructed above from $D(S) \setminus \text{Int } \frac{1}{2}D(S)$ and $D(T) \setminus \text{Int } \frac{1}{2}D(T)$. By Theorem 2.18, there is an isotopy $H_t: M \rightarrow M$ starting from the identity, such that $H_1 \circ \Psi_1 = \Psi_2$, i.e. sends the first collar to the second.

We now obtain a codimension zero submanifold $C_X \times D^3$ homeomorphic to $I \times D^3$ such that, with respect to the collar neighbourhood Ψ_2 , we have:

- For all $c \in C_X$ such that $\{c\} \times D^3 \cap \Psi_2(\partial X \times [0, 1]) \neq \emptyset$, we have that $\{c\} \times D^3 \subseteq \Psi_2(\partial X \times [t, 1])$ for some $t \in [0, 1]$.
- For every $d \in D^3$, $(C_X \times \{d\}) \cap (\partial X \times [0, 1]) = \Psi_2(\{x\} \times [0, 1])$ for some x in either F_S or F_T .

In addition, above we constructed two discs $E_S \subseteq F_S$ and $E_T \subseteq F_T$. Any two embedded discs in a 3-ball are ambiently isotopic: place this isotopy inside $C_X \times D^3$ to obtain a locally flat embedding $C_X \times D^2 \cong I \times D^2 \subseteq C_X \times D^3$.

Now consider $X \subseteq M$ and take the union

$$(E_S \times [0, 1]) \cup (C_X \times D^2) \cup (E_T \times [0, 1]) \subseteq M$$

to obtain an embedding $C \times D^2 \cong I \times D^2$ whose intersection with S equals E_S and whose intersection with T equals E_T . The core $C = C \times \{0\}$ is a locally flat embedded path in M from P to Q with interior in $M \setminus (S \cup T)$ and with the correct relative homology class in $H_1(M, \{P, Q\}; \mathbb{Z})$. We may then perform the tubing $S \# T := (S \setminus E_S) \cup (T \setminus E_T) \cup C \times S^1$ as promised. \square

CHAPTER 12

Classification results for 4-manifolds

It is well-known (e.g. [CZ90, Theorem 5.1.1]) that any finitely presented group is the fundamental group of a closed orientable smooth 4-manifold. Markov [Mar58] used this fact to show that closed 4-manifolds cannot be classified up to homeomorphism. To circumvent this group theoretic issue one aims to classify 4-manifolds with a given isomorphism type of a fundamental group.

In this chapter we present the known 4-manifold classification results that have been obtained using the techniques of classical, or modified, surgery theory in the topological category, combined with Freedman’s Disc Embedding Theorem [Fre82, FQ90, BKK⁺21]. The use of this theorem requires the fundamental group of the 4-manifold be “good” [FQ90, Part II, Introduction], a condition that has a precise geometric description using the “ π_1 -null disc property”. We will not reproduce that description here, but will instead note which groups are currently known to be good. Freedman showed that the infinite cyclic group and finite groups are good [Fre82, pp. 658-659] (see also [FQ90, Section 5.1]). In addition, by [FT95, Lemma 1.2] the class of good groups is closed under extensions, direct limits, subgroups and quotients. It follows that solvable groups are good. Furthermore in [FT95, Theorem 0.1] and [KQ00] it was shown that groups with subexponential growth are good.

12.1. SIMPLY CONNECTED 4-MANIFOLDS

The following theorem was the first noteworthy result towards a classification of 4-dimensional manifolds.

Theorem 12.1. *Suppose M and N are two closed oriented simply-connected 4-dimensional manifolds. If the intersection forms are isometric, then M and N are homotopy equivalent.*

PROOF. This theorem was proved for smooth manifolds by Milnor [Mil58, Theorem 3], building on work of Whitehead [Whi49]. A proof that works in the general case is given in [MH73, Chapter V, Theorem 1.5]. \square

We state Freedman’s classification for closed, simply connected 4-manifolds [Fre82, Theorem 1.5]. We give the statement as in [FQ90, Theorem 10.1]. We note that Freedman’s original statement only applied to 4-manifolds smoothable away from a point, and so the statement below requires the subsequent developments by Quinn. The last sentence comes from [Qui86].

Theorem 12.2. *Fix a triple (F, θ, k) , where F is a finitely generated free abelian group, θ is a symmetric, nonsingular, bilinear form $\theta: F \times F \rightarrow \mathbb{Z}$, and $k \in \mathbb{Z}/2$. If θ is even, that is $\theta(x, x) \in 2\mathbb{Z}$ is even for every $x \in F$, then suppose that $\sigma(\theta)/8 \equiv k \in \mathbb{Z}/2$.*

Then there exists a closed, simply connected, oriented 4-manifold M with $H_2(M; \mathbb{Z}) \cong F$, with intersection form isometric to θ and with Kirby-Siebenmann invariant equal to k .

Let M and M' be two closed, simply connected, oriented 4-manifolds and let $\phi: H_2(M; \mathbb{Z}) \xrightarrow{\cong} H_2(M'; \mathbb{Z})$ be an isometry of the intersection forms. Suppose that $\text{ks}(M) = \text{ks}(M')$. Then there is an orientation preserving homeomorphism $M \xrightarrow{\cong} M'$ inducing ϕ on second homology. This homeomorphism is unique up to isotopy.

In other words, every even, symmetric, integral matrix with determinant ± 1 is realised as the intersection form of a unique closed, simply connected, oriented 4-manifold. For such matrices which are odd instead, there are precisely two closed, simply connected, oriented 4-manifolds up to homeomorphism, exactly one of which has vanishing Kirby-Siebenmann invariant and is therefore stably smoothable. These two manifolds are homotopy equivalent by Theorem 12.1.

In particular, the last paragraph with $M = M'$ implies that every automorphism of the intersection form of a closed, simply connected, oriented 4-manifold is realised by a self-homeomorphism of M .

The following special case of Theorem 12.2, when $F = 0$, is worth pointing out explicitly.

Corollary 12.3. (4-dimensional Poincaré conjecture) *If N is a 4-manifold homotopy equivalent to S^4 then N is homeomorphic to S^4 .*

PROOF. Note N and S^4 are closed, simply connected, and oriented. Furthermore, $H_2(N; \mathbb{Z}) = 0$ and the zero map $H_2(N; \mathbb{Z}) \rightarrow H_2(S^4; \mathbb{Z})$ is an isometry (between zero forms). By the last paragraph of Theorem 12.2, there is a homeomorphism $N \cong S^4$ realising this isometry. Note that since N has trivial and therefore even intersection form, $\text{ks}(N) = \sigma(N)/8 = 0$ by Theorem 9.2 (4). \square

12.2. NON SIMPLY-CONNECTED 4-MANIFOLDS

We summarise known classification results for different types of nontrivial fundamental groups.

12.2.1. Infinite cyclic group. First, we present a classification result [FQ90, Theorem 10.7A] for closed, oriented 4-manifolds with fundamental group \mathbb{Z} which is quite similar to Theorem 12.2. To state the theorem we need some extra definitions.

Definition 12.4. For a finitely generated free $\mathbb{Z}[\mathbb{Z}]$ module F , a hermitian sesquilinear form $\theta: F \times F \rightarrow \mathbb{Z}[\mathbb{Z}]$ is called *even* if there is a left $\mathbb{Z}[\mathbb{Z}]$ -module homomorphism $q: F \rightarrow \overline{\text{Hom}_{\mathbb{Z}[\mathbb{Z}]}(F, \mathbb{Z}[\mathbb{Z}])}$ with $q + q^*: F \rightarrow \overline{\text{Hom}_{\mathbb{Z}[\mathbb{Z}]}(F, \mathbb{Z}[\mathbb{Z}])}$ equal to the adjoint of θ . Otherwise we call the form *odd*.

Definition 12.5. Two homeomorphisms $h_0, h_1: M \rightarrow N$ are *pseudo-isotopic* if there is a homeomorphism $H: M \times I \rightarrow N \times I$ with $H|_{M \times \{i\}} = h_i: M \times \{i\} \rightarrow N \times \{i\}$ for $i = 0, 1$.

An isotopy of homeomorphisms gives rise to a pseudo-isotopy. Perron and Quinn proved that the converse holds for compact simply connected 4-manifolds [Per86], [Qui86]). Budney and Gabai [BG23] showed that pseudo-isotopy does not in general imply isotopy for homeomorphisms between 4-manifolds with nontrivial fundamental groups.

Theorem 12.6. *Fix a triple (F, θ, k) , where F is a finitely generated free $\mathbb{Z}[\mathbb{Z}]$ -module, θ is a hermitian, nonsingular, sesquilinear form $\theta: F \times F \rightarrow \mathbb{Z}[\mathbb{Z}]$, and $k \in \mathbb{Z}/2$. If θ is even, then suppose that $\sigma(\mathbb{R} \otimes \theta)/8 \equiv k \in \mathbb{Z}/2$.*

Then there exists a closed, oriented 4-manifold M with $\pi_1(M) \cong \mathbb{Z}$, with $H_2(M; \mathbb{Z}[\mathbb{Z}])$ isomorphic to F , whose equivariant intersection form

$$\lambda_M: H_2(M; \mathbb{Z}[\mathbb{Z}]) \times H_2(M; \mathbb{Z}[\mathbb{Z}]) \rightarrow \mathbb{Z}[\mathbb{Z}]$$

is isometric to θ , and with $\text{ks}(M) = k$.

Let M and M' be two closed, oriented 4-manifolds with $\pi_1(M) \cong \mathbb{Z} \cong \pi_1(M')$ and let $\phi: H_2(M; \mathbb{Z}[\mathbb{Z}]) \xrightarrow{\cong} H_2(M'; \mathbb{Z}[\mathbb{Z}])$ be an isometry of the equivariant intersection forms. Suppose that $\text{ks}(M) = \text{ks}(M')$. Then there is an orientation and basepoint preserving homeomorphism $M \xrightarrow{\cong} M'$ inducing the given identification of the fundamental groups and inducing ϕ on $\mathbb{Z}[\mathbb{Z}]$ coefficient second homology. There are exactly two pseudo-isotopy classes of such homeomorphisms.

The last sentence of this theorem is a correction to [FQ90, Theorem 10.7A] by Stong and Wang [SW00].

12.2.2. Baumslag-Solitar groups. Here is another family of groups for which a complete classification of closed orientable 4-manifolds up to homeomorphism is known. This is the family of solvable Baumslag-Solitar groups

$$B(k) := \langle a, b \mid aba^{-1}b^{-k} \rangle.$$

Note that $B(0) = \mathbb{Z}$ and $B(1) = \mathbb{Z}^2$. Baumslag-Solitar groups are solvable and, as we pointed out above, solvable groups are good. The next classification result was proven by Hambleton, Kreck, and Teichner in [HKT09].

Definition 12.7. The w_2 -type of a closed, oriented 4-manifold M with universal covering \widetilde{M} is type I, II, III, as follows: (I) $w_2(\widetilde{M}) \neq 0$; (II) $w_2(M) = 0$; and (III) $w_2(M) \neq 0$ but $w_2(\widetilde{M}) = 0$.

Theorem 12.8. *Let $B(k)$ be a solvable Baumslag-Solitar group and let M and N be closed, oriented 4-manifolds with fundamental group isomorphic to $B(k)$. Suppose that there is an isomorphism $\phi: H_2(M; \mathbb{Z}[B(k)]) \rightarrow H_2(N; \mathbb{Z}[B(k)])$ of $\mathbb{Z}[B(k)]$ -modules such that:*

- (1) *The map ϕ induces an isometry between the intersection form $\lambda: H_2(M; \mathbb{Z}[B(k)]) \times H_2(M; \mathbb{Z}[B(k)]) \rightarrow \mathbb{Z}[B(k)]$ and the corresponding intersection form on $H_2(N; \mathbb{Z}[B(k)])$.*
- (2) *The Kirby-Siebenmann invariants agree $\text{ks}(M) = \text{ks}(N)$.*
- (3) *The w_2 -types of M and N coincide.*

Then M and N are homeomorphic via an orientation preserving homeomorphism that induces $\phi: H_2(M; \mathbb{Z}[B(k)]) \rightarrow H_2(N; \mathbb{Z}[B(k)])$.

There is also a precise realisation result for these invariants [HKT09, Theorem B] and 4-manifolds with fundamental group $B(k)$.

12.2.3. Finite cohomological dimension. In the same paper as that discussed in the previous section [HKT09], further classification results were given for 4-manifolds with geometrically 2-dimensional fundamental groups.

Some partial results towards a classification for 4-manifolds whose fundamental groups are good and have cohomological dimension 3 appear in Hambleton-Hildum [HH19].

Kasprowski-Land [KL22] studied 4-manifolds M with 4-dimensional fundamental group, under the assumption that the classifying map $M \rightarrow B\pi$ is degree one, i.e. induces an isomorphism $H_4(M; \mathbb{Z}) \xrightarrow{\cong} H_4(B\pi; \mathbb{Z})$.

12.2.4. Finite groups. Next, 4-manifolds with finite fundamental groups were studied by Hambleton and Kreck in [HK88, HK93a]. Given a finitely generated abelian group G , let TG be its torsion subgroup and let $FG := G/TG$.

The most complete result was for 4-manifolds with finite cyclic fundamental group, given below.

Theorem 12.9. *Let G be a finite cyclic group and let M and N be closed, oriented 4-manifolds with fundamental group isomorphic to G . Suppose that there is an isomorphism $\phi: FH_2(M; \mathbb{Z}) \rightarrow FH_2(N; \mathbb{Z})$ such that the following hold.*

- (1) *The map ϕ induces an isometry between the intersection form $\lambda_M: FH_2(M; \mathbb{Z}) \times FH_2(M; \mathbb{Z}) \rightarrow \mathbb{Z}$ and the intersection form $\lambda_N: FH_2(N; \mathbb{Z}) \times FH_2(N; \mathbb{Z}) \rightarrow \mathbb{Z}$.*
- (2) *The Kirby-Siebenmann invariants agree $ks(M) = ks(N)$.*
- (3) *The w_2 -types of M and N coincide.*

Then M and N are homeomorphic via an orientation preserving homeomorphism that induces $\phi: FH_2(M; \mathbb{Z}) \rightarrow FH_2(N; \mathbb{Z})$.

A full realisation result for the invariants in Theorem 12.9 is not known, however Hambleton-Kreck showed how to realise in the majority of cases. The following relations between the invariants hold.

- (1) If $w_2(M) = 0$, then $ks(M) \equiv \sigma(M)/8 \in \mathbb{Z}/2$ and λ_M is even.
- (2) If M is type I, then λ_M is odd.
- (3) If the order of G is odd, then $w_2(\widetilde{M}) = 0$ implies $w_2(M) = 0$, so there are no 4-manifolds with w_2 -type III.

We outline a construction that realises all configurations of the invariants, with the restriction that in w_2 -type III, the intersection form λ_M is even. The key is a construction of rational homology 4-spheres. For every finite cyclic group G , [HK93a, Proposition 4.1] produces the following manifolds.

- (1) A rational homology sphere Σ_G^{II} with w_2 -type II and with fundamental group G . Note that $\text{ks}(\Sigma_G^{II}) \equiv \sigma(\Sigma_G^{II})/8 = 0$.
- (2) A rational homology sphere $\Sigma_G^{III,0}$ with w_2 -type III, trivial Kirby-Siebenmann invariant, and fundamental group G .
- (3) A rational homology sphere $\Sigma_G^{III,1}$ with w_2 -type III, nontrivial Kirby-Siebenmann invariant, and fundamental group G .

There can be no rational homology sphere with w_2 -type I by [HK88, Theorem 4.2]; in the notation of that theorem, $Q''(\pi_1, 0)$ gives rise to manifolds with nontrivial $H_2(-; \mathbb{Q})$. Now we describe the partial realisation of the invariants from Theorem 12.9.

- (1) By taking the connected sum of Σ_G^{II} with a closed, spin, simply connected manifold, we can realise any even, nonsingular, symmetric, bilinear form as the intersection form $\lambda_M: FH_2(M; \mathbb{Z}) \times FH_2(M; \mathbb{Z}) \rightarrow \mathbb{Z}$ of a closed, oriented 4-manifold M with fundamental group G and with w_2 type II. In this case $\text{ks}(M)$ is determined by the signature of λ_M .
- (2) Likewise, taking connected sum of $\Sigma_G^{III,0}$ or $\Sigma_G^{III,1}$ with a closed, spin, simply connected manifold, we can realise every even λ_M as the intersection form of a closed, oriented 4-manifold M with fundamental group G and with w_2 type III, with prescribed Kirby-Siebenmann invariant.
- (3) Finally, by taking connected sum of $\Sigma_G^{III,0}$ or $\Sigma_G^{III,1}$ with a closed, oriented, simply connected 4-manifold, we can realise any odd, nonsingular, symmetric, bilinear form as the intersection form $\lambda_M: FH_2(M; \mathbb{Z}) \times FH_2(M; \mathbb{Z}) \rightarrow \mathbb{Z}$ of a closed, oriented 4-manifold M with fundamental group G and with w_2 type I, with prescribed Kirby-Siebenmann invariant.

Question 12.10. *Must the intersection form in w_2 -type III be even? If not, how can we realise all intersection forms and Kirby-Siebenmann invariants in w_2 -type III?*

In his survey paper, Hambleton [Ham09, Theorem 5.2] also outlined a homeomorphism classification for closed, spin 4-manifolds with finite odd order fundamental group.

The following result on 4-manifold with finite fundamental group from [HK93b, Theorem B] also deserves to be mentioned.

Theorem 12.11. *Let M and N be closed, oriented, topological 4-manifolds with finite fundamental group. Suppose that $M \#^r(S^2 \times S^2)$ and $N \#^r(S^2 \times S^2)$ are homeomorphic for some $r \in \mathbb{N}$. Suppose that $X = X_0 \#(S^2 \times S^2)$. Then X is homeomorphic to Y .*

12.2.5. Nonorientable 4-manifolds. For nonorientable closed 4-manifolds, the homeomorphism classification results we are aware of are for fundamental group $\mathbb{Z}/2$ in [HKT94] and for fundamental group \mathbb{Z} in [Wan95]. For nonorientable closed 4-manifolds with fundamental group $\mathbb{Z}/2$, the paper [HKT94] gives a complete list of invariants for distinguishing such manifolds up to homeomorphism [HKT94, Theorem 2], and gives a list of the possible manifolds [HKT94, Theorem 3].

12.2.6. 4-manifolds with nonempty boundary. Simply-connected compact 4-manifolds with a fixed 3-manifold as boundary were classified by Boyer in [**Boy86**, **Boy93**], with an independent contribution by Stong [**Sto93**]. Homeomorphisms of such 4-manifolds were classified up to isotopy by Orson-Powell [**OP22**]. Since the statements are somewhat involved, we refer the reader to the original articles.

For compact 4-manifolds with fundamental group \mathbb{Z} , an analogous classification was given by Conway-Powell [**CP23**] and Conway-Piccirillo-Powell [**CPP22**], under the assumptions that $\pi_1(\partial M) \rightarrow \pi_1(M) \cong \mathbb{Z}$ is surjective, and that the homology $H_1(\partial M; \mathbb{Z}[\mathbb{Z}])$ of the corresponding \mathbb{Z} -cover is a $\mathbb{Z}[\mathbb{Z}]$ -torsion module.

Stable smoothing of homeomorphisms

Wall [Wal64] proved that simply connected, closed, smooth 4-manifolds with isometric intersection forms are stably diffeomorphic. It follows that every pair of simply connected, closed, homeomorphic smooth 4-manifolds are stably diffeomorphic. We shall discuss the analogous statement without the simply connected hypothesis.

Definition 13.1.

- (1) Let M and N be connected, smooth 4-manifolds. We say that M and N are *stably diffeomorphic* if there is an integer k such that the connected sums $M\#^k(S^2 \times S^2)$ and $N\#^k(S^2 \times S^2)$ are diffeomorphic.
- (2) Let M and N be connected 4-manifolds. We say that M and N are *stably homeomorphic* if there is an integer k such that the connected sums $M\#^k(S^2 \times S^2)$ and $N\#^k(S^2 \times S^2)$ are homeomorphic.

The next theorem is due to Gompf [Gom84].

Theorem 13.2. *Every homeomorphic pair of compact, connected, orientable, smooth 4-manifolds with diffeomorphic boundaries are stably diffeomorphic.*

Moreover, let $f: M \rightarrow N$ be a homeomorphism between two such 4-manifolds, that restricts to a diffeomorphism $f|: \partial M \rightarrow \partial N$. Then $f|$ extends to a stable diffeomorphism.

One might imagine a stronger statement, that given a homeomorphism $f: M \rightarrow N$ we can smoothen it stably up to isotopy. However such a statement is only known for simply connected 4-manifolds [FQ90, Chapter 8], and does not hold in general.

For non simply-connected manifolds, one must consider the bundle map of stable tangent microbundles induced by f , and lift it to a bundle map between the stable tangent bundles. Such a lift does not exist in general; there is a Casson-Sullivan obstruction in $H^3(M, \partial M; \mathbb{Z}/2)$ to its existence. Cappell-Shaneson [CS71], using unpublished work of R. Lee, showed that there is a homeomorphism of $(S^1 \times S^3)\#(S^2 \times S^2)$ for which this obstruction is nontrivial. Hence this homeomorphism cannot be stably smoothed. If a lift does exist, then for each lift there is a stabilisation by $\#^k(S^2 \times S^2)$, and then a pseudo-isotopy of the stabilised F to a diffeomorphism [FQ90, Chapter 8]. So even when the Casson-Sullivan invariant vanishes, one only has a stable smoothing up to pseudo-isotopy.

The proof of Theorem 13.2 that we shall give using Kreck's modified surgery [Kre99] was outlined in Teichner's thesis [Tei92, Theorem 5.1.1]. We think this proof is worth publicising with expanded details, because the method is arguably more conceptual than

Gompf's original, and because it allows us to expand on Gompf's statement in the nonorientable case.

Gompf also proved that for every pair of compact, connected, nonorientable, smooth 4-manifolds M and N that are homeomorphic, $M \# S^2 \tilde{\times} S^2$ and $N \# S^2 \tilde{\times} S^2$ are stably diffeomorphic. We shall slightly improve on this statement.

Theorem 13.3. *Let M and N be compact, connected, nonorientable, smooth 4-manifolds. Suppose that M and N are homeomorphic via a homeomorphism restricting to a diffeomorphism $\partial M \cong \partial N$. If $w_2(\tilde{M}) \neq 0 \neq w_2(\tilde{N})$, that is the universal covers of M and N are not spin, then M and N are stably diffeomorphic via a stable diffeomorphism extending the given diffeomorphism $\partial M \cong \partial N$.*

Gompf's statement [Gom84, p. 116] for the nonorientable case, given in the next corollary, follows easily from Theorem 13.3. However note that Theorem 13.3 shows that for many nonorientable 4-manifolds, the extra summand given by the twisted bundle $S^2 \tilde{\times} S^2$ is not necessary.

Corollary 13.4. *Let M and N be compact, connected, nonorientable, smooth 4-manifolds. Suppose that M and N are homeomorphic via a homeomorphism restricting to a diffeomorphism $\partial M \cong \partial N$. Then $M \# S^2 \tilde{\times} S^2$ and $N \# S^2 \tilde{\times} S^2$ are stably diffeomorphic.*

PROOF. Taking the connected sum of any 4-manifold with $S^2 \tilde{\times} S^2 \cong \mathbb{C}\mathbb{P}^2 \# \overline{\mathbb{C}\mathbb{P}^2}$ gives rise to a 4-manifold whose universal cover is not spin. The corollary therefore follows from Theorem 13.3. \square

The hypothesis in Theorem 13.3 that $w_2(\tilde{M}) \neq 0 \neq w_2(\tilde{N})$ cannot be dropped in general. Cappell and Shaneson found an example of a smooth 4-manifold R that is homotopy equivalent to the real projective space $\mathbb{R}\mathbb{P}^4$ but that is not stably diffeomorphic to $\mathbb{R}\mathbb{P}^4$ [CS71, CS76]. When these papers were published, it was not possible to prove that the fake $\mathbb{R}\mathbb{P}^4$ manifold R is homeomorphic to $\mathbb{R}\mathbb{P}^4$, but this was later established [Rub84, p. 221] as a consequence of the work of Freedman and Quinn [FQ90], and the fact that the Whitehead group of $\mathbb{Z}/2$ is trivial.

Later, Kreck [Kre84] showed a much more general statement in this direction. Let $K3 := \{[z_0 : z_1 : z_2 : z_3] \in \mathbb{C}\mathbb{P}^3 \mid z_0^4 + z_1^4 + z_2^4 + z_3^4 = 0\}$ denote the Kummer surface. As discussed in [GS99, Chapter 1.3], this is a closed, smooth, spin 4-manifold with signature 16, $b_2(K3) = 22$ and intersection form $3 \cdot H \oplus 2 \cdot E_8$. Here is Kreck's result from [Kre84]. These were the first known examples of exotic pairs of 4-manifolds.

Theorem 13.5. *Let π be a finitely presented group with a surjective homomorphism $w: \pi \rightarrow \mathbb{Z}/2$. Then there exists a closed, smooth, connected 4-manifold W with fundamental group π and orientation character w , with the property that $W \# K3$ and $W \#^{11}(S^2 \times S^2)$ are homeomorphic 4-manifolds that are not stably diffeomorphic.*

One part of this is easy to see: if W is nonorientable then there are homeomorphisms

$$\begin{aligned} W \# K3 &\cong W \# E_8 \# E_8 \#^3(S^2 \times S^2) \cong W \# E_8 \# \overline{E}_8 \#^3(S^2 \times S^2) \\ &\cong W \#^8(S^2 \times S^2) \#^3(S^2 \times S^2) \cong W \#^{11}(S^2 \times S^2). \end{aligned}$$

Here we used Theorem 12.2 that simply connected closed 4-manifolds with Kirby-Siebenmann invariant vanishing are determined by their intersection forms, and we used that the connected sum $M \# N$ of an oriented manifold M with a nonorientable manifold N is homeomorphic to $\overline{M} \# N$.

In the following three sections we will prove Theorems 13.2 and 13.3. To keep the notation manageable we will only provide a proof for closed manifolds, and describe the case of nonempty boundary in Section 13.4.

13.1. KRECK'S MODIFIED SURGERY

Below we will state a theorem due to Kreck that relates stable diffeomorphisms of 4-manifolds with bordism theory. This came as a corollary of Kreck's modified surgery theory [Kre99]. First we need some definitions from [Kre99].

Recall that a topological space A is m -connected if $\pi_k(A) = 0$ for $1 \leq k \leq m$ and is m -coconnected if $\pi_k(A) = 0$ for $k \geq m$. A map of spaces $f: A \rightarrow B$ is m -connected if the homotopy cofibre (i.e. the mapping cone) is m -connected; equivalently $f_*: \pi_k(A) \rightarrow \pi_k(B)$ is an isomorphism for $k < m$ and is surjective for $k = m$. A map of spaces $f: A \rightarrow B$ is m -coconnected if the homotopy fibre is m -coconnected; equivalently $f_*: \pi_k(A) \rightarrow \pi_k(B)$ is an isomorphism for $k > m$ and is injective for $k = m$.

Definition 13.6. A *normal 1-type* of a closed, connected, smooth 4-manifold M is a 2-coconnected fibration $\xi: B \rightarrow \text{BO}$ for which there is a 2-connected lift $\tilde{\nu}_M: M \rightarrow B$ of the stable normal vector bundle $\nu_M: M \rightarrow \text{BO}$ such that $\xi \circ \tilde{\nu}_M = \nu_M: M \rightarrow \text{BO}$. We call such a choice of lift $\tilde{\nu}_M: M \rightarrow B$ a *normal 1-smoothing*.

Remark 13.7. The data of a normal 1-type is $\xi: B \rightarrow \text{BO}$. The existence of $\tilde{\nu}_M$ is a condition on that data.

Definition 13.8. A *normal 1-type* of a closed, connected 4-manifold M is a 2-coconnected fibration $\xi^{\text{TOP}}: B^{\text{TOP}} \rightarrow \text{BTOP}$ for which there is a 2-connected lift $\tilde{\nu}_M: M \rightarrow B^{\text{TOP}}$ of the stable topological normal bundle $\nu_M: M \rightarrow \text{BTOP}$ (Definition 6.22) such that $\xi^{\text{TOP}} \circ \tilde{\nu}_M = \nu_M: M \rightarrow \text{BTOP}$. We call such a choice of lift $\tilde{\nu}_M: M \rightarrow B^{\text{TOP}}$ a *normal TOP 1-smoothing*.

Normal 1-types $\xi: B \rightarrow \text{BO}$ of a closed, connected smooth 4-manifold are fibre homotopy equivalent over BO , and using this we abuse notation and refer to *the* normal 1-type of a smooth 4-manifold, and similarly for the topological version. Here are some of the key examples in the oriented case. We will give the details of the nonorientable case in Section 13.3.

Write $\pi = \pi_1(M)$ and let $w_2 \in H^2(M; \mathbb{Z}/2)$ be the second Stiefel-Whitney class of M . There are three main cases for the normal 1-types of oriented, closed smooth 4-manifolds. For more details, see [KLPT17, Sections 2 and 3].

Lemma 13.9. *Let M be a closed, oriented, connected, smooth 4-manifold with universal covering \widetilde{M} . We write $\pi := \pi_1(M)$.*

- (1) *Suppose that we have $w_2(\widetilde{M}) \neq 0$. Then $\xi: B = B\pi \times \text{BSO} \rightarrow \text{BO}$ is the normal 1-type of M , with the map ξ given by projection to BSO followed by the canonical map $\text{BSO} \rightarrow \text{BO}$.*
- (2) *Suppose that $w_2(M) = 0$, i.e. M is spin. Then $\xi: B = B\pi \times \text{BSpin} \rightarrow \text{BO}$ is the normal 1-type of M , with the map ξ given by projection to BSpin followed by the canonical map $\text{BSpin} \rightarrow \text{BO}$.*
- (3) *Suppose that we have $w_2(M) \neq 0$ but $w_2(\widetilde{M}) = 0$. Then there is a model for $B\pi$ and a fibration $w_2: B\pi \rightarrow K(\mathbb{Z}/2, 2)$ that pulls back along $M \rightarrow B\pi$ to $w_2(M)$. The fibration $\xi: B \rightarrow \text{BO}$ is obtained from pulling back w_2 along the universal class $\overline{w}_2: \text{BSO} \rightarrow K(\mathbb{Z}/2, 2)$, to obtain the space B and a fibration $B \rightarrow \text{BSO}$, and then composing with $\text{BSO} \rightarrow \text{BO}$. Since we have a fibration sequence $\text{BSpin} \rightarrow \text{BSO} \xrightarrow{\overline{w}_2} K(\mathbb{Z}/2, 2)$, the pullback gives rise to a fibration sequence $\text{BSpin} \rightarrow B \rightarrow B\pi$. Then $\xi: B \rightarrow \text{BO}$ is the normal 1-type of M .*

Recall $\pi_1(\text{STOP}) \cong \pi_1(\text{TOP}) \cong \pi_1(\text{O}) \cong \mathbb{Z}/2$; see Chapter 7.

Lemma 13.10. *Let M be a closed, oriented, connected 4-manifold with universal covering \widetilde{M} . We write $\pi := \pi_1(M)$.*

- (1) *Suppose that we have $w_2(\widetilde{M}) \neq 0$. Then $\xi^{\text{TOP}}: B^{\text{TOP}} = B\pi \times \text{BSTOP} \rightarrow \text{BTOP}$ is the normal 1-type of M , with the map given by projection to BSTOP followed by the canonical map $\text{BSTOP} \rightarrow \text{BTOP}$.*
- (2) *Suppose that $w_2(M) = 0$, i.e. M is spin. Then $\xi^{\text{TOP}}: B^{\text{TOP}} = B\pi \times \text{BTOPSpin} \rightarrow \text{BTOP}$ is the normal 1-type of M , with the map given by projection to BTOPSpin followed by the canonical map $\text{BTOPSpin} \rightarrow \text{BTOP}$.*
- (3) *Suppose that we have $w_2(M) \neq 0$ but $w_2(\widetilde{M}) = 0$. Then there is a model for $B\pi$ and a fibration $w_2: B\pi \rightarrow K(\mathbb{Z}/2, 2)$ that pulls back along $M \rightarrow B\pi$ to $w_2(M)$. The fibration $\xi^{\text{TOP}}: B^{\text{TOP}} \rightarrow \text{BTOP}$ is obtained from pulling back w_2 along the universal class $\overline{w}_2: \text{BSTOP} \rightarrow K(\mathbb{Z}/2, 2)$, to obtain the space B^{TOP} and a fibration $B^{\text{TOP}} \rightarrow \text{BSTOP}$, and then composing with $\text{BSTOP} \rightarrow \text{BTOP}$. Since we have a fibration sequence $\text{BTOPSpin} \rightarrow \text{BSTOP} \xrightarrow{\overline{w}_2} K(\mathbb{Z}/2, 2)$, the pullback gives rise to a fibration sequence $\text{BTOPSpin} \rightarrow B \rightarrow B\pi$. Then $\xi^{\text{TOP}}: B^{\text{TOP}} \rightarrow \text{BTOP}$ is the normal 1-type of M .*

Here is the relevant theorem of Kreck [Kre99, Theorem C], which relates bordism over the normal 1-type to stable diffeomorphism. We write $\Omega_4(B, \xi)$ for the group under disjoint union of closed 4-manifolds M together with a lift $\overline{\nu}: M \rightarrow B$ along ξ of the stable normal vector bundle, considered up to bordism over (B, ξ) .

Theorem 13.11. *Two closed, connected, smooth 4-manifolds M and N with $\chi(M) = \chi(N)$ and normal 1-types both fibre homotopy equivalent to a fixed fibration $\xi: B \rightarrow \text{BO}$ are stably*

diffeomorphic if and only if

$$[(M, \tilde{\nu}_M)] = [(N, \tilde{\nu}_N)] \in \Omega_4(B, \xi)$$

for some choices of normal 1-smoothings $\tilde{\nu}_M$ and $\tilde{\nu}_N$.

SKETCH OF THE PROOF. One direction is quite easy: one has to check that M and $M\#(S^2 \times S^2)$ are bordant over the normal 1-type of M .

For the other direction, start with a 5-dimensional bordism W over (B, ξ) and perform surgery below the middle dimension [Kre99, Section 3] to arrange that the map $W \rightarrow B$ is 1-connected. We outline the procedure next. First perform surgery on pairs of discs, extending the map to B over the new copies of $D^1 \times S^4$ to make the map $\pi_1(W) \rightarrow \pi_1(B)$ surjective. Then represent normal generators of the kernel of $\pi_1(W) \rightarrow \pi_1(B)$ by framed circles using Chapters 5 and 10. Since $\pi_1(W)$ and $\pi_1(B)$ are finitely presented, this can be done with finitely many circles. To prove that the normal vector bundles admit framings one must use the bundle data ξ ; for this we refer to [Kre99, Section 3]. Perform surgery on the framed circles to make the map to B 1-connected. This completes the surgery below the middle dimension step.

Now represent the elements of $\ker(\pi_2(W) \rightarrow \pi_2(B))$ by framed embedded spheres, and remove thickenings of these spheres. Also, for each sphere, remove a tube $D^1 \times D^4$ connecting that sphere to either M or N . Choose whether to tube to M or N so as to preserve the Euler characteristic equality. This operation of removing copies of $S^2 \times D^3$, tubed to the boundary, has the effect of adding copies of $S^2 \times S^2$ to M and N giving rise to M' and N' respectively. The operation also converts W to an s -cobordism W' . That $(W'; M', N')$ is an s -cobordism means by definition that the inclusion maps $M' \rightarrow W'$ and $N' \rightarrow W'$ are simple homotopy equivalences. The stable s -cobordism theorem [Qui83] states that every 5-dimensional s -cobordism becomes diffeomorphic to a product after adding copies of $(S^2 \times S^2) \times I$ along a smoothly embedded interval $I \subseteq W'$ with one endpoint on each of M and N . This completes the sketch proof of Theorem 13.11. \square

The proof of the topological version is similar.

Theorem 13.12. *Two closed, topological 4-manifolds M and N with $\chi(M) = \chi(N)$ and normal 1-types both fibre homotopy equivalent to a fixed fibration $\xi^{\text{TOP}}: B^{\text{TOP}} \rightarrow \text{BTOP}$ are stably homeomorphic if and only if*

$$[(M, \tilde{\nu}_M)] = [(N, \tilde{\nu}_N)] \in \Omega_4^{\text{TOP}}(B^{\text{TOP}}, \xi^{\text{TOP}})$$

for some choices of normal 1-smoothings $\tilde{\nu}_M$ and $\tilde{\nu}_N$.

From now on, to ease notation, we will sometimes abbreviate $\Omega_4^{\text{TOP}}(B^{\text{TOP}}, \xi^{\text{TOP}})$ to $\Omega_4^{\text{TOP}}(B, \xi)$.

PROOF. One direction is again quite easy: we need that homeomorphic manifolds are bordant over B , and that M and $M\#(S^2 \times S^2)$ are bordant in $\Omega_4^{\text{TOP}}(B, \xi)$. For the other direction, apply the same argument as above to improve a cobordism W to an s -cobordism. The stable s -cobordism Theorem applies to topological s -cobordisms as well as to smooth

s -cobordisms. This is not written in [Qui83], but the same proof applies, with the following additions (see the Exercise on [FQ90, p. 107]). First, 5-dimensional cobordisms admit a topological handle structure [FQ90, Theorem 9.1]. The proof of [Qui83] consists of simplifying a handle decomposition, and tubing surfaces in 4-manifolds around and into parallel copies of one another to remove intersections. This is possible in the topological category by using transversality (Theorem 10.3) to arrange that intersections between surfaces are isolated points, and the existence of normal vector bundles (Theorem 5.14) to take parallel copies using sections. \square

13.2. STABLE DIFFEOMORPHISM OF HOMEOMORPHIC ORIENTABLE 4-MANIFOLDS

Now we will explain the proof of Theorem 13.2. For the convenience of the reader, we recall the statement.

Theorem 13.2. *Every homeomorphic pair of closed, connected, orientable, smooth 4-manifolds are stably diffeomorphic.*

The proof will rest on the following proposition.

Proposition 13.13. *Let (B, ξ) be one of the oriented smooth normal 1-types from Lemma 13.9, and let $(B^{\text{TOP}}, \xi^{\text{TOP}})$ be the corresponding topological normal 1-type from Lemma 13.10 obtained by replacing BSO with BSTOP or BSpin with BTOPSpin as appropriate. The forgetful map*

$$F: \Omega_4(B, \xi) \rightarrow \Omega_4^{\text{TOP}}(B^{\text{TOP}}, \xi^{\text{TOP}}) = \Omega_4^{\text{TOP}}(B, \xi)$$

is injective.

The combination of this proposition with Theorems 13.11 and 13.12 implies the following corollary, which is the closed version of Theorem 13.2, with a slightly more precise statement concerning orientations.

Corollary 13.14. *Every pair of smooth, closed, connected, oriented 4-manifolds that are homeomorphic via an orientation preserving homeomorphism are stably diffeomorphic via an orientation preserving diffeomorphism.*

PROOF. We prove the corollary assuming Proposition 13.13. Homeomorphic 4-manifolds are in particular stably homeomorphic and have the same normal 1-types. Therefore two homeomorphic smooth 4-manifolds as in the statement of the corollary are bordant over the normal 1-type, so give rise to equal elements in $\Omega_4^{\text{TOP}}(B, \xi)$. By Proposition 13.13, they give rise to equal elements of $\Omega_4(B, \xi)$. Then by Theorem 13.11, the two 4-manifolds are stably diffeomorphic, as asserted. \square

PROOF OF PROPOSITION 13.13. Let S be SO in case (1) of the smooth list of 1-types given in Lemma 13.10, and let S denote Spin in cases (2) and (3).

Let ST be STOP in case (1) of the topological list of 1-types above, and let ST denote TOPSpin in cases (2) and (3).

The James spectral sequence [Tei92, Theorem 3.1.1], [KLPT17, Section 3] is of the form:

$$E_{p,q}^2 = H_p(\mathbb{B}\pi; \Omega_q^S) \Rightarrow \Omega_{p+q}(B, \xi).$$

We have that $\Omega_4^S \cong \mathbb{Z}$, detected by the signature. Indeed, the signature is a \mathbb{Z} -valued invariant that agrees for stably diffeomorphic 4-manifolds. The signature of a 4-manifold with a normal 1-smoothing into B gives rise to an element of $\Omega_4(B, \xi)$. The E^2 term $H_0(\mathbb{B}\pi; \Omega_4^S) \cong \mathbb{Z}$ is computed using $\Omega_4^S \cong \mathbb{Z}$.

Claim. This term $H_0(\mathbb{B}\pi; \Omega_4^S)$ survives to the E^∞ page. That is, all differentials with this as codomain are trivial.

Let us prove the claim. Since Ω_q^S is torsion for $q = 1, 2, 3$, no terms from those q -lines can map to $H_0(\mathbb{B}\pi; \Omega_4^S)$ under a differential.

Aside from $H_0(\mathbb{B}\pi; \Omega_4^S)$, there is one other potentially infinite term on the 4-line of the E^∞ page, namely the subgroup of $H_4(\mathbb{B}\pi; \Omega_0^S)$ arising as the kernel of relevant differentials. Since there are no differentials with the $(4, 0)$ term as codomain, this subgroup is a quotient of $\Omega_4(B, \xi)$. The image of $[M, c] \in \Omega_4(B, \xi)$ is the image $c_*([M])$ of the fundamental class under the classifying map $c_*: H_4(M; \mathbb{Z}) \rightarrow H_4(\mathbb{B}\pi; \mathbb{Z}) \cong H_4(\mathbb{B}\pi; \Omega_0^S)$.

There could be a nontrivial differential $d_{5,0}^5: H_5(\mathbb{B}\pi; \Omega_0^S) \rightarrow H_0(\mathbb{B}\pi; \Omega_4^S)$. However if there were a nonzero differential, then only finitely many signatures would occur for 4-manifolds with normal 1-type B and fixed invariant in $H_4(\mathbb{B}\pi; \Omega_0^S)$. But we can add copies of the $K3$ -surface, mapping to a point in B , to a given fixed element of $\Omega_4(B, \xi)$, keeping the normal 1-type and $c_*([M])$ the same, but changing the signature by $+16$ for each copy of $K3$. This contradiction implies that $d_{5,0}^5$ is the zero map. This completes the proof of the claim that the term $H_0(\mathbb{B}\pi; \Omega_4^S)$ survives to the E^∞ page.

Since $H_0(\mathbb{B}\pi; \Omega_4^S)$ survives to the E^∞ term, we have a short exact sequence:

$$0 \rightarrow \Omega_4^S \rightarrow \Omega_4(B, \xi) \rightarrow \tilde{\Omega}_4(B, \xi) \rightarrow 0,$$

where $\tilde{\Omega}_4(B, \xi)$ denotes the quotient. That is, there is a filtration with iterated graded quotients given by the E^∞ page:

$$0 \subseteq E_{4,0}^\infty = \Omega_4^S \subseteq \cdots \subseteq \Omega_4(B, \xi),$$

and it is the quotient by the $E_{4,0}^\infty$ subgroup that we denote $\tilde{\Omega}_4(B, \xi)$.

Similarly, for the topological case, we have

$$0 \rightarrow \Omega_4^{ST} \rightarrow \Omega_4^{\text{TOP}}(B, \xi) \rightarrow \tilde{\Omega}_4^{\text{TOP}}(B, \xi) \rightarrow 0.$$

The only difference in the proof from the smooth case is that we also have to argue that the Kirby-Siebenmann invariant $\mathbb{Z}/2 \subseteq \Omega_4^{ST}$ survives to the E^∞ page. But the Kirby-Siebenmann invariant is additive, and realised on simply connected manifolds, by the E_8 manifold. Thus there exist bordism classes (i.e. stable homeomorphism classes) realising both trivial and nontrivial Kirby-Siebenmann invariants within a normal 1-type, and so this $\mathbb{Z}/2$ cannot be killed by a differential.

Since the structure forgetting map $\Omega_q^S \rightarrow \Omega_q^{ST}$ is an isomorphism for $0 \leq q \leq 3$, we have an isomorphism $\tilde{\Omega}_4(B, \xi) \cong \tilde{\Omega}_4^{\text{TOP}}(B, \xi)$. This uses that the differentials agree, by naturality of the James spectral sequence with respect to homology theories. Indeed, note that the differentials depend only on the classifying space $B\pi$, and on the complex line bundle $E \rightarrow B\pi$ in case (3). Both are category independent.

Then there is a map of short exact sequences:

$$\begin{array}{ccccccc} 0 & \longrightarrow & \Omega_4^S & \longrightarrow & \Omega_4(B, \xi) & \longrightarrow & \tilde{\Omega}_4(B, \xi) \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \cong \\ 0 & \longrightarrow & \Omega_4^{ST} & \longrightarrow & \Omega_4^{\text{TOP}}(B, \xi) & \longrightarrow & \tilde{\Omega}_4^{\text{TOP}}(B, \xi) \longrightarrow 0. \end{array}$$

The left vertical map is injective, either inclusion into the first summand $\mathbb{Z} \rightarrow \mathbb{Z} \oplus \mathbb{Z}/2$ for non-spin or $16\mathbb{Z} \rightarrow 8\mathbb{Z}$ in the spin case, when $B = B\pi \times B\text{Spin}$ and $B^{\text{TOP}} = B\pi \times B\text{TOPSpin}$. Since the left and right vertical maps are injective, it follows from a diagram chase that the central vertical map is also injective, as required. \square

13.3. NONORIENTABLE 4-MANIFOLDS AND STABLE DIFFEOMORPHISM

For the convenience of the reader, we recall the statement of Theorem 13.3.

Theorem 13.3. *Let M and N be closed, connected, nonorientable, smooth 4-manifolds. Suppose that M and N are homeomorphic. If $w_2(\tilde{M}) \neq 0 \neq w_2(\tilde{N})$, that is the universal covers of M and N are not spin, then M and N are stably diffeomorphic.*

Here is the normal 1-type for nonorientable manifolds with a certain w_2 -type [Tei92, Chapter 2], first in the smooth and then in the topological case.

Lemma 13.15. *Let M be a nonorientable closed, connected smooth 4-manifold with $w_2(\tilde{M}) \neq 0$. We set $\pi := \pi_1(M)$. Then the normal 1-type of M is $\xi: B = B\pi \times B\text{SO} \rightarrow B\text{O}$ with the map $\xi = w_1 \oplus Bi$ given by the Whitney sum of a bundle on $B\pi$ determined by $w_1: \pi \rightarrow \mathbb{Z}/2$ and the canonical map $Bi: B\text{SO} \rightarrow B\text{O}$ induced by the inclusion $i: \text{SO} \rightarrow \text{O}$.*

Lemma 13.16. *Let M be a nonorientable closed, connected 4-manifold with $w_2(\tilde{M}) \neq 0$. We set $\pi := \pi_1(M)$. Then the normal 1-type of M is $\xi: B = B\pi \times B\text{STOP} \rightarrow B\text{TOP}$ with the map $\xi = w_1 \oplus Bi$ given by the Whitney sum of a bundle on $B\pi$ determined by the orientation character $w_1: \pi \rightarrow \mathbb{Z}/2$ and the canonical map $Bi: B\text{STOP} \rightarrow B\text{TOP}$ induced by the inclusion $i: \text{STOP} \rightarrow \text{TOP}$.*

These normal 1-types give rise to a James spectral sequence (see [Tei93, Chapter II] for details) governing the bordism groups of (B, ξ)

$$E_{p,q}^2 = H_p(B\pi; \Omega_q^{w_1}) \Rightarrow \Omega_{p+q}(B, \xi).$$

Note that the coefficients are twisted using $\mathbb{Z}^{w_1} \otimes \Omega_q$, where by definition, $g \in \pi$ acts on \mathbb{Z}^{w_1} by multiplication by $(-1)^{w_1(g)}$. The corresponding topological James spectral sequence is:

$$E_{p,q}^2 = H_p(B\pi; (\Omega_q^{\text{STOP}})^{w_1}) \Rightarrow \Omega_{p+q}^{\text{TOP}}(B, \xi).$$

As in the previous section, here we abbreviate $\Omega_4^{\text{TOP}}(B^{\text{TOP}}, \xi^{\text{TOP}})$ to $\Omega_4^{\text{TOP}}(B, \xi)$.

By Kreck's Theorem 13.11) and the argument in the proof of Theorem 13.2, in order to prove Theorem 13.3 it suffices to prove the next injectivity statement, which is an analogue of Proposition 13.13.

Proposition 13.17. *Let (B, ξ) be one of the normal 1-types in Lemma 13.15 and let $(B^{\text{TOP}}, \xi^{\text{TOP}})$ be the corresponding topological normal 1-type over BTOP. The forgetful map*

$$F: \Omega_4(B, \xi) \rightarrow \Omega_4^{\text{TOP}}(B^{\text{TOP}}, \xi^{\text{TOP}}) = \Omega^{\text{TOP}}(B, \xi)$$

is injective.

PROOF OF THEOREM 13.3 ASSUMING PROPOSITION 13.17. Homeomorphic 4-manifolds have the same normal 1-types and are trivially TOP bordant over this normal 1-type, injectivity of F implies that homeomorphic nonorientable, closed, connected, smooth 4-manifolds are smoothly bordant over their normal 1-type, and therefore by Theorem 13.11 are stably diffeomorphic. \square

PROOF OF PROPOSITION 13.17. The structure of the proof is very similar to that of the proof of Proposition 13.13. This proof is therefore somewhat terse. In the smooth James spectral sequence computing $\Omega_4(B, \xi)$, we consider the term on the E^2 page $H_0(\mathbb{B}\pi; \Omega_4^{w_1}) \cong \mathbb{Z}/2$. This is detected by the Euler characteristic of the manifold modulo two.

Since we can always perform connected sum with a copy of $\mathbb{C}\mathbb{P}^2$ (note that for nonorientable manifolds connected sum with $\mathbb{C}\mathbb{P}^2$ and $\overline{\mathbb{C}\mathbb{P}^2}$ is the same), both mod 2 Euler characteristics are realised by bordism classes over (B, ξ) . Also note that adding $\mathbb{C}\mathbb{P}^2$ does not change the normal 1-type when $w_2(\widetilde{M}) \neq 0$. Therefore $H_0(\mathbb{B}\pi; \Omega_4^{w_1}) \cong \mathbb{Z}/2$ survives to the E^∞ page.

In the topological case, the corresponding term in the James spectral sequence computing $\Omega_4^{\text{TOP}}(B, \xi)$ is

$$H_0(\mathbb{B}\pi; (\Omega_4^{\text{STOP}})^{w_1}) \cong \mathbb{Z}/2 \oplus \mathbb{Z}/2.$$

We can add copies of $\mathbb{C}\mathbb{P}^2$ and $*\mathbb{C}\mathbb{P}^2$ to a given element of $\Omega_4^{\text{TOP}}(B, \xi)$ to show that this term survives to the E^∞ page. The structure forgetting map $\mathbb{Z}/2 \cong H_0(\mathbb{B}\pi; \Omega_4^{w_1}) \rightarrow H_0(\mathbb{B}\pi; (\Omega_4^{\text{STOP}})^{w_1}) \cong \mathbb{Z}/2 \oplus \mathbb{Z}/2$ is injective.

Therefore the filtrations of $\Omega_4(B, \xi)$ and $\Omega_4^{\text{TOP}}(B, \xi)$ arising from the spectral sequence give rise to short exact sequences, that form the rows of the following commutative diagram:

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mathbb{Z}/2 & \longrightarrow & \Omega_4(B, \xi) & \longrightarrow & \widetilde{\Omega}_4(B, \xi) \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \cong \\ 0 & \longrightarrow & \mathbb{Z}/2 \oplus \mathbb{Z}/2 & \longrightarrow & \Omega_4^{\text{TOP}}(B, \xi) & \longrightarrow & \widetilde{\Omega}_4^{\text{TOP}}(B, \xi) \longrightarrow 0. \end{array}$$

We noted above that the left vertical map is injective. Since $\Omega_q \rightarrow \Omega_q^{\text{STOP}}$ is an isomorphism for $0 \leq q \leq 3$, the right vertical map is an isomorphism and is therefore injective. It follows from a diagram chase that the central vertical map is also injective, as required. \square

For the other normal 1-types of nonorientable 4-manifolds, the $E_{0,4}^2$ terms are given by the homology with twisted coefficients $H_0(\mathbb{B}\pi; (\Omega_4^{\text{Spin}})^{w_1})$ and $H_0(\mathbb{B}\pi; (\Omega_4^{\text{TOPSpin}})^{w_1})$. The forgetful map $H_0(\mathbb{B}\pi; (\Omega_4^{\text{Spin}})^{w_1}) \rightarrow H_0(\mathbb{B}\pi; (\Omega_4^{\text{TOPSpin}})^{w_1})$ is not injective, because $16\mathbb{Z} \cong \Omega_4^{\text{Spin}} \rightarrow \Omega_4^{\text{TOPSpin}} \cong 8\mathbb{Z}$ is not surjective. Not only does the proof break down in these cases, but the examples in Theorem 13.5 show that the hypothesis on the w_2 -type cannot be removed.

13.4. HOMEOMORPHIC 4-MANIFOLDS WITH BOUNDARY

In this section we briefly explain how to extend the proofs to manifolds with boundary. Here is the relevant version of Kreck's theorem [Kre99].

Theorem 13.18. *Let M and N be compact (smooth) 4-manifolds, let $f: \partial M \rightarrow \partial N$ be a diffeomorphism, and suppose that the normal 1-types of M and N are both fibre homotopy equivalent to the same fibration $\xi^{\text{TOP}}: B^{\text{TOP}} \rightarrow \text{BTOP}$ ($\xi: B \rightarrow \text{BO}$).*

Then M and N are stably homeomorphic (diffeomorphic) via a stable homeomorphism (diffeomorphism) extending f if and only if M and N admit normal 1-smoothings $\bar{\nu}_M$ and $\bar{\nu}_N$ into B^{TOP} (B) such that $\bar{\nu}_M = \bar{\nu}_N \circ f: \partial M \rightarrow B^{\text{TOP}}$ (B), for which the union $(M \cup_f N, \bar{\nu}_M \cup -\bar{\nu}_N)$ represents the trivial element of $\Omega_4(B^{\text{TOP}}, \xi^{\text{TOP}})$ ($\Omega_4(B, \xi)$).

In Propositions 13.13 and 13.17 we proved injectivity of the forgetful maps in bordism $\Omega_4(B, \xi) \rightarrow \Omega_4^{\text{TOP}}(B^{\text{TOP}}, \xi^{\text{TOP}})$ relevant to Theorems 13.2 and 13.3.

PROOF OF THEOREMS 13.2 AND 13.3. Let M and N be smooth, compact 4-manifolds with nonempty boundary. Let $(B^{\text{TOP}}, \xi^{\text{TOP}})$ and (B, ξ) be the relevant normal 1-types. Let $f: \partial M \rightarrow \partial N$ be a diffeomorphism that extends to a homeomorphism from M to N . Then by Theorem 13.18 in the topological category, there exist normal 1-smoothings $\bar{\nu}_M^{\text{TOP}}: M \rightarrow B^{\text{TOP}}$ and $\bar{\nu}_N^{\text{TOP}}: N \rightarrow B^{\text{TOP}}$ such that $(M \cup_f N, \bar{\nu}_M^{\text{TOP}} \cup -\bar{\nu}_N^{\text{TOP}})$ represents the trivial element of $\Omega_4(B^{\text{TOP}}, \xi^{\text{TOP}})$. Since M and N are smooth, their stable normal vector bundle classifying maps lift along $\text{BO} \rightarrow \text{BTOP}$, and we obtain normal 1-smoothings $\bar{\nu}_M: M \rightarrow B$ and $\bar{\nu}_N: N \rightarrow B$ such that $(M \cup_f N, \bar{\nu}_M \cup -\bar{\nu}_N) \in \Omega_4(B, \xi)$ maps under the forgetful map to $(M \cup_f N, \bar{\nu}_M^{\text{TOP}} \cup -\bar{\nu}_N^{\text{TOP}}) = 0 \in \Omega_4(B^{\text{TOP}}, \xi^{\text{TOP}})$. By injectivity of the forgetful maps $\Omega_4(B, \xi) \rightarrow \Omega_4^{\text{TOP}}(B^{\text{TOP}}, \xi^{\text{TOP}})$, as proven in Propositions 13.13 and 13.17, it follows that $(M \cup_f N, \bar{\nu}_M \cup -\bar{\nu}_N) = 0 \in \Omega_4(B, \xi)$. By Theorem 13.18, we learn that indeed M and N are stably diffeomorphic via a stable diffeomorphism extending the given diffeomorphism $f: \partial M \rightarrow \partial N$. \square

CHAPTER 14

Reidemeister torsion in the topological category

14.1. THE SIMPLE HOMOTOPY TYPE OF A MANIFOLD

In the following we need the notion of a *simple* homotopy equivalence. We will not give a definition, instead we refer to [Tur01, p. 40] for details. Roughly, a simple homotopy equivalence between CW complexes is a sequence of elementary expansions and collapses of pairs of cells whose dimension differs by one.

As we discussed in Chapter 3, it is not clear whether topological 4-manifolds have CW-structure. Fortunately the following definition allows us to define a simple homotopy type even for topological spaces which are not homeomorphic to a CW complex.

Definition 14.1. Let (W, V) be a pair of topological spaces. Consider tuples (W, V, f, X, Y) , where (X, Y) is a finite CW complex pair with $Y \subseteq X$, and $f: W \rightarrow X$ and $f|_V: V \rightarrow Y$ homotopy equivalences. Two such tuples (W, V, f, X, Y) and (W, V, f', X', Y') , with (X', Y') another finite CW pair and $f': (W, V) \rightarrow (X', Y')$, are *equivalent* if there exists a simple homotopy equivalence of pairs $s: (X, Y) \rightarrow (X', Y')$ such that $s \circ f$ is homotopic to f' and $s|_Y \circ f|_V: V \rightarrow Y'$ is homotopic to $f'|_V$. Such an equivalence class of (W, V, f, X, Y) is called a *simple homotopy type* of (W, V) . In particular, a simple homotopy type of (W, \emptyset) is called a *simple homotopy type* of W .

Now consider a compact, connected n -manifold M . If M admits a smooth structure, then by Theorem 3.13 we know that M admits in particular a CW structure, and we equip M with the simple homotopy type given by $(M, \emptyset, \text{Id}, M, \emptyset)$. By Chapman's Theorem [Cha74, p. 488] below, this simple homotopy type is independent of the choice of CW structure on M .

Theorem 14.2. (Chapman's Theorem) *Let W be a compact topological space. Any two CW structures on W are simple homotopy equivalent.*

As we pointed out in Chapter 3, it is unknown whether every compact manifold admits a CW structure. In the remainder of this section, we will nonetheless introduce the simple homotopy type of a compact manifold M following [KS77, Essay III, Section 4, p. 117]. The first step is to construct a disc bundle $D(M) \rightarrow M$ together with a PL structure on the total space $D(M)$. We will work with a compact m -dimensional manifold M with boundary ∂M , and seek to construct the simple homotopy type of $(M, \partial M)$.

Construction 14.3. We deal with the case $\partial M = \emptyset$ first, and then later address the additional complications arising from having nonempty boundary.

As a first step to constructing the disc bundle $D(M) \subseteq \mathbb{R}^n$, we need an embedding of M into \mathbb{R}^{n-1} for some large integer $n - 1 > 2m + 5$. For a closed m -manifold M such an embedding is readily available [**Hat02**, Corollary A.9]. It follows from Theorem 6.20 that for $n - 1 > 2m + 5$ all such embeddings of M are isotopic, and that they admit a normal microbundle $\nu_{\mathbb{R}^{n-1}}(M)$ that is unique up to isotopy. By Theorem 6.17 this normal microbundle $\nu_{\mathbb{R}^{n-1}}(M)$ can be upgraded to a topological \mathbb{R}^{n-1-m} -bundle. By taking the product with \mathbb{R} , construct an embedding $M \subseteq \mathbb{R}^n$ whose normal microbundle is $\nu(M) = \nu_{\mathbb{R}^{n-1}}(M) \times \mathbb{R}$. Since we stabilised once, the normal microbundle $\nu(M)$ contains a normal disc bundle $B(M)$ [**KS77**, Essay III, Proposition 4.4, p. 120].

The next big step will be to upgrade $B(M) \subseteq \mathbb{R}^n$ from a submanifold to a PL submanifold. Since the interior is codimension 0, the interior of $B(M)$ is automatically also a PL submanifold. However, we have to arrange $\partial B(M)$ to be a PL submanifold of \mathbb{R}^n itself. In the next paragraphs, we modify the PL structure on \mathbb{R}^n such that $\partial B(M)$ becomes a PL submanifold and then isotope this new PL structure on \mathbb{R}^n back to the standard PL structure.

Using the Collar Neighbourhood Theorem 2.16, pick a collar $W_\partial = \partial B(M) \times (-1, 1)$ and $D_\partial = \partial B(M) \times [-\frac{1}{2}, \frac{1}{2}]$. The Local Product Structure Theorem 4.18 [**KS77**, Essay I, Theorem 5.2, p. 36], applied with $W = W_\partial$ and $D = D_\partial$, gives a PL structure σ_∂ on \mathbb{R}^n such that $\partial B(M)$ is a PL submanifold and σ_∂ is concordant to the standard PL structure σ_{std} .




FIGURE 14.1. Illustration of $B(M)$.

Now we will isotope the pair $\partial B(M) \subseteq B(M)$ so that they become PL submanifolds of $(\mathbb{R}^n, \sigma_{\text{std}})$. The PL structure σ_∂ is concordant to σ_{std} . Since concordance implies isotopy [**KS77**, Essay I, Theorem 4.1, p. 25] in dimension $m \geq 6$, there is an isotopy $\phi_t \in \text{Homeo}(\mathbb{R}^n)$ such that $\phi_0 = \text{Id}$, and $\phi_1^* \sigma_{\text{std}} = \sigma_\partial$. Consequently, $D(M) := \phi_1(B(M))$ and $D(\partial M) := \phi_1(B(\partial M))$ are PL submanifolds of $(\mathbb{R}^n, \sigma_{\text{std}})$, which defines a simple type of M , the tuple $(M, z, D(M))$, where $z: M \rightarrow D(M)$ is the zero section.

Having finished the case $\partial M \neq \emptyset$, next we discuss the procedure for a manifold M with nonempty boundary.

Take the union of M with an external open collar $\partial M \times [0, 1)$ of its boundary. Write $M' := M \cup_{\partial M} \partial M \times [0, 1)$. Embed M' into \mathbb{R}^n as in the closed case [**Hat02**, Corollary A.9]. Note that M' has empty boundary and so it is properly embedded. As in the closed case, obtain a disc bundle $B(M')$ and let $B(M)$ be the restriction of this disc bundle to M .

Now we have to take much more care. Note that $\partial B(M)$ decomposes as $\partial B(M) = B(\partial M) \cup_X B_\partial(M)$. Here $B_\partial(M)$ denotes the fibrewise boundary and $X = \partial(B(\partial M))$

denotes the intersection of $B(\partial M)$ and $B_\partial(M)$. As above, we will find a PL structure σ_∂ of \mathbb{R}^n such that each subset $B(\partial M)$, X and $B_\partial(M)$ is PL-submanifold of \mathbb{R}^n .

Our first goal is to modify the PL structure on \mathbb{R}^n so that the corners X become a PL submanifold of \mathbb{R}^n . Denote the standard PL structure on \mathbb{R}^n by σ_{std} . Pick a bicollar $\partial B(M) \times [-1, 1] \subseteq \mathbb{R}^n$ of the boundary of the codimension 0 submanifold $B(M)$. Again by the Collar Neighbourhood Theorem 2.16, we can pick a bicollar $X \times [-1, 1] \subseteq \partial B(M)$. We consider the open set $W_X := X \times (-1, 1)^2 \subseteq \partial B(M) \times (-1, 1) \subseteq \mathbb{R}^n$, and $D_X = X \times [-\frac{1}{2}, \frac{1}{2}]^2$. The Local Product Structure Theorem 4.18 [KS77, Essay I, Theorem 5.2, p. 36], applied with $W = W_X$ and $D = D_X$, gives a PL structure on X and a PL structure σ_X on \mathbb{R}^n , which is concordant to σ_{std} rel. $\mathbb{R}^n \setminus (X \times (-\frac{2}{3}, \frac{2}{3})^2)$. This PL structure σ_X has the property that it agrees with the product PL structure on $X \times (-1, 1)^2$ in a neighbourhood of D_X . Thus X is a PL submanifold of (\mathbb{R}^n, σ_X) .




FIGURE 14.2. Illustration of construction of $D(M)$ if $\partial M \neq \emptyset$.

Now we arrange the next stratum $\partial B(M) \supseteq X$ to be a PL submanifold of \mathbb{R}^n . Near $D_X = X \times [-\frac{1}{2}, \frac{1}{2}]$, the PL structure σ_X is the product PL structure, and therefore $\partial B(M) \cap \text{Int } D_X = X \times (-\frac{1}{2}, \frac{1}{2}) \times \{0\}$ is already a PL submanifold of (\mathbb{R}^n, σ_X) . Furthermore, σ_X is a product along $(-\frac{1}{2}, \frac{1}{2})$ near $X \times [-\frac{1}{3}, \frac{1}{3}]$. Pick $W_\partial = \partial B(M) \times (-1, 1)$, $C_\partial = X \times [-\frac{1}{3}, \frac{1}{3}] \times [-\frac{1}{3}, \frac{1}{3}]$ and $D_\partial = \partial B(M) \times [-\frac{1}{2}, \frac{1}{2}]$. As above, the Local Product Structure Theorem 4.18 [KS77, Essay I, Theorem 5.2, p. 36], applied with $W = W_\partial$, $C = C_\partial$, and $D = D_\partial$, gives a PL structure σ_∂ on \mathbb{R}^n such that $\partial B(M)$ is a PL submanifold and σ_∂ is concordant to σ_X rel. $(\mathbb{R}^n \setminus \partial B(M) \times (-\frac{2}{3}, \frac{2}{3})) \cup C_\partial$. Since $X \subseteq C_\partial$ the submanifold X is still a PL submanifold of $(\mathbb{R}^n, \sigma_\partial)$.

As in the closed case, use a concordance from σ_∂ to σ_{std} to obtain an isotopy $\phi_t \in \text{Homeo}(\mathbb{R}^n)$ such that $\phi_0 = \text{Id}$, and $\phi_1^* \sigma_{\text{std}} = \sigma_\partial$. Define $D(M) := \phi_1(B(M))$ and $D(\partial M) := \phi_1(B(\partial M))$, which are both PL submanifolds of $(\mathbb{R}^n, \sigma_{\text{std}})$. We obtain a simple homotopy type $(M, \partial M, z, D(M), D(\partial M))$, where again z is the zero section of the disc bundle. This finishes the case where M has nonempty boundary.

In both cases, ∂M empty and nonempty, our construction involved many choices. Let $D'(\partial M) \subseteq D'(M)$ be obtained by other choices. Following the discussion [KS77, p. 123],

we can suitably stabilise the bundles and find a commutative diagram of PL maps:

$$\begin{array}{ccc} D(M) \times D^s & \xrightarrow{\cong} & D'(M) \times D^r \\ \uparrow & & \uparrow \\ D(\partial M) \times D^s & \xrightarrow{\cong} & D'(\partial M) \times D^r, \end{array}$$

where D^k denotes the disc with its standard PL structure and the horizontal maps are PL isomorphisms that preserve the zero sections up to homotopy.

Definition 14.4. The *simple homotopy type* of a compact connected n -manifold M is given by (M, s) , where $s: M \rightarrow D(M)$ is the inclusion of the 0-section. The simple homotopy type of the pair $(M, \partial M)$ is given by the square

$$\begin{array}{ccc} M & \xrightarrow{s} & D(M) \\ \uparrow & & \uparrow \\ \partial M & \xrightarrow{s|_{\partial M}} & D(\partial M), \end{array}$$

where $D(\partial M) \subseteq D(M)$ are the disc bundles from Construction 14.3, with CW structures arising from a choice of PL triangulations corresponding to the PL structures.

By the commutative square at the end of Construction 14.3, the simple homotopy type of $(M, \partial M)$ is well-defined. Here we use that PL isomorphisms are simple: for any choice of triangulations underpinning the PL structures, the resulting homeomorphism is a simple homotopy equivalence. Also stabilising by D^s does not change the simple homotopy type, since as PL manifolds $D^s \cong D^{s-1} \times [-1, 1]$, and $D^{s-1} \times \{0\} \rightarrow D^{s-1} \times [-1, 1]$ is a simple equivalence.

Remark 14.5. Why is the simple homotopy type of ∂M obtained in this way the same as that obtained by applying Construction 14.3 with ∂M considered as a manifold without boundary?

For suitably high n , we may assume that the embedding of $(M, \partial M)$ into \mathbb{R}^n is isotopic, and thus by Theorem 2.20 ambiently isotopic, to an embedding with $i: \partial M \hookrightarrow \{\vec{x} \in \mathbb{R}^n \mid x_1 = 0\} \cong \mathbb{R}^{n-1}$ and an (interior) collar $\partial M \times [0, 1]$ embedded as a product in $\{\vec{x} \in \mathbb{R}^n \mid 0 \leq x_1 \leq 1\}$ with $(x, t) \mapsto (i(x), t)$, as in Theorem 2.21. Such an isotopy does not affect the simple homotopy type obtained, by the argument sketched above, which can also be found on [KS77, p. 123]. The simple homotopy type of ∂M obtained from Construction 14.3, via an embedding of ∂M into \mathbb{R}^{n-1} , uses a disc bundle $D(\partial M)$ that stabilises using the x_1 direction to a disc bundle $D'(\partial M)$, with fibre a disc of one dimension higher, for ∂M embedded in \mathbb{R}^n . This latter disc bundle gives rise to the canonical simple homotopy type of ∂M from Definition 14.4.

Remark 14.6. If M is a smooth manifold, then M has an underlying PL structure, and with a bit more care in Construction 14.3, we can arrange that the bundle $D(M)$ is a PL

bundle. Note that this is stronger than just a PL structure on the total space. For PL bundles, the bundle projection $D(M) \rightarrow M$ is a simple homotopy equivalence. Indeed, for trivial bundles this is discussed above, and in general the projection is an α -equivalence (a notion defined in [Fer77]) for any cover α of M and so is simple [Fer77, Corollary 3.2]. It follows that the simple homotopy type defined by (M, Id) agrees with the one of (M, s) , and the same holds for the relative simple homotopy type of the pair $(M, \partial M)$.

According to [KS77, Essay III, Theorem 5.11, p. 123], if a manifold has a triangulation, then the simple homotopy type of the manifold agrees with the simple homotopy type of that triangulation. It is not clear to us whether the analogous statement holds if M has a CW structure not coming from a triangulation.

14.2. THE CELLULAR CHAIN COMPLEX AND POINCARÉ TRIADS

Throughout this section let M be a compact connected n -manifold. Furthermore assume that we are given a decomposition $\partial M = R_- \cup R_+$ into codimension zero submanifolds such that $\partial R_- = R_- \cap R_+ = \partial R_+$.

The following proposition follows from the argument of Construction 14.3, applied with even more iterations to deal with corners of corners. See also the proof of [KS77, Essay III, Theorem 5.13, p. 136].

Proposition 14.7. *There exists a finite CW complex triad (X, X_-, X_+) and a homotopy equivalence of triads $f: (M, R_-, R_+) \rightarrow (X, X_-, X_+)$ such that the following two statements hold:*

- (1) *The restrictions of f to M , R_\pm and $R_- \cap R_+$ give the simple homotopy types of these manifolds, as defined in Definition 14.1.*
- (2) *The restrictions of f to the pairs $(M, \partial M)$, $(\partial M, R_\pm)$ and $(R_\pm, R_- \cap R_+)$ give the simple homotopy types of these pairs of manifolds, as defined in the previous section.*

We continue with a general definition regarding CW complexes.

Definition 14.8. Let (X, Y) be a pair of CW complexes such that X is connected. We write $\pi = \pi_1(X)$ and we denote the universal covering by $p: \tilde{X} \rightarrow X$. The group π acts on the left on the cells of the CW complex $(\tilde{X}, p^{-1}(Y))$. This equips $C_*^{\text{cell}}(\tilde{X}, p^{-1}(Y))$ with the structure of a left $\mathbb{Z}[\pi]$ -module. We define

$$\begin{aligned} C_*^{\text{cell}}(X, Y; \mathbb{Z}[\pi]) &:= \mathbb{Z}[\pi] \otimes_{\mathbb{Z}[\pi]} C_*^{\text{cell}}(\tilde{X}, p^{-1}(Y)) \\ C_{\text{cell}}^*(X, Y; \mathbb{Z}[\pi]) &:= \text{Hom}_{\text{right-}\mathbb{Z}[\pi]}(C_*^{\text{cell}}(\tilde{X}, p^{-1}(Y)), \mathbb{Z}[\pi]). \end{aligned}$$

Here, given a left $\mathbb{Z}[\pi]$ -module M we denote the right $\mathbb{Z}[\pi]$ -module given by $m \cdot g := g^{-1} \cdot m$ by \bar{M} . Note that the group π acts freely on the left on the cells of the CW complex $(\tilde{X}, p^{-1}(Y))$. For each cell in $X \setminus Y$, pick a lift to \tilde{X} . This turns $C_*^{\text{cell}}(X, Y; \mathbb{Z}[\pi])$ and $C_{\text{cell}}^*(X, Y; \mathbb{Z}[\pi])$ into based left $\mathbb{Z}[\pi]$ -module (co-) chain complexes.

Now we can state the main theorem of this section.

Theorem 14.9. *The finite CW complex triad (X, X_-, X_+) from Proposition 14.7 is a simple Poincaré triad, meaning that there is a chain level representative $\sigma \in C_n^{\text{cell}}(X, X_- \cup X_+)$ of the fundamental class $[X] \in H_n(X, X_+ \cup X_-; \mathbb{Z}) \cong H_n(M, \partial M; \mathbb{Z})$ such that*

$$- \cap \sigma: C_{\text{cell}}^{n-r}(X, X_-; \mathbb{Z}[\pi_1(X)]) \rightarrow C_r^{\text{cell}}(X, X_+; \mathbb{Z}[\pi_1(X)])$$

is a simple chain homotopy equivalence.

The theorem is proved in [KS77, Essay III, Theorem 5.13, p. 136]. In the Universal Poincaré Duality Theorem A.16 we will prove that there exists a chain homotopy equivalence between the two chain complexes. But we will not prove that there exists a *simple* homotopy equivalence; for that the reader will need to consult [KS77].

14.3. REIDEMEISTER TORSION

In this section we introduce Reidemeister torsion invariants for compact manifolds and discuss some of the key properties of these invariants.

Let M be a compact connected n -manifold and write $\pi = \pi_1(M)$. Let R_- be a compact codimension 0 submanifold of ∂M . In many applications $R_- = \emptyset$ or $R_- = \partial M$. We write $R_+ = \overline{\partial M} \setminus R_-$. Let F be a field and let $\alpha: \pi \rightarrow \text{GL}(d, F)$ be a representation of the fundamental group of M . With respect to this representation, we consider the twisted homology $H_k(M, R_-; F^d)$, as defined in Section A.1.

Assumption 14.10. *Suppose that $H_k(M, R_-; F^d) = 0$ for all k .*

Pick a homotopy equivalence of triads $f: (M, R_-, R_+) \rightarrow (X, X_-, X_+)$ as in Proposition 14.7. We use the homotopy equivalence f to make the identification $\pi_1(X) = \pi$. By a serious abuse of notation, we refer to the cellular chain complex $C_*^{\text{cell}}(X, X_-; \mathbb{Z}[\pi])$ of (X, X_-) as the *cellular chain complex* $C_*^{\text{cell}}(M, R_-; \mathbb{Z}[\pi])$ of (M, R_-) . As in Section 14.2 we view $C_*^{\text{cell}}(M, R_-; \mathbb{Z}[\pi])$ as a based left $\mathbb{Z}[\pi]$ -module chain complex. Equip the F -module chain complex $C_*^{\text{cell}}(M, R_-; F^d) = F^d \otimes_{\mathbb{Z}[\pi]} C_*^{\text{cell}}(M, R_-; \mathbb{Z}[\pi])$ with the basing given by the tensor products of the $\mathbb{Z}[\pi]$ -bases of $C_*^{\text{cell}}(M, R_-; \mathbb{Z}[\pi])$ and the canonical F -basis for F^d .

We write \sim_α for the equivalence relation on $F^\times := F \setminus \{0\}$ that is given by the subgroup $\{\pm \det(\alpha(g)) \mid g \in \pi_1(M)\} \subseteq F^\times$. We define $\tau(M, R_-, \alpha) \in F^\times / \sim_\alpha$ to be the Reidemeister torsion of the above acyclic, based F -module chain complex. We refer to [Tur01, Section 6] for the definition of the Reidemeister torsion of an acyclic, based F -module chain complex. It follows from a slight generalisation of [Tur01, Theorem 9.1] that $\tau(M, R_-, \alpha) \in F^\times / \sim_\alpha$ is well-defined, in that it is independent of the choice of the representative of the simple homotopy type of (X, X_-, X_+) and it is independent of the choice of the lifts of the cells. If $R_- = \emptyset$ then we write $\tau(M, \alpha) := \tau(M, \emptyset, \alpha)$.

The following two theorems give the two arguably most important properties of Reidemeister torsion.

Theorem 14.11. *Let M be a compact connected n -manifold, let R_- be a compact codimension zero submanifold of ∂M and let $\alpha: \pi_1(M) \rightarrow \text{GL}(d, F)$ be a representation. Let*

R_-^1, \dots, R_-^m be the components of R_- . By abuse of notation we also write α for the composition $\alpha: \pi_1(R_-^i) \rightarrow \pi_1(M) \rightarrow \mathrm{GL}(d, F)$ defined using a path from the base point of M to a base point of R_-^i . If $H_*(R_-^1; F^d) = \dots = H_*(R_-^m; F^d) = H_*(M; F^d) = 0$, then

$$\tau(M, \alpha) = \prod_{i=1}^m \tau(R_-^i, \alpha) \cdot \tau(M, R_-, \alpha) \in F^\times / \sim_\alpha.$$

PROOF. We have the following short exact sequence of chain complexes with compatible bases:

$$0 \rightarrow C_*^{\mathrm{cell}}(X_-; F^d) \rightarrow C_*^{\mathrm{cell}}(X; F^d) \rightarrow C_*^{\mathrm{cell}}(X, X_-; F^d) \rightarrow 0.$$

Given such a short exact sequence, the multiplicativity of the torsion is proven in [Tur01, Theorem 3.4]. \square

Definition 14.12. Let F be a field with (possibly trivial) involution. Given a representation $\alpha: \pi \rightarrow \mathrm{GL}(d, F)$ we denote the representation $g \mapsto \overline{\alpha(g^{-1})}^T$ by α^\dagger . We say that α is *unitary* if $\alpha = \alpha^\dagger$.

Example 14.13. Let $\phi: \pi \rightarrow \mathbb{Z}$ be a group homomorphism. Equip $\mathbb{Q}(t)$ with the usual involution given by $\bar{t} = t^{-1}$. The representation $\alpha: \pi \rightarrow \mathrm{GL}(1, \mathbb{Q}(t))$ given by $g \mapsto t^{\phi(g)}$ is unitary.

Theorem 14.14. Let M be a compact n -manifold with (possibly empty) boundary. Assume that we are given a decomposition $\partial M = R_- \cup R_+$ into codimension zero submanifolds such that $\partial R_- = R_- \cap R_+ = \partial R_+$. Furthermore let F be a field with (possibly trivial) involution. Let $\alpha: \pi_1(M) \rightarrow \mathrm{GL}(d, F)$ be a representation such that $H_*(\partial M; F^d) = 0 = H_*(M; F^d)$. Then

$$\tau(M, R_-, \alpha) = \overline{\tau(M, R_+, \alpha^\dagger)}^{(-1)^{n+1}} \in F^\times / \sim_\alpha.$$

In particular, if α is unitary we have

$$\tau(M, R_-, \alpha) = \overline{\tau(M, R_+, \alpha)}^{(-1)^{n+1}} \in F^\times / \sim_\alpha.$$

PROOF. We write $\pi = \pi_1(M)$. Write $C_*^\pm = C_*^{\mathrm{cell}}(M, R_\pm; \mathbb{Z}[\pi])$, recalling the convention described below Assumption 14.10.

It follows from Theorem 14.9 that the torsion of the based F -module chain complex $F^d \otimes_{\mathbb{Z}[\pi]} C_*^-$ agrees with the torsion of the based F -module chain complex

$$F^d \otimes_{\mathbb{Z}[\pi]} \mathrm{Hom}_{\mathrm{right-}\mathbb{Z}[\pi]}(\overline{C_{n-*}^+}, \mathbb{Z}[\pi]).$$

Consider the following isomorphism of based left F -module chain complexes

$$\begin{aligned} F_\alpha^d \otimes_{\mathbb{Z}[\pi]} \mathrm{Hom}_{\mathrm{right-}\mathbb{Z}[\pi]}(\overline{C_{n-*}^+}, \mathbb{Z}[\pi]) &\rightarrow \overline{\mathrm{Hom}_{\mathrm{left-}F}(F_{\alpha^\dagger}^d \otimes_{\mathbb{Z}[\pi]} C_{n-*}^+, F)} \\ v \otimes \varphi &\mapsto \left(\begin{array}{ccc} F_{\alpha^\dagger}^d \otimes_{\mathbb{Z}[\pi]} C_{n-*}^+ & \rightarrow & F \\ (w \otimes \sigma) & \mapsto & \overline{v\alpha(\varphi(\sigma))\bar{w}^T} \end{array} \right) \end{aligned}$$

Using this isomorphism $\tau(M, R_-, \alpha)$ also equals the torsion of the chain complex on the right hand side. It follows from algebraic duality for torsions [**Tur01**, Theorem 1.9] that the torsion of the based chain complex on the right hand side equals $\overline{\tau(M, R_+, \alpha^\dagger)}^{(-1)^{n+1}}$. \square

Obstructions to being topologically slice

15.1. THE FOX-MILNOR THEOREM

In this section we provide an example of the use of many of the theorems described in the previous chapters by applying them to obtain an obstruction for a knot to be topologically slice.

Definition 15.1. Let Y be a homology 3-sphere that is the boundary of an integral homology 4-ball X .

- (1) We say a knot K in Y is *topologically slice in X* if K bounds a *slice disc*, that is a proper submanifold of X homeomorphic to a disc.
- (2) Suppose X is equipped with a smooth structure, e.g. $X = D^4$. We say a knot K in Y is *smoothly slice in X* if K bounds a *smooth slice disc*, that is a proper smooth submanifold of X diffeomorphic to a disc.

There are many classical obstructions to a knot in S^3 being smoothly slice in D^4 . For example, there are obstructions based on the Alexander polynomial [FM66] and the Levine-Tristram signatures [Tri69, Lev69] and there are the more subtle Casson-Gordon [CG78, CG86] obstructions. It is not hard to see that these results also apply if we replace S^3 by any integral homology 3-sphere and if we replace D^4 by any smooth homology 4-ball.

Even though these results, having appeared prior to the work of Freedman and Quinn, were formulated as obstructions to being smoothly slice, it has been understood for many years that the original proofs can be modified to prove that these are in fact obstructions to being topologically slice.

In this section we will prove a sample theorem on the Alexander polynomial $\Delta_K(t)$ of a knot K . (On page 127 we recall the definition of the Alexander polynomial of a knot.) The following theorem, which in the smooth setting was first proved by Fox-Milnor [FM66], is arguably the most basic obstruction to a knot being topologically slice knot.

Theorem 15.2. (Fox-Milnor) *Suppose that K is a knot in a homology 3-sphere Y that bounds an integral homology 4-ball X . If K is topologically slice in X , then the Alexander polynomial $\Delta_K(t)$ of K factors as $\Delta_K(t) = \pm t^k \cdot f(t) \cdot f(t^{-1})$ for some $k \in \mathbb{Z}$ and for some $f(t) \in \mathbb{Z}[t, t^{-1}]$ such that $f(1) = \pm 1$.*

Even though this result is very well known we want to provide a detailed proof. In particular we want to highlight where some of the results discussed in this book are used.

The reader is encouraged to go through the above papers [Tri69, Lev69, CG78, CG86] and to modify the proofs to deal with topologically slice knots.

15.2. A PROOF OF THE FOX-MILNOR THEOREM

For the proof of the Fox-Milnor Theorem 15.2 we adopt the following notation.

- (1) Let Y be a homology 3-sphere bounding some integral homology 4-ball X .
- (2) Given a knot K in Y , denote its zero framed surgery by N_K .
- (3) Given an oriented knot K let μ_K be an oriented meridian.
- (4) For a slice disc D in X , let $N(D)$ be a tubular neighbourhood provided by Theorem 5.5. We refer to $W_D = X \setminus N(D)$ as the *exterior of D* .
- (5) The ring of integral Laurent polynomials in one variable is denoted $\mathbb{Z}[t, t^{-1}]$ or $\mathbb{Z}[t^{\pm 1}]$.

Many topological slicing obstructions, such as knot signatures [Tri69], the Fox-Milnor condition [FM66], the Blanchfield form [Kea75], Casson-Gordon invariants [CG78, CG86], L^2 -signature defects [COT03] and $L^{(2)}$ -von Neumann ρ -invariants [CT07], rely implicitly and explicitly on the next three propositions or slight variations thereof.

Proposition 15.3. *Let K be an oriented knot in Y and let D be a slice disc in X .*

- (1) *We have $\partial W_D = N_K$.*
- (2) *The inclusion map $\mu_K \rightarrow W_D$ induces a \mathbb{Z} -homology equivalence.*

In the remainder of this section, given an oriented knot, we use $\phi: \pi_1(N_K) \rightarrow \langle t \rangle$ and $\phi: \pi_1(W_D) \rightarrow \langle t \rangle$ to denote the unique homomorphisms that send the oriented meridian to t . These homomorphisms allow us to view $\mathbb{Z}[t^{\pm 1}]$ and $\mathbb{Q}(t)$ as a $\mathbb{Z}[\pi_1(N_K)]$ -module and a $\mathbb{Z}[\pi_1(W_D)]$ -module.

PROOF. First note that it follows from Proposition 5.10 (or more directly, the fact that D is contractible) that the tubular neighbourhood of D is trivial, thus we can identify it with $D \times D^2$.

- (1) We have to check that the framing of K induced by the unique trivialisation $N(D) \cong D^2 \times D$ is the 0-framing. Consider the double $DX = X \cup_Y X$. Note that Y splits DX into two copies of X . Let D be contained in one copy of X , and push a Seifert surface Σ into the other copy of X . By picking a collar neighbourhood for $Y = \partial X$ we obtain a bicollar neighbourhood $Y \times [-1, 1] \subseteq DX$. Using Theorem 2.21 we can arrange that $D \cap (Y \times [-1, 1]) = K \times [-1, 0]$, and $\Sigma \cap (Y \times [0, 1]) = K \times [0, 1]$. Let $F = \Sigma \cup -D \subseteq DX$. We compute the Euler number $e(F) \in \mathbb{Z}$ in two ways. First, note that $e(F) = [F] \cdot [F] = 0$, since $H_2(DX; \mathbb{Z}) = 0$. On the other hand, the number $e(F)$ is also the difference between the induced framings of $N(\Sigma)|_K$ and $N(D)|_K$. Consequently, the two framings agree and $N(D)$ induces the 0-framing, which by definition is the framing induced by $N(\Sigma)|_K$.
- (2) Let $\mu_K \rightarrow W_D$ be the inclusion of the meridian μ_K of K . Then we have $H_*(W_D, * \times \mu_K; \mathbb{Z}) = H_*(W_D, D \times S^1; \mathbb{Z}) = H_*(X, D \times D^2) = 0$ by excision and the hypothesis that X be a homology 4-ball. By the homology long exact sequence for the pair

(W_D, μ_K) , the meridional map $\mu_K \rightarrow W_D$ induces a homology equivalence, so W_D is a homology circle. \square

Proposition 15.4. *The exterior W_D of a slice disc D is homotopy equivalent to a finite 3-dimensional CW complex. In particular the homology groups*

$$H_*(W_D; \mathbb{Z}[t^{\pm 1}]), H_*(W_D, N_K; \mathbb{Z}[t^{\pm 1}]) \text{ and } H_*(N_K; \mathbb{Z}[t^{\pm 1}])$$

are all finitely generated.

PROOF. Note that W_D is a compact 4-manifold with nonempty boundary. It follows from Theorem 3.16 that W_D is homotopy equivalent to a 3-dimensional CW complex. The statements regarding the homology groups follow from Proposition A.9. \square

Proposition 15.5.

- (1) *For any knot K in a homology 3-sphere the modules $H_*(N_K; \mathbb{Z}[t^{\pm 1}])$ are $\mathbb{Z}[t^{\pm 1}]$ -torsion.*
- (2) *If D is a slice disc, then all the modules $H_*(W_D; \mathbb{Z}[t^{\pm 1}])$ are $\mathbb{Z}[t^{\pm 1}]$ -torsion.*

PROOF. We start out with the proof of the second statement. Let $P \subseteq \mathbb{Z}[t^{\pm 1}]$ be the multiplicative subset of Laurent polynomials that augment to ± 1 , that is $p(1) = \pm 1$ and only if $p \in P$. We shall prove the slightly stronger statement, that $H_k(W_D; \mathbb{Z}[t^{\pm 1}])$ is P -torsion for $k > 0$. Since $H_0(W_D; \mathbb{Z}[t^{\pm 1}]) \cong \mathbb{Z}[t^{\pm 1}]/(t - 1)$ is $\mathbb{Z}[t^{\pm 1}]$ -torsion, the result will follow. We write $\pi = \pi_1(W_D)$. Let

$$Q := P^{-1}\mathbb{Z}[t^{\pm 1}]$$

be the result of inverting the polynomials in P . By Proposition A.5 there exists a chain complex C_* of finite length consisting of finitely generated free left $\mathbb{Z}[\pi]$ -modules such that for any ring R and any $(R, \mathbb{Z}[\pi])$ -bimodule A we have

$$H_k(W_D, \mu_K; A) \cong H_k(A \otimes_{\mathbb{Z}[\pi]} C_*).$$

By Proposition 15.3 we know that $H_k(\mathbb{Z} \otimes_{\mathbb{Z}[\pi]} C_*) = H_k(W_D, \mu_K; \mathbb{Z}) = 0$. Since C_* is a chain complex of finite length consisting of finitely generated free left $\mathbb{Z}[\pi]$ -modules we obtain from chain homotopy lifting [COT03, Proposition 2.10], see also [NP17, Lemma 3.1], that $H_k(Q \otimes_{\mathbb{Z}[\pi]} C_*) = 0$. A straightforward calculation shows that $H_*(S^1, \text{pt}; Q) = 0$. It follows that $H_*(W_D, \text{pt}; Q) = 0$, so that $H_k(W_D; \mathbb{Z}[t^{\pm 1}])$ is P -torsion for $k > 0$.

The first statement is very well known. One of the many proofs would be to use the above argument and the fact that $S^1 \rightarrow Y \setminus \nu K$ is a homology equivalence to show that the modules $H_*(Y \setminus \nu K; \mathbb{Z}[t^{\pm 1}])$ are torsion. A basic Mayer-Vietoris argument then shows that the modules $H_*(N_K; \mathbb{Z}[t^{\pm 1}])$ are also torsion. \square

We want to recall the definition of the Alexander polynomial of a knot. To do so we need the notion of the order of a module.

Definition 15.6. Let H be a finitely generated free abelian group and let M be a finitely generated $\mathbb{Z}[H]$ -module. By [Lan02a, Corollary IV.9.5] the ring $\mathbb{Z}[H]$ is Noetherian which implies that M admits a free resolution

$$\mathbb{Z}[H]^r \xrightarrow{A} \mathbb{Z}[H]^s \rightarrow M \rightarrow 0.$$

Without loss of generality we can assume that $r > s$. Since $\mathbb{Z}[H]$ is unique factorisation domain, see [Lan02a, Lemma IV.2.3], the order $\text{ord}(M)$ is defined as the greatest common divisor of the $s \times s$ -minors of A . By [Tur01, Lemma 4.4] the order is well-defined, i.e. independent of the choice of the free resolution, up to multiplication by a unit in $\mathbb{Z}[H]$.

The fact that the order is only well-defined up to multiplication by a unit leads us to the following notation:

Notation 15.7. Let H be a free abelian group. Given $p, q \in \mathbb{Z}[H]$ we write $p \doteq q$ if $p = \pm h \cdot q$ for some $h \in H$.

In the proof of the Fox-Milnor Theorem we will need the following lemma, collecting basic facts about orders of finitely generated $\mathbb{Z}[H]$ -modules.

Lemma 15.8. *Let H be a finitely generated free abelian group.*

- (1) *If $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$ is a short exact sequence of finitely generated $\mathbb{Z}[H]$ -modules, then $\text{ord}(B) \doteq \text{ord}(A) \cdot \text{ord}(C)$.*
- (2) *If $0 \rightarrow C_k \rightarrow \cdots \rightarrow C_0 \rightarrow 0$ is an exact sequence of finitely generated $\mathbb{Z}[H]$ -torsion modules, then the alternating product of the orders is a unit in $\mathbb{Z}[H]$.*
- (3) *For any finitely generated $\mathbb{Z}[H]$ -module A we have $\text{ord}(\overline{A}) \doteq \overline{\text{ord}(A)}$. Here we write $f \mapsto \overline{f}$ for the natural involution on $\mathbb{Z}[H]$ given by $h \mapsto h^{-1}$ for $h \in H$, and given a $\mathbb{Z}[H]$ -module A we write \overline{A} for the $\mathbb{Z}[H]$ -module given by $f \cdot a := \overline{f} \cdot a$.*
- (4) *For any finitely generated torsion $\mathbb{Z}[H]$ -module A we have $\text{ord}(\text{Ext}_{\mathbb{Z}[H]}^1(A, \mathbb{Z}[H])) \doteq \text{ord}(A)$.*

PROOF. Statement (1) is proven for $H \cong \mathbb{Z}$ in [Lev67, Lemma 5]. The general case follows from [Hil12, Theorem 3.12]. Note that (2) is an immediate consequence of (1), by separating the long exact sequence into short exact sequences such as $0 \rightarrow \text{Im } C_j \rightarrow C_{j-1} \rightarrow \text{Im } C_{j-1} \rightarrow 0$, applying (1), and performing substitutions using the resulting equations involving orders.

Next (3) follows immediately from the definition. Finally (4) is well-known to the experts, but we could not find a reference, therefore we sketch the key ingredients in the proof. We introduce the following notation.

- (a) Given any prime ideal \mathfrak{p} of $\mathbb{Z}[H]$, let $\mathbb{Z}[H]_{\mathfrak{p}}$ be the localisation at \mathfrak{p} , that is we invert all elements that do not lie in \mathfrak{p} . We view $\mathbb{Z}[H]$ as a subring of $\mathbb{Z}[H]_{\mathfrak{p}}$.
- (b) Given a ring R and $f, g \in R$ we write $f \doteq_R g$ if f and g differ by multiplication by a unit in R .

Now we sketch the proof of (4). We will use the following five observations.

- (i) Since $\mathbb{Z}[H]$ is a unique factorisation domain, for any prime element $p \in \mathbb{Z}[H]$ the ideal (p) is a prime ideal.
- (ii) Being a unique factorisation domain and being Noetherian are preserved under localisation [Pes96, Theorem 7.53], [Row06, Corollary 8.8']. In particular each $\mathbb{Z}[H]_{\mathfrak{p}}$ is a Noetherian unique factorisation domain. This allows us, by the same definitions as above, to define the order of a finitely generated module over $\mathbb{Z}[H]_{\mathfrak{p}}$.

- (iii) Localisation is flat [**Lan02a**, Proposition XVI.3.2]. It follows that for any finitely generated $\mathbb{Z}[H]$ -module M and any prime element $p \in \mathbb{Z}[H]$ one has $\text{ord}(M) \doteq_{\mathbb{Z}[H]_{(p)}} \text{ord}(\mathbb{Z}[H]_{(p)} \otimes_{\mathbb{Z}[H]} M)$ and $\mathbb{Z}[H]_{(p)} \otimes_{\mathbb{Z}[H]} \text{Ext}_{\mathbb{Z}[H]}^1(M, \mathbb{Z}[H]) \cong \text{Ext}_{\mathbb{Z}[H]_{(p)}}^1(\mathbb{Z}[H]_{(p)} \otimes_{\mathbb{Z}[H]} M, \mathbb{Z}[H]_{(p)})$ as $\mathbb{Z}[H]_{(p)}$ -modules.
- (iv) By [**Os00**, Corollary A.14] every commutative ring with the property that every prime ideal is principal, is a PID. It follows easily that for each prime element p , the localisation $\mathbb{Z}[H]_{(p)}$ is a PID.
- (v) Let L be a torsion $\mathbb{Z}[H]_{(p)}$ -module. Since $\mathbb{Z}[H]_{(p)}$ is a PID every two elements have a greatest common divisor. We can therefore perform row and column operations to find a resolution for L such that the presentation matrix is diagonal. From this observation one easily deduces that $L \cong \text{Ext}_{\mathbb{Z}[H]_{(p)}}^1(L, \mathbb{Z}[H]_{(p)})$ as left $\mathbb{Z}[H]_{(p)}$ -modules. Since L is torsion the presentation matrix is injective and so its transpose presents the Ext group. To convert the Ext group to a left module, we use the trivial involution, which we may do since $\mathbb{Z}[H]_{(p)}$ is a commutative ring.
- (vi) Suppose that f and g are in $\mathbb{Z}[H]$. If $f \doteq_{\mathbb{Z}[H]_{(p)}} g$ for all prime elements $p \in \mathbb{Z}[H]$, then since $\mathbb{Z}[H]$ is a unique factorisation domain we must have $f \doteq_{\mathbb{Z}[H]} g$.

Now with $L = \mathbb{Z}[H]_{(p)} \otimes_{\mathbb{Z}[H]} A$ a finitely generated $\mathbb{Z}[H]$ -torsion module, we have

$$\mathbb{Z}[H]_{(p)} \otimes_{\mathbb{Z}[H]} A \cong \text{Ext}_{\mathbb{Z}[H]_{(p)}}^1(\mathbb{Z}[H]_{(p)} \otimes_{\mathbb{Z}[H]} A, \mathbb{Z}[H]_{(p)})$$

for every prime element p , by (iv). On the other hand, again for each prime element p , we have

$$\text{ord}(A) \doteq_{\mathbb{Z}[H]_{(p)}} \text{ord}(\mathbb{Z}[H]_{(p)} \otimes_{\mathbb{Z}[H]} A)$$

by (iii). Combining these two observations yields

$$\text{ord}(A) \doteq_{\mathbb{Z}[H]_{(p)}} \text{ord}(\text{Ext}_{\mathbb{Z}[H]_{(p)}}^1(\mathbb{Z}[H]_{(p)} \otimes_{\mathbb{Z}[H]} A, \mathbb{Z}[H]_{(p)})).$$

By the second part of (iii) we have that

$$\text{ord}(\text{Ext}_{\mathbb{Z}[H]_{(p)}}^1(\mathbb{Z}[H]_{(p)} \otimes_{\mathbb{Z}[H]} A, \mathbb{Z}[H]_{(p)})) \doteq_{\mathbb{Z}[H]_{(p)}} \text{ord}(\mathbb{Z}[H]_{(p)} \otimes_{\mathbb{Z}[H]} \text{Ext}_{\mathbb{Z}[H]}^1(A, \mathbb{Z}[H])).$$

By the first part of (iii) again we have

$$\text{ord}(\mathbb{Z}[H]_{(p)} \otimes_{\mathbb{Z}[H]} \text{Ext}_{\mathbb{Z}[H]}^1(A, \mathbb{Z}[H])) \doteq_{\mathbb{Z}[H]_{(p)}} \text{ord}(\text{Ext}_{\mathbb{Z}[H]}^1(A, \mathbb{Z}[H])).$$

Thus combining the last three equalities we have

$$\text{ord}(A) \doteq_{\mathbb{Z}[H]_{(p)}} \text{ord}(\text{Ext}_{\mathbb{Z}[H]}^1(A, \mathbb{Z}[H]))$$

for all prime elements p . Now (4) follows by applying (vi). \square

We use the notion of order to define the Alexander polynomial of a knot in a homology 3-sphere.

Definition 15.9. The *Alexander polynomial* $\Delta_K(t)$ of a knot K is defined as the order of the Alexander module $H_1(N_K; \mathbb{Z}[t^{\pm 1}])$. Note that this polynomial is only well-defined up to units in $\mathbb{Z}[t^{\pm 1}]$.

After these preparations we turn to the actual proof of the Fox-Milnor Theorem 15.2. We need the following elementary lemma.

Lemma 15.10. *Let π be a group, let C_* be a chain complex of left free $\mathbb{Z}[\pi]$ -modules and let $\phi: \pi \rightarrow \langle t \rangle$ be a homomorphism. The map*

$$\begin{aligned} \text{Hom}_{\text{right-}\mathbb{Z}[\pi]}(\overline{C_*}, \mathbb{Z}[t^{\pm 1}]) &\rightarrow \overline{\text{Hom}_{\text{left-}\mathbb{Z}[t^{\pm 1}]}(\mathbb{Z}[t^{\pm 1}] \otimes_{\mathbb{Z}[\pi]} C_*, \mathbb{Z}[t^{\pm 1}])} \\ f &\mapsto (p \otimes \sigma \mapsto p \cdot f(\sigma)) \end{aligned}$$

is well-defined and is an isomorphism of left $\mathbb{Z}[t^{\pm 1}]$ -cochain complexes. \square

FIRST PROOF OF THE FOX-MILNOR THEOREM 15.2. In this proof we abbreviate $\Lambda := \mathbb{Z}[t^{\pm 1}]$. We start out with the following three observations.

- (a) We have $H_0(W_D; \Lambda) \cong H_0(N_K; \Lambda) \cong \Lambda/(t-1)$.
- (b) We have $H_0(W_D, N_K; \Lambda) = 0$.
- (c) By Proposition 15.5 and Proposition 15.4 we know that for all k

$$\text{Ext}_{\Lambda}^0(H_k(W_D, N_K; \Lambda), \Lambda) = \text{Hom}_{\Lambda}(H_k(W_D, N_K; \Lambda), \Lambda) = 0.$$

Claim. For any $i \in \mathbb{N}$ we have

$$\begin{aligned} H_i(N_K; \Lambda) &\cong \overline{\text{Ext}_{\Lambda}^1(H_{2-i}(N_K; \Lambda), \Lambda)} \\ H_i(W_D; \Lambda) &\cong \overline{\text{Ext}_{\Lambda}^1(H_{3-i}(W_D, N_K; \Lambda), \Lambda)}. \end{aligned}$$

We prove the second statement of the claim. The proof of the first statement is almost identical. By the Poincaré Duality Theorem A.15 we have an isomorphism $H_i(W_D; \Lambda) \cong H^{4-i}(W_D, N_K; \Lambda)$ of Λ -modules. By Lemma 15.10, applied to $C_* = C_*(W_D, N_K; \mathbb{Z}[\pi])$, we know that

$$H^{4-i}(W_D, N_K; \Lambda) \cong \overline{H_{4-i}(\text{Hom}_{\Lambda}(\Lambda \otimes_{\mathbb{Z}[\pi]} C_*(W_D, N_K; \mathbb{Z}[\pi]), \Lambda)}).$$

Finally we apply the universal coefficient spectral sequence [Lev77, Theorem 2.3] to the Λ -module chain complex $C_*(W_D, N_K; \Lambda)$. It follows from the above observations (b) and (c) that the spectral sequence collapses and that we have an isomorphism

$$H_{4-i}(\text{Hom}_{\Lambda}(\Lambda \otimes_{\mathbb{Z}[\pi]} C_*(W_D, N_K; \Lambda))) \cong \text{Ext}_{\Lambda}^1(H_{3-i}(W_D, N_K; \Lambda), \Lambda).$$

This concludes the proof of the claim.

Next we consider the long exact sequence of the pair (W_D, N_K) of twisted homology with Λ -coefficients:

$$\cdots \rightarrow H_2(W_D; \Lambda) \rightarrow H_2(W_D, N_K; \Lambda) \rightarrow H_1(N_K; \Lambda) \rightarrow H_1(W_D; \Lambda) \rightarrow H_1(W_D, N_K; \Lambda) \rightarrow \cdots$$

It follows from Propositions 15.4 that all the above modules are finitely generated. Thus it makes sense to consider their orders. Also note that in Proposition 15.5 we saw that the modules for N_K and W_D are all Λ -torsion. It follows from the long exact sequence that the relative homology groups $H_*(W_D, N_K; \Lambda)$ are also Λ -torsion. By Lemma 15.8 (3) the alternating product of the orders equals $\pm t^k$.

By the above claim and Lemma 15.8 (3) and (4) the orders are anti-symmetric around $H_1(N_K; \Lambda)$. More precisely, we have

$$\text{ord}(H_2(W_D, N_K; \Lambda)) \doteq \text{ord}(\overline{\text{Ext}_\Lambda^1(H_1(W_D; \Lambda), \Lambda)}) \doteq \overline{\text{ord}(H_1(W_D; \Lambda))},$$

and the same type of relation holds as we progress further from the middle term $H_1(N_K; \Lambda)$ in the above long exact sequence. But this implies that there exist nonzero polynomials $f, g \in \Lambda$ with $f \cdot \bar{f} \doteq \Delta_K(t) \cdot g \cdot \bar{g}$. It follows easily from the fact that $\Lambda = \mathbb{Z}[t^{\pm 1}]$ is a UFD that there exists an $h \in \Lambda$ with $h \cdot \bar{h} \doteq \Delta_K(t)$. \square

A knot K in Y is *homotopy ribbon* if there is a slice disc D in X such that $\pi_1(Y \setminus \nu K) \rightarrow \pi_1(W_D)$ is surjective.

Corollary 15.11. *Let $D \subseteq X$ be a homotopy ribbon disc for $K \subseteq Y$. Let $f(t) = \text{ord } H_1(W_D; \Lambda)$. Then $\Delta_K(t) \doteq f(t)f(t^{-1})$.*

PROOF. Since $\pi_1(S^3 \setminus \nu K) \rightarrow \pi_1(W_D)$ factors through $\pi_1(N_K)$, the map $\pi_1(N_K) \rightarrow \pi_1(W_D)$ is surjective. Hence $\pi_1(N_K)^{(1)} \rightarrow \pi_1(W_D)^{(1)}$, the map on commutator subgroups is surjective. The respective homology groups with Λ coefficients are the abelianisations of the commutator subgroups, and so $H_1(N_K; \Lambda) \rightarrow H_1(W_D; \Lambda)$ is surjective. Hence $H_1(W_D, N_K; \Lambda) = 0$, and so

$$1 = \text{ord}(H_1(W_D, N_K; \Lambda)) = \overline{\text{ord}(H_2(W_D; \Lambda))}.$$

The pervious proof then implies that

$$\Delta_K(t) \doteq \text{ord}(H_1(N_K; \Lambda)) \doteq \text{ord } H_1(W_D; \Lambda) \cdot \overline{\text{ord } H_1(W_D; \Lambda)} = f(t)f(t^{-1})$$

as required. \square

We conclude with an alternative argument for the Fox-Milnor Theorem in the topological category using Reidemeister torsion. The advantage of the Reidemeister torsion invariant is that proofs are often easier, and it has in general a smaller indeterminacy than the order of homology, although this will not manifest itself in the upcoming proof.

SECOND PROOF OF THEOREM 15.2. We continue with the notation introduced above. As before we have a homomorphism $\alpha: \pi_1(W_D) \rightarrow H_1(W_D; \mathbb{Z}) \xrightarrow{\cong} \mathbb{Z}$, sending an oriented meridian of K to $1 \in \mathbb{Z}$. As usual $\mathbb{Q}(t)$ denotes the field of fractions of the Laurent polynomial ring $\mathbb{Z}[t, t^{-1}]$. We take $d = 1$, and so obtain a representation $\phi: \pi_1(W_D) \rightarrow \text{GL}(1, \mathbb{Q}(t))$, that sends $g \mapsto (t^{\alpha(g)})$. In the previous proof we had already seen that the modules $H_*(N_K; \mathbb{Z}[t^{\pm 1}])$, $H_*(W_D; \mathbb{Z}[t^{\pm 1}])$ and $H_*(N_K; \mathbb{Z}[t^{\pm 1}])$ are $\mathbb{Z}[t^{\pm 1}]$ -torsion. Since $\mathbb{Q}(t)$ is flat over $\mathbb{Z}[t^{\pm 1}]$ it follows that the corresponding twisted homology groups with $\mathbb{Q}(t)$ -coefficients are zero.

By the discussion in Section 14.3 we can consider the Reidemeister torsions $\tau(W_D, \phi)$, $\tau(N_K, \phi)$ and $\tau(W_D, N_K, \phi)$. By Theorem 14.14, $\tau(W_D, N_K, \phi) \doteq \overline{\tau(W_D, \phi)^{(-1)^5}} \doteq \overline{\tau(W_D, \phi)^{-1}}$. Since the torsion is multiplicative in short exact sequences by Theorem 14.11, we have that

$$\tau(W_D, \phi) \doteq \tau(N_K, \phi) \cdot \tau(W_D, N_K, \phi) \doteq \tau(N_K, \phi) \cdot \overline{\tau(W_D, \phi)^{-1}}.$$

By [Tur01, Theorem 14.12] the torsion of the zero surgery of a knot is equal to $\Delta_K(t)/((t-1)(t^{-1}-1))$. It follows that $\Delta_K(t)$ is a norm as claimed. \square

Remark 15.12. The two proofs presented above avoid the use of the smooth category, and so are in keeping with the spirit of this book. However, one can give a further alternative proof by allowing smooth techniques. First one can use Theorem 9.9 to find a simply connected 4-manifold W' such that $W := W_D \# W'$ is smoothable. Then one can triangulate W and apply Reidemeister torsion machinery without appealing to [KS77, Essay III]. The disadvantage of this approach is that typically $H_2(W'; \mathbb{Z})$ will be nontrivial, so that W is not acyclic over $\mathbb{Q}(t)$. One can proceed by choosing a self-dual basis for homology, so that one can still obtain a torsion invariant that is well-defined up to norms. Apply [CF13, Theorem 2.4], and argue that since the intersection form of W is nonsingular, the contribution of W' to the torsion is a norm.

APPENDIX A

Poincaré Duality with twisted coefficients

Surveying the literature, we felt it would be beneficial to have a more detailed proof of Poincaré duality with twisted coefficients for manifolds with boundary, but without a smooth or PL structure, so we offer one in this appendix. One can find other proofs of Poincaré duality for some subsets of these conditions, e.g. Sun [Sun17] and Kwasik-Sun [KS18] provide a proof in the closed case. A detailed discussion of twisted (co-) homology and Poincaré duality can also be found in [Fri23, Part XXIV].

A.1. TWISTED HOMOLOGY AND COHOMOLOGY GROUPS

We start out with the following notation.

Notation A.1. Given a group π and a left $\mathbb{Z}[\pi]$ -module A , write \bar{A} for the right $\mathbb{Z}[\pi]$ -module that has the same underlying abelian group but for which the right action of $\mathbb{Z}[\pi]$ is defined by $a \cdot g := g^{-1} \cdot a$ for $a \in A$ and $g \in \pi$. The same notation is also used with the rôles of left and right reversed and $g \cdot a := a \cdot g^{-1}$.

We recall the definition of twisted homology and cohomology groups.

Definition A.2. Let X be a connected topological space that admits a universal cover $p: \tilde{X} \rightarrow X$. Write $\pi = \pi_1(X)$. Let Y be a subset of X and let A be a right $\mathbb{Z}[\pi]$ -module. Let π act on \tilde{X} by deck transformations, which is naturally a left action. Thus, the singular chain complex $C_*(\tilde{X}, p^{-1}(Y))$ becomes a left $\mathbb{Z}[\pi]$ -module chain complex. Define the *twisted chain complex*

$$C_*(X, Y; A) := (A \otimes_{\mathbb{Z}[\pi]} C_*(\tilde{X}, p^{-1}(Y)), \text{Id} \otimes \partial_*).$$

The corresponding *twisted homology groups* are $H_k(X, Y; A)$. With $\delta^k = \text{Hom}(\partial_k, \text{Id})$ define the *twisted cochain complex* to be

$$C^*(X, Y; A) := (\text{Hom}_{\text{right-}\mathbb{Z}[\pi]}(\overline{C_*(\tilde{X}, p^{-1}(Y))}, A), \delta^*).$$

The corresponding *twisted cohomology groups* are $H^k(X, Y; A)$.

Note that if R is some ring (not necessarily commutative) and if A is an $(R, \mathbb{Z}[\pi])$ -bimodule, then the above twisted homology and cohomology groups are naturally left R -modules.

Given a CW complex one can similarly define twisted cellular (co-) chain complexes and twisted cellular (co-) homology groups. The following proposition implies that twisted singular (co-) homology groups are isomorphic to twisted cellular (co-) homology groups.

Proposition A.3. *Let (X, Y) be a CW complex pair and write $\pi = \pi_1(X)$. The singular chain complex $C_*^{\text{sing}}(X, Y; \mathbb{Z}[\pi])$ and the cellular chain complex $C_*^{\text{cell}}(X, Y; \mathbb{Z}[\pi])$ are chain homotopy equivalent as chain complexes of left $\mathbb{Z}[\pi]$ -modules.*

The proof of Proposition A.3 relies on the following very useful lemma.

Lemma A.4. *Let $f: C_* \rightarrow D_*$ be a chain map of chain complexes of free left $\mathbb{Z}[\pi]$ -modules (here chain complexes are understood to start in degree 0) that induces an isomorphism on homology. Then f is a chain equivalence.*

PROOF. Since f induces an isomorphism of homology groups we know that the mapping cone $\text{cone}(f)_*$ is acyclic. By assumption C_* and D_* are free left $\mathbb{Z}[\pi]$ -modules. It follows that $\text{cone}(f)_*$ is also a chain complex of free left $\mathbb{Z}[\pi]$ -modules. But this guarantees the existence of a chain homotopy $\text{Id}_{\text{cone}(f)_*} \simeq_P 0$, since we can view $\text{cone}(f)_*$ as a free resolution of 0 and any two such resolutions are chain homotopic. Recall that chain homotopy means

$$(A.1) \quad \partial^{\text{cone}(f)_*} \circ P + P \circ \partial^{\text{cone} f_*} = \text{Id}_{\text{cone}(f)_*}$$

If we write P as a matrix

$$P_n = \begin{pmatrix} P_n^{11} & P_n^{12} \\ P_n^{21} & P_n^{22} \end{pmatrix} : C_{n-1} \oplus D_n \rightarrow C_n \oplus D_{n+1}$$

then one easily verifies using Equation (A.1), that $P_*^{12}: D_* \rightarrow C_*$ is a chain homotopy inverse of f_* , where the chain homotopies are given by P_*^{11} and P_*^{22} . \square

PROOF OF PROPOSITION A.3. Given a CW complex A we consider the intermediate chain complex $C_*^{\text{int}}(A) := \ker(C_n(A^n) \xrightarrow{\partial} C_{n-1}(A^n) \rightarrow C_{n-1}(A^n, A^{n-1}))$. Given a subcomplex B of A we set $C_*^{\text{int}}(A, B) := \text{coker}(C_n^{\text{int}}(A) \rightarrow C_n^{\text{int}}(B))$.

Let $p: \tilde{X} \rightarrow X$ denote the universal cover. We write $\tilde{Y} := p^{-1}(Y)$. In [Sch68, p. 303] (see also [Lüc98, Lemma 4.2]) it is shown that the natural maps $\iota: C_*^{\text{int}}(\tilde{X}, \tilde{Y}) \rightarrow C_*(\tilde{X}, \tilde{Y})$ and $\pi: C_*^{\text{int}}(\tilde{X}, \tilde{Y}) \rightarrow C_*^{\text{cell}}(\tilde{X}, \tilde{Y})$ induce isomorphisms of homology groups.

Note that $C_*(\tilde{X}, \tilde{Y})$ and $C_*^{\text{cell}}(\tilde{X}, \tilde{Y})$ are free $\mathbb{Z}[\pi]$ -left modules. But it is not clear whether each $C_*^{\text{int}}(\tilde{X}, \tilde{Y})$ is a free $\mathbb{Z}[\pi]$ -left module. But it is straightforward to show that there exists a chain complex F_* consisting of free $\mathbb{Z}[\pi]$ -left modules and a chain map $\varphi: F_* \rightarrow C_*^{\text{int}}(\tilde{X}, \tilde{Y})$ of left $\mathbb{Z}[\pi]$ -modules which induces isomorphisms of homology groups.

It follows from Lemma A.4 that $\iota \circ \varphi: F_* \rightarrow C_*(\tilde{X}, \tilde{Y})$ and $\pi \circ \varphi: F_* \rightarrow C_*^{\text{cell}}(\tilde{X}, \tilde{Y})$ are chain homotopy equivalences of left $\mathbb{Z}[\pi]$ -modules. It follows that $C_*(\tilde{X}, \tilde{Y})$ and $C_*^{\text{cell}}(\tilde{X}, \tilde{Y})$ are chain homotopy equivalent as chain complexes of left $\mathbb{Z}[\pi]$ -modules, which is equivalent to $C_*^{\text{sing}}(X, Y; \mathbb{Z}[\pi])$ and $C_*^{\text{cell}}(X, Y; \mathbb{Z}[\pi])$ being chain homotopy equivalent as chain complexes of left $\mathbb{Z}[\pi]$ -modules. \square

Proposition A.5. *Let M be a compact n -manifold and let $N \subseteq M$ be a subspace that is a compact manifold in its own right. Write $\pi = \pi_1(M)$. There exists a chain complex C_**

of finite length consisting of finitely generated free left $\mathbb{Z}[\pi]$ -modules such that for any ring R , for any $(R, \mathbb{Z}[\pi])$ -bimodule A and for any $k \in \mathbb{N}_0$ we have left R -module isomorphisms

$$H_k(M, N; A) \cong H_k(A \otimes_{\mathbb{Z}[\pi]} C_*)$$

and

$$H^k(M, N; A) \cong H_k(\text{Hom}_{\mathbb{Z}[\pi]}(\overline{C_*}, A)).$$

Remark A.6. Note that we do *not* demand that N be a submanifold of M . For example N could be a union of boundary components of M , or N could be a submanifold of the boundary. Evidently N could also be the empty set.

PROOF. By Theorem 3.16 the manifolds M and N are homotopy equivalent to finite CW complexes X and Y respectively. Let $i: N \rightarrow M$ be the inclusion map. By the Cellular Approximation Theorem there exists a cellular map $j: Y \rightarrow X$ such that the following diagram commutes up to homotopy:

$$\begin{array}{ccc} N & \xrightarrow{i} & M \\ \simeq \downarrow & & \downarrow \simeq \\ Y & \xrightarrow{j} & X. \end{array}$$

Next we replace M and X by the mapping cylinders of i and j respectively, to create cofibrations. Given a map $f: U \rightarrow V$ between topological spaces let $\text{cyl}(f)$ be the mapping cylinder. We view U as a subset of $\text{cyl}(f)$ in the obvious way. With this notation we have

$$H_k(M, N; A) \cong H_k(\text{cyl}(i: N \rightarrow M), N; A) \cong H_k(\text{cyl}(j: Y \rightarrow X), Y; A).$$

The mapping cylinder $Z := \text{cyl}(j: Y \rightarrow X)$ admits the structure of a finite CW complex such that Y is a subcomplex. Thus we can compute the twisted homology groups $H_k(\text{cyl}(j: X \rightarrow Y); A)$ using the relative twisted cellular chain complex, and similarly for cohomology. Put differently, $C_* = C_*^{\text{cell}}(Z, Y; \mathbb{Z}[\pi])$ has the desired properties. \square

In order to give a criterion for twisted homology modules to be finitely generated, we need the notion of a Noetherian ring.

Definition A.7. A ring R is said to be *left Noetherian* if for any descending chain

$$R \supseteq I_1 \supseteq I_2 \supseteq I_3 \supseteq \dots$$

of left R -ideals the inclusions eventually become equality. If R is commutative, then we just say Noetherian.

Example A.8. The following rings are left Noetherian:

- (1) The ring \mathbb{Z} is Noetherian.
- (2) Any (skew) field is left Noetherian.
- (3) If A is a commutative Noetherian ring, then the multivariable Laurent polynomial ring $A[t_1^{\pm 1}, \dots, t_k^{\pm 1}]$ is also Noetherian [**Lan02a**, Corollary IV.9.5].

The following theorem is often implicitly used.

Proposition A.9. *Let M be a compact n -manifold, let $N \subseteq M$ be a subspace that is a compact manifold in its own right, let R be a ring and let A be an $(R, \mathbb{Z}[\pi])$ -bimodule. If R is left Noetherian and if A is finitely generated as a left R -module, then all the twisted homology modules $H_*(M, N; A)$ are finitely generated left R -modules.*

In the proof of Proposition A.9 we will need the following lemma; cf. [Lam91, Proposition 1.21] or [Lan02a, Proposition X.1.4].

Lemma A.10. *Let R be a left Noetherian ring. If P is a finitely generated left R -module, then any left submodule of P is also a finitely generated left R -module.*

PROOF OF PROPOSITION A.9. By Proposition A.5, there exists a chain complex C_* of finite length consisting of finitely generated free left $\mathbb{Z}[\pi]$ -modules such that

$$H_k(M, N; A) \cong H_k(A \otimes_{\mathbb{Z}[\pi]} C_*).$$

Given $k \in \mathbb{N}_0$ we denote the rank of C_k as a free left $\mathbb{Z}[\pi]$ -module by r_k . Then we have $A \otimes_{\mathbb{Z}[\pi]} C_k \cong A \otimes_{\mathbb{Z}[\pi]} \mathbb{Z}[\pi]^{r_k} \cong A^{r_k}$. In particular $H_k(M, N; A)$ is isomorphic to a quotient of a submodule of a finitely generated R -module. The desired statement follows from Lemma A.10. \square

A.1.1. Cup and cap products on twisted (co-) chain complexes. Throughout this section let X be a connected topological space admitting a universal cover, and write $\pi = \pi_1(X)$. We want to introduce the cup product and the cap product on twisted (co-) chain complexes. Given an n -simplex σ , define the p -simplices $\sigma]_p$ and $\sigma]_p$ by

$$\begin{aligned} \sigma]_p(t_0, \dots, t_p) &:= \sigma(t_0, \dots, t_p, 0, \dots, 0), \\ \sigma]_p(t_0, \dots, t_p) &:= \sigma(0, \dots, 0, t_0, \dots, t_p). \end{aligned}$$

Throughout this section let A and B be two right $\mathbb{Z}[\pi]$ -modules A . We view $A \otimes_{\mathbb{Z}} B$ as a right $\mathbb{Z}[\pi]$ -module via the diagonal action of π .

First we introduce the cup product on twisted cohomology. The following lemma can be verified easily by hand, say along the lines of the proof of [Hat02, Lemma 3.6].

Lemma A.11. *Let Y be a subset of X . For all $p, q \in \mathbb{N}_0$ We consider the map*

$$\begin{aligned} \cup : C^p(X, Y; A) \times C^q(X, Y; B) &\longrightarrow C^{p+q}(X, Y; A \otimes_{\mathbb{Z}} B) \\ (\phi, \psi) &\longmapsto (\sigma \mapsto \varphi(\sigma]_p) \otimes_{\mathbb{Z}} \psi(\sigma]_{k-p}). \end{aligned}$$

(Note that the right-hand side is indeed a $\mathbb{Z}[\pi]$ -homomorphism, i.e. it defines an element $C^{p+q}(X; A \otimes_{\mathbb{Z}} B)$.) Furthermore the map descends to a well defined map

$$\cup : H^p(X, Y; A) \times H^q(X, Y; B) \longrightarrow H^{p+q}(X, Y; A \otimes_{\mathbb{Z}} B).$$

We refer to this map as the cup product.

Next we introduce the cap product. As with cup product, first we define it on the chain level.

Lemma A.12. *Let $S, T \subseteq X$ be subsets. We write $C_k(X, \{S, T\}) = C_k(X)/(C_k(S) + C_k(T))$. The map*

$$\begin{aligned} \cap : C^p(X, S; A) \times C_k(X, \{S, T\}; A) &\longrightarrow C_{k-p}(X, T; A \otimes_{\mathbb{Z}} B) \\ (\psi, b \otimes_{\mathbb{Z}[\pi]} \sigma) &\longmapsto (\psi(\sigma|_p) \otimes_{\mathbb{Z}} b) \otimes_{\mathbb{Z}[\pi]} \sigma|_{k-p}. \end{aligned}$$

is well-defined. We refer to this map as the cap product.

PROOF. We verify that the given map respects the tensor product. Thus let $\psi \in C^p(X; A)$, $\sigma \in C_k(\tilde{X})$, $\gamma \in \pi$ and $b \in B$. We calculate that

$$\begin{aligned} \psi \cap b \otimes_{\mathbb{Z}[\pi]} \gamma \sigma &= (\psi(\gamma \sigma|_p) \otimes_{\mathbb{Z}} b) \otimes_{\mathbb{Z}[\pi]} \gamma \sigma|_{k-p} &= (\gamma \psi(\sigma|_p) \otimes_{\mathbb{Z}} b) \cdot \gamma \otimes_{\mathbb{Z}[\pi]} \sigma|_{k-p} \\ &= (\gamma^{-1} \gamma \psi(\sigma|_p) \otimes_{\mathbb{Z}} b \gamma) \otimes_{\mathbb{Z}[\pi]} \sigma|_{k-p} &= \psi \cap b \gamma \otimes_{\mathbb{Z}[\pi]} \sigma. \end{aligned}$$

It follows easily from the definitions that the cap product descends to the given quotient (co-) chain complexes. \square

Lemma A.13. *Let $f \in C^p(X; A)$ and let $c \in C_k(X; B)$. We have*

$$\partial(f \cap c) = (-1)^p \cdot (-\delta(f) \cap c + f \cap \partial c) \in C_{k-1}(X; A \otimes_{\mathbb{Z}} B).$$

PROOF. The lemma follows from a calculation using the definition of the cap product and the boundary maps, see e.g. [Fri23, Lemma 192.9] for details. Note that the precise signs differ from similar formulas in some textbooks in algebraic topology since there are many different sign conventions in usage. \square

Corollary A.14. *Let $S, T \subseteq X$ be subsets, let R be a ring and let A be an $(R, \mathbb{Z}[\pi_1(M)])$ -bimodule. For any cycle $\sigma \in C_n(X, \{S, T\}; \mathbb{Z})$ the cap product*

$$\begin{aligned} \cap[\sigma] : H^k(X, S; A) &\rightarrow H_{n-k}(X, T; A) = H_{n-k}(X, T; A \otimes_{\mathbb{Z}} \mathbb{Z}) \\ [\varphi] &\mapsto [\varphi \cap \sigma] \end{aligned}$$

is well-defined. Furthermore this map only depends on the homology class $[\sigma] \in H_n(X, S \cup T; \mathbb{Z})$.

A.2. THE POINCARÉ DUALITY THEOREM

The following theorem is a generalisation of the familiar Poincaré duality for untwisted coefficients to the case of twisted coefficients.

Theorem A.15. (Twisted Poincaré Duality Theorem) *Let M be a compact, oriented, connected n -dimensional manifold. Let S and T be codimension 0 compact submanifolds of ∂M such that $\partial S = \partial T = S \cap T$ and $\partial M = S \cup T$. Let $[M] \in H_n(M, \partial M; \mathbb{Z})$ be the fundamental class of M . If R is a ring and if A is an $(R, \mathbb{Z}[\pi_1(M)])$ -bimodule, then the map*

$$- \cap [M] : H^k(M, S; A) \rightarrow H_{n-k}(M, T; A)$$

defined by Lemma A.13 is an isomorphism of left R -modules.

We also have the following Poincaré Duality statement on the (co-) chain level.

Theorem A.16. (Universal Poincaré Duality Theorem) *Let M a compact, oriented, connected n -dimensional manifold. Let S and T be codimension 0 compact submanifolds of ∂M such that $\partial S = \partial T = S \cap T$ and $\partial M = S \cup T$. Let $\sigma \in C_n(M, \{S, T\}; \mathbb{Z})$ be a representative of the fundamental class of M . If R is a ring and if A is an $(R, \mathbb{Z}[\pi_1(M)])$ -bimodule, then the map*

$$- \cap \sigma: C^k(M, S; \mathbb{Z}[\pi_1(M)]) \rightarrow C_{n-k}(M, T; \mathbb{Z}[\pi_1(M)])$$

defined by Lemma A.13 is a chain homotopy equivalence of left R -chain complexes.

Note that Theorem 14.9 can be used to give an alternative proof that the chain complexes of the theorem are chain homotopy equivalent.

PROOF OF THEOREM A.16 USING THEOREM A.15. The Universal Poincaré Duality Theorem A.16 follows immediately from the Twisted Poincaré Duality Theorem A.15 together with Lemma A.4. \square

In the following sections we will provide a proof of the Twisted Poincaré Duality Theorem A.15. But just for fun we would like to show that the Universal Poincaré Duality Theorem A.16 also implies the Twisted Poincaré Duality Theorem A.15, in other words, the two theorems are equivalent:

PROOF OF THEOREM A.15 USING THEOREM A.16. Let M be a compact, oriented, connected n -dimensional manifold. To simplify the discussion we just deal with the case that $S = \partial M$ and $T = \emptyset$. We pick a representative σ for $[M]$ and we write $\pi = \pi_1(M)$. Let A be an $(R, \mathbb{Z}[\pi])$ -bimodule. Given a chain complex D_* of right $\mathbb{Z}[\pi]$ -modules we consider the cochain map

$$\begin{aligned} \Xi: A \otimes_{\mathbb{Z}[\pi]} \text{Hom}_{\text{right-}\mathbb{Z}[\pi]}(D_*; \mathbb{Z}[\pi]) &\rightarrow \text{Hom}_{\text{right-}\mathbb{Z}[\pi]}(D_*, A) \\ a \otimes f &\mapsto (\sigma \mapsto a \cdot f(\sigma)). \end{aligned}$$

Note that Ξ is an isomorphism if each D_k is a finitely generated free $\mathbb{Z}[\pi]$ -module. But in general Ξ is not an isomorphism.

Furthermore we consider the following diagram

$$\begin{array}{ccc} H^k(A \otimes_{\mathbb{Z}[\pi]} C^*(M, \partial M; \mathbb{Z}[\pi])) & \xrightarrow{\text{Id}_A \otimes (\cap \sigma)} & H_{n-k}(A \otimes_{\mathbb{Z}[\pi]} C_*(M; \mathbb{Z}[\pi])) = H_{n-k}(M; A) \\ \downarrow \Xi_* & \nearrow \cap [M] & \\ H^k(C^*(M, \partial M; A)) = H^k(M, \partial M; A) & & \end{array}$$

One easily verifies that the diagram commutes. The top horizontal map is an isomorphism by the Universal Poincaré Duality Theorem A.16. It remains to show that the vertical map is an isomorphism. As we had pointed out above, on the chain level Ξ is in general not an isomorphism.

As in the proof of Proposition A.5 we can use Theorem 3.16 to find a pair (X, Y) of finite CW complexes and a homotopy equivalence $f: (X, Y) \rightarrow (M, \partial M)$. By Proposition A.3 there exists a homotopy equivalence $\Theta: C_*^{\text{cell}}(X, Y; \mathbb{Z}[\pi]) \rightarrow C_*(X, Y; \mathbb{Z}[\pi])$ of

$\mathbb{Z}[\pi]$ -chain complexes. We consider the following diagram where all tensor products and homomorphism are over $\mathbb{Z}[\pi]$:

$$\begin{array}{ccccc} A \otimes C_{\text{cell}}^*(X, Y; \mathbb{Z}[\pi]) & \xrightarrow{\Theta^*} & A \otimes C^*(X, Y; \mathbb{Z}[\pi]) & \xrightarrow{f^*} & A \otimes C^*(M, \partial M; \mathbb{Z}[\pi]) \\ \downarrow \Xi_* & & \downarrow \Xi_* & & \downarrow \Xi_* \\ C_{\text{cell}}^*(X, Y; A) & \xrightarrow{\Theta^*} & C^*(X, Y; A) & \xrightarrow{f^*} & C^*(M, \partial M; A). \end{array}$$

One easily verifies that the diagram commutes. As pointed out above, the horizontal maps are chain homotopy equivalences over $\mathbb{Z}[\pi]$. Since X is a finite CW complex we see that each $C_k^{\text{cell}}(X, Y; \mathbb{Z}[\pi])$ is a finitely generated free $\mathbb{Z}[\pi]$ -module. Thus we obtain from the above that the left vertical map is an isomorphism. Therefore the right vertical map is a chain homotopy equivalence. In particular it induces an isomorphism of homology groups. \square

The remainder of this appendix is dedicated to the proof of the Twisted Poincaré Duality Theorem A.15. Even though the theorem is well-known and often used, there are not many satisfactory proofs in the literature. The proof which is closest to ours in spirit is the proof of Sun [Sun17]. For closed manifolds Kwasik-Sun [KS18] provide a proof by using the work of Kirby-Siebenmann to reduce the proof to the case of triangulated manifolds.

The proof of the Twisted Poincaré Duality Theorem A.15 is modelled on the proof of untwisted Poincaré Duality that is given in Bredon's book [Bre97, Chapter VI.8]. The logic of his proof is unchanged, but some arguments and definitions have to be adjusted for the twisted setting.

A.3. PREPARATIONS FOR THE PROOF OF THE TWISTED POINCARÉ DUALITY THEOREM

We fix some notation that we will use for the remainder of the appendix. Let M be a *connected* manifold, let $x_0 \in M$ and denote by $\pi := \pi_1(M, x_0)$ the fundamental group. Finally let R be a ring and let A be an $(R, \mathbb{Z}[\pi])$ -bimodule.

We write $p: \widetilde{M} \rightarrow M$ for the universal cover of M . For a subset $X \subseteq M$ (not necessarily connected) we consider the (co)-homology of X with respect to the coefficient system coming from M by setting

$$\begin{aligned} C_*(X; A) &:= A \otimes_{\mathbb{Z}[\pi]} C_*(p^{-1}(X); \mathbb{Z}), \\ C^*(X; A) &:= \text{Hom}_{\mathbb{Z}[\pi]} \left(\overline{C_*(p^{-1}(X); \mathbb{Z})}, A \right), \end{aligned}$$

with generalisation to pairs $Y \subseteq X \subseteq M$ by

$$\begin{aligned} C_*(X, Y; A) &:= A \otimes_{\mathbb{Z}[\pi]} C_*(p^{-1}(X), p^{-1}(Y); \mathbb{Z}), \\ C^*(X, Y; A) &:= \text{Hom}_{\mathbb{Z}[\pi]} \left(\overline{C_*(p^{-1}(X), p^{-1}(Y); \mathbb{Z})}, A \right). \end{aligned}$$

We summarise the basic properties of (co-) homology with twisted coefficients in the following theorem, which should be compared to the untwisted case.

Theorem A.17. *Let M be a connected manifold with fundamental group π , and let A be an $(R, \mathbb{Z}[\pi])$ -bimodule.*

(1) *Given $Y \subseteq X \subseteq M$ there is a long exact sequence of pairs in homology*

$$\cdots \rightarrow H_k(Y; A) \rightarrow H_k(X; A) \rightarrow H_k(X, Y; A) \rightarrow H_{k-1}(Y; A) \rightarrow \cdots$$

and cohomology

$$\cdots \rightarrow H^k(X, Y; A) \rightarrow H^k(X; A) \rightarrow H^k(Y; A) \rightarrow H^{k+1}(X, Y; A) \rightarrow \cdots .$$

(2) *Suppose we have a chain of subspaces $Z \subseteq Y \subseteq X \subseteq M$ such that the closure of Z is contained in the interior of Y . Then the inclusion $(X \setminus Z, Y \setminus Z) \rightarrow (X, Y)$ induces an isomorphism in homology and cohomology i.e. for all $k \in \mathbb{N}_0$ we have*

$$H_k(X \setminus Z, Y \setminus Z; A) \xrightarrow{\cong} H_k(X, Y; A) \quad \text{and} \quad H^k(X \setminus Z, Y \setminus Z; A) \xleftarrow{\cong} H^k(X, Y; A)$$

(3) *If $U_1 \subseteq U_2 \subseteq M$ and $V_1 \subseteq V_2 \subseteq M$ are open subsets in M , then there are long exact sequences in homology*

$$\begin{aligned} \cdots \rightarrow H_k(U_1 \cap V_1, U_2 \cap V_2; A) &\rightarrow \begin{array}{c} H_k(U_1, U_2; A) \\ \oplus \\ H_k(V_1, V_2; A) \end{array} \rightarrow H_k(U_1 \cup V_1, U_2 \cup V_2; A) \\ &\rightarrow H_{k-1}(U_1 \cap V_1, U_2 \cap V_2; A) \longrightarrow \cdots \end{aligned}$$

and cohomology

$$\begin{aligned} \cdots &\longrightarrow H^{k-1}(U_1 \cap V_1, U_2 \cap V_2; A) \\ &\rightarrow H^k(U_1 \cup V_1, U_2 \cup V_2; A) \rightarrow \begin{array}{c} H^k(U_1, U_2; A) \\ \oplus \\ H^k(V_1, V_2; A) \end{array} \rightarrow H^k(U_1 \cap V_1, U_2 \cap V_2; A) \rightarrow \cdots \end{aligned}$$

(4) *Suppose the inclusion $Y \rightarrow X$ is a homotopy equivalence, then the inclusion induced maps*

$$H_k(Y; A) \xrightarrow{\cong} H_k(X; A) \quad \text{and} \quad H^k(Y; A) \xleftarrow{\cong} H^k(X; A)$$

are isomorphisms.

(5) *Let $U_1 \subseteq U_2 \subseteq \dots$ be a sequence of open sets in M and let $U = \bigcup_{i \in \mathbb{N}} U_i$, then for each $k \in \mathbb{N}_0$ the inclusions induce an isomorphism*

$$\varinjlim_{i \in \mathbb{N}} H_k(U_i; A) \xrightarrow{\cong} H_k(U; A).$$

The proofs are essentially the same as in the classical case. Therefore we will only sketch the arguments and focus on what is different. We also warn the reader that we give the “philosophically wrong proof” of statement (4). This is due to the fact that we developed the theory of twisted coefficients only for inclusions and hence a homotopy inverse does not

fit in our theory. Therefore statement (4) will be deduced in a slightly round-about way using the following elementary lemma [Bre97, Theorem III.3.4 & remark after proof].

Lemma A.18. (Covering Homotopy Theorem) *Given a covering $p: \tilde{X} \rightarrow X$, a homotopy $H: Y \times I \rightarrow X$, and a lift $\tilde{h}: Y \rightarrow \tilde{X}$ of $H(-, 0)$, there exists a unique lift $\tilde{H}: Y \times I \rightarrow \tilde{X}$ of H with $\tilde{h} = \tilde{H}(-, 0)$.*

PROOF OF THEOREM A.17. Recall that $p: \tilde{M} \rightarrow M$ denotes the universal cover. For statement (1) we consider the short exact sequence $0 \rightarrow C_*(Y; \mathbb{Z}[\pi]) \rightarrow C_*(X; \mathbb{Z}[\pi]) \rightarrow C_*(X, Y; \mathbb{Z}[\pi]) \rightarrow 0$ of free $\mathbb{Z}[\pi]$ -modules. Since the modules are free the sequence stays exact after applying the functors $A \otimes_{\mathbb{Z}[\pi]} -$ and $\text{Hom}_{\mathbb{Z}[\pi]}(-, A)$.

Recall the proof of statement (2) and (3) in the classical case as in [Bre97, Chapter IV.17]. The main ingredient is to show that the inclusion of chain complexes $C_*^{\mathcal{U}}(X; \mathbb{Z}[\pi]) \rightarrow C^*(X; \mathbb{Z}[\pi])$ induces an isomorphism on homology [Bre97, Theorem IV.17.7]. Here \mathcal{U} is an open cover of X and $C_*^{\mathcal{U}}(X; \mathbb{Z}[\pi])$ is the free abelian group generated by simplices σ for which there is a $U \in \mathcal{U}$ such that $\sigma: \Delta^* \rightarrow p^{-1}(U)$. This is done by defining the barycentric subdivision $\Upsilon_*: C_*(\tilde{X}; \mathbb{Z}) \rightarrow C_*(\tilde{X}; \mathbb{Z})$ and a chain homotopy T between Υ_* and the identity [Bre97, Lemma IV.17.1]. The important thing for us to observe is that both maps are natural [Bre97, Claim (1) in proof of Lemma IV.17.1]. Hence for a twisted chain $\Upsilon(e \otimes_{\mathbb{Z}[\pi]} \sigma) := e \otimes_{\mathbb{Z}[\pi]} \Upsilon(\sigma)$ is well-defined, because

$$\begin{aligned} \Upsilon(e \otimes_{\mathbb{Z}[\pi]} \gamma\sigma) &= e \otimes_{\mathbb{Z}[\pi]} \Upsilon(\gamma\sigma) \\ &= e \otimes_{\mathbb{Z}[\pi]} \gamma\Upsilon(\sigma) \quad (\text{naturality of } \Upsilon) \\ &= e\gamma \otimes_{\mathbb{Z}[\pi]} \Upsilon(\sigma) = \Upsilon(e\gamma \otimes_{\mathbb{Z}[\pi]} \sigma). \end{aligned}$$

The same holds for T and from now on one can follow the classical proofs. Alternatively, one could invoke Lemma A.4.

Next we prove statement (4). Let $f: X \rightarrow Y \subseteq X$ be a homotopy inverse of the inclusion and $H: X \times I \rightarrow X$ a homotopy between Id_X and f . Since $p: \tilde{X} \rightarrow X$ is a covering and $\text{Id}_{\tilde{X}}$ is a lift of $H(p(-), 0)$, we get by Lemma A.3 a lift $\tilde{H}: \tilde{X} \times I \rightarrow \tilde{X}$ of the homotopy H . One easily verifies that the inclusion $\tilde{Y} \rightarrow \tilde{X}$ induces a homotopy equivalence where a homotopy inverse is given by $\tilde{H}(-, 1)$. Hence the inclusion induced map $H_k(C_*(\tilde{Y}; \mathbb{Z})) \rightarrow H_k(C_*(\tilde{X}; \mathbb{Z}))$ is an isomorphism for every k . Thus the claim follows from Lemma A.4.

The proof of Statement (5) is almost verbatim the same proof as in the classical case. \square

A.4. THE MAIN TECHNICAL THEOREM

Given a group π we can view \mathbb{Z} as a $\mathbb{Z}[\pi]$ -module with trivial π -action. We denote this module by \mathbb{Z}^{triv} . Let $p: \tilde{M} \rightarrow M$ be the covering projection. We have the following useful lemma, concerning the chain map $C_*(X; \mathbb{Z}^{\text{triv}}) \rightarrow C_*(X; \mathbb{Z})$ defined by $k \otimes_{\mathbb{Z}[\pi]} \tilde{\sigma} \mapsto k \cdot p(\sigma)$.

Lemma A.19. *Given any subset $X \subseteq M$ the chain map above is an isomorphism between $C_*(X; \mathbb{Z}^{\text{triv}})$ and $C_*(X; \mathbb{Z})$, and induces one between $C^*(X; \mathbb{Z})$ and $C^*(X; \mathbb{Z}^{\text{triv}})$, where $C_*(X; \mathbb{Z})$ and $C^*(X; \mathbb{Z})$ are the untwisted singular chain complexes.*

PROOF. The isomorphism is given by lifting a simplex, which is always possible since a simplex is simply connected. If one has two different choices of lifts, then they differ by an element in π . But the action of $\mathbb{Z}[\pi]$ on \mathbb{Z} is trivial and hence this indeterminacy vanishes. \square

We will keep the notational difference between $C_*(X; \mathbb{Z})$ and $C_*(X; \mathbb{Z}^{\text{triv}})$ to emphasise where our simplices live.

As above let R be a ring and let A be an $(R, \mathbb{Z}[\pi])$ -bimodule. Let $K \subseteq M$ be a compact subset of M . We define the (twisted) Čech cohomology groups

$$\check{H}^p(K; A) := \lim_{\substack{\longrightarrow \\ K \subseteq U \subseteq M}} H^p(U; A),$$

where the direct limit runs over all open sets in M containing K . Since cohomology is contravariant, we define the order on open sets in the reversed way i.e. $U \leq V$ if $V \subseteq U$.

Now we assume that M is oriented. Being oriented gives us, for any compact subset $K \subseteq M$, a preferred element $\theta_K \in H_n(M, M \setminus K; \mathbb{Z}^{\text{triv}}) \cong H_n(M, M \setminus K; \mathbb{Z})$, which restricts for all $x \in K$ to the generator in $H_n(M, M \setminus \{x\}; \mathbb{Z}^{\text{triv}})$.

For any open set $U \subseteq M$ containing K , the inclusion induced map $H_n(U, U \setminus K; \mathbb{Z}^{\text{triv}}) \rightarrow H_n(M, M \setminus K; \mathbb{Z}^{\text{triv}})$ is an isomorphism by Theorem A.17 (2). Let $\text{ex}_U: H_n(M, M \setminus K; \mathbb{Z}^{\text{triv}}) \rightarrow H_n(U, U \setminus K; \mathbb{Z}^{\text{triv}})$ be the inverse of this inclusion induced isomorphism i.e. if $j: U \rightarrow M$ is the inclusion then $j_* \circ \text{ex}_U = \text{Id}$. We then obtain a map

$$\begin{aligned} D_U: H^p(U; A) &\longrightarrow H_n(M, M \setminus K; A) \\ \phi &\longmapsto j_*(\phi \cap \text{ex}_U(\theta_K)). \end{aligned}$$

Given another open set $V \subseteq U$ denote by $i: V \rightarrow U$ the inclusion. Then one easily calculates:

$$\text{PD}_V(i^* \phi) = j_* i_*(i^* \phi \cap \text{ex}_V(\theta_K)) = j_*(\phi \cap i_* \text{ex}_V(\theta_K)) = j_*(\phi \cap \text{ex}_U(\theta_K)) = \text{PD}_U(\phi).$$

Or, in other words, the following diagram commutes:

$$\begin{array}{ccc} H^p(U; A) & \xrightarrow{\text{PD}_U} & H_{n-p}(M, M \setminus K; A). \\ \downarrow i^* & \nearrow \text{PD}_V & \\ H^p(V; A) & & \end{array}$$

By the universal property of the direct limit we obtain the *dualising* map $\text{PD}_K: \check{H}^p(K; A) \rightarrow H_{n-p}(M, M \setminus K; A)$.

In the remainder of this section we will prove the following theorem.

Theorem A.20. (Poincaré Duality Theorem) *The map $\text{PD}_K: \check{H}^p(K; A) \rightarrow H_{n-p}(M, M \setminus K; A)$ is a left R -module isomorphism for all compact subsets $K \subseteq M$.*

Here, as above, A is an $(R, \mathbb{Z}[\pi])$ -bimodule. In the subsequent section we will see that the Twisted Poincaré Duality Theorem A.15 is a reasonably straightforward consequence of Theorem A.20.

The proof of Theorem A.20 will be an application of the following lemma.

Lemma A.21. (Bootstrap lemma) *For each compact subspace $K \in M$ let $P_M(K)$ be a statement. If $P_M(\cdot)$ satisfies the following three conditions:*

- (1) $P_M(K)$ holds true for all compact subsets $K \subseteq M$ with the property that for all $x \in K$ the inclusions $\{x\} \rightarrow K$ and $M \setminus K \rightarrow M \setminus \{x\}$ are deformation retracts,
- (2) If $P_M(K_1), P_M(K_2)$ and $P_M(K_1 \cap K_2)$ are true, then $P_M(K_1 \cup K_2)$ is true,
- (3) If $\dots \subseteq K_2 \subseteq K_1$ and $P_M(K_i)$ is true for all $i \in \mathbb{N}$, then $P_M(\bigcap_{i \in \mathbb{N}} K_i)$ is true.

Then $P_M(K)$ is true for all $K \subseteq M$.

PROOF. See [Bre97, Lemma VI.7.9]. □

The idea is to apply the bootstrap lemma to the statement that the conclusion of Theorem A.20 holds for a given compact set K . It turns out that condition (3) is the easiest to verify. It follows from formal properties about direct limits. For the verification of condition (1) we have to do one explicit calculation. This is the content of the next lemma.

Lemma A.22. *Let $x \in M$ be a point. The map $\text{PD}_{\{x\}}: \check{H}^0(\{x\}; A) \rightarrow H_n(M, M \setminus \{x\}; A)$ is an R -module isomorphism.*

PROOF. Let $p: \widetilde{M} \rightarrow M$ be the universal cover. Since x is a point in a manifold we can calculate the dualising map $\text{PD}_{\{x\}}$ by taking the limit over open neighbourhoods U of x with the following two properties:

- (1) U is contractible,
- (2) for any connected component $\bar{U} \subseteq p^{-1}(U)$ the map $p|_{\bar{U}}$ is a homeomorphism.

This can be done, since any neighbourhood of x contains a neighbourhood with these two properties. Let U be such a neighbourhood of x and $\bar{U} \subseteq p^{-1}(U)$ a fixed connected component. This choice of connected component gives us an isomorphism $H^0(U; A) \cong A$ as follows. Let $f \in H^0(U; A)$ be arbitrary and $\bar{x} \in \bar{U}$ be a point in our connected component. Then we get an element in A by evaluating $f([\bar{x}])$. Conversely, given an element $e \in A$ we can construct a function in $H^0(U; A)$ by setting $f([\bar{x}]) = e$ for all $\bar{x} \in \bar{U}$. Note that there is a unique way to extend f equivariantly to $C_0(p^{-1}(U); \mathbb{Z})$.

We are now going to construct a representative of the orientation class $\theta_K \in H_n(M, M \setminus \{x\}; \mathbb{Z}^{\text{triv}})$ for which it is very simple to calculate the dualising map. Let \bar{x} be the preimage of x in \bar{U} . Now take a cycle $\sum_{i=1}^d k_i \sigma_i$ which generates $H_n(\bar{U}, \bar{U} \setminus \{\bar{x}\}; \mathbb{Z})$. By Theorem A.17 (2) and Lemma A.19 one easily sees that $1 \otimes_{\mathbb{Z}[\pi]} \sum_{i=1}^d k_i \sigma_i$ is a generator of $H_n(M, M \setminus \{x\}; \mathbb{Z}^{\text{triv}})$.

Using the isomorphism $H^0(U; A) \cong A$ from above the dualising map becomes $\text{PD}_{\{x\}}: A \rightarrow H_n(M, M \setminus \{x\}; A)$, $e \mapsto e \otimes_{\mathbb{Z}[\pi]} \sum_{i=1}^d k_i \sigma_i$. This is clearly an isomorphism, since on the chain level we have:

$$C_*(U, U \setminus \{x\}; A) = A \otimes_{\mathbb{Z}[\pi]} \bigoplus_{\gamma \in \pi} C_*(\gamma \bar{U}, \gamma \bar{U} \setminus \{\gamma \bar{x}\}; \mathbb{Z}) \cong A \otimes_{\mathbb{Z}} C_*(\bar{U}, \bar{U} \setminus \{\bar{x}\}; \mathbb{Z}). \quad \square$$

In order to verify condition (2) of the bootstrap lemma we will need the following lemma (compare [Bre97, Lemma VI.8.2]).

Lemma A.23. *If K and L are two compact subsets of M , then for all $p \in \mathbb{N}_0$ the diagram*

$$\begin{array}{ccccccc} \cdots & \longrightarrow & \check{H}^p(K \cup L; A) & \longrightarrow & \check{H}^p(K; A) \oplus \check{H}^p(L; A) & \longrightarrow & \check{H}^p(K \cap L; A) \longrightarrow \check{H}^{p+1}(K \cup L; A) \longrightarrow \cdots \\ & & \downarrow \text{PD}_{K \cup L} & & \downarrow \text{PD}_K \oplus \text{PD}_L & & \downarrow \text{PD}_{K \cap L} & & \downarrow \text{PD}_{K \cup L} \\ \cdots & \rightarrow & H_{n-p}(M, M \setminus (K \cup L); A) & \rightarrow & H_{n-p}(M, M \setminus K; A) \oplus H_{n-p}(M, M \setminus L; A) & \rightarrow & H_{n-p}(M, M \setminus (K \cap L); A) \rightarrow H_{n-p-1}(M, M \setminus (K \cup L); A) \rightarrow \cdots \end{array}$$

has exact rows and it commutes up to a sign depending only on p .

PROOF. The rows are exact by Mayer-Vietoris (see Theorem A.17 (3)) and the fact that direct limit is an exact functor. The commutativity of the squares is clear except for the last one involving the boundary map. This will be a painful diagram chase. Let $U \supseteq K$ and $V \supseteq L$ be open neighbourhoods containing K resp. L . The sequence in the top row comes from the short exact sequence ($\mathcal{U} = \{U, V\}$):

$$0 \rightarrow C_{\mathcal{U}}^*(U \cup V; A) \rightarrow C^*(U; A) \oplus C^*(V; A) \rightarrow C^*(U \cap V; A) \rightarrow 0.$$

An element $\phi \in \check{H}^p(K \cap L; A)$ will already be represented by some element $f \in C^p(U \cap V; A)$ for some U and V as above. We can extend f to an element $\bar{f} \in C^p(M; A)$ by

$$\bar{f}(\sigma) = \begin{cases} f(\sigma) & \text{if } \text{Im } \sigma \subseteq \tilde{U} \cap \tilde{V} \\ 0 & \text{else.} \end{cases}$$

Note that $\bar{f} \in C^p(M; A)$ since $p^{-1}(U \cap V)$ is an equivariant subspace and hence \bar{f} is equivariant. If we consider \bar{f} as an element in $C^p(U; A)$ then the cohomology class $\delta(\phi)$ is represented by the cochain $h \in C^{p+1}(U \cup V; A)$ which is given by

$$h(\sigma) = \begin{cases} \delta(\bar{f})(\sigma) & \text{if } \text{Im } \sigma \subseteq \tilde{U} \\ 0 & \text{else.} \end{cases}$$

Since ϕ is a cocycle we have $\delta(f)(\sigma) = 0$ for $\sigma \in C_*(U \cap V; A)$. It follows in particular that if σ is a simplex whose image is completely contained in \tilde{V} , then $h(\sigma) = 0$. We can

represent our orientation class $\theta \in H_n(M, M \setminus (K \cup L))$ by a cycle

$$a = b + c + d + e \quad \text{with} \quad b \in C_n(U \cap V; \mathbb{Z}^{\text{triv}}) \quad c \in C_n(U \setminus (U \cap L); \mathbb{Z}^{\text{triv}}) \\ d \in C_n(V \setminus (V \cap K); \mathbb{Z}^{\text{triv}}), \quad e \in C_n(M \setminus (K \cup L); \mathbb{Z}^{\text{triv}}).$$

Obviously e does not play a role since we kill it in the end. With these representatives one computes that $\delta(\phi)(\theta)$ is represented by

$$h \cap (b + c + d) = \delta(\bar{f}) \cap c + h \cap d + \delta(f) \cap b = \delta(\bar{f}) \cap c.$$

The pairing of h with b is zero since f was a cocycle in $C^*(U \cap V; A)$ and the pairing of h with d is zero since d consists of simplices with image in \tilde{V} .

The lower sequence comes from the short exact sequence:

$$0 \rightarrow C_*(M, M \setminus (K \cup L); A) \rightarrow \begin{array}{c} C_*(M, M \setminus K; A) \\ \oplus \\ C_*(M, M \setminus L; A) \end{array} \rightarrow C_*(M, M \setminus (K \cap L); A) \rightarrow 0.$$

Before we compute the other side $\partial(\phi \cap \text{ex}_{U \cap V}(\theta))$ we want to recall that the cap product is natural on the chain complex level i.e. the following diagram commutes:

$$\begin{array}{ccc} C^p(U; A) \times C_n(U, U \setminus K; \mathbb{Z}^{\text{triv}}) & \longrightarrow & C_*(U, U \setminus K; A) \\ \uparrow & & \downarrow \\ C^p(M; A) \times C_n(M, M \setminus K; \mathbb{Z}^{\text{triv}}) & \longrightarrow & C_*(M, M \setminus K; A). \end{array}$$

Therefore we use the representatives from above. To construct the boundary map ∂ , we take as the preimage of $\bar{f} \cap a \in C_*(M, M \setminus (K \cap L); A)$ the element $(\bar{f} \cap a, 0) \in C_*(M, M \setminus K; A) \oplus C_*(M, M \setminus L; A)$. Then one computes in $C_*(M, M \setminus K; A)$

$$\begin{aligned} \partial(\bar{f} \cap a) &= (-1)^{p+1} \cdot \delta(\bar{f}) \cap a \pm f \cap \partial a \quad (\text{by Lemma A.13}) \\ &= (-1)^{p+1} \cdot \delta(\bar{f}) \cap a \quad (\text{since } f \cap \partial a \in C_{n-p-1}(M \setminus (K \cup L); A)) \\ &= (-1)^{p+1} \cdot \delta(\bar{f}) \cap b + c + d + e \\ &= (-1)^{p+1} \cdot \delta(\bar{f}) \cap (c + d) \quad (\text{same reason as above}) \\ &= (-1)^{p+1} \cdot \delta(\bar{f}) \cap c \quad (\text{since } d \in C_{n-p}(V \setminus (K \cap V); A)) \end{aligned}$$

Therefore the element $\partial(\phi \cap \text{ex}_{U \cap V}(\theta))$ is also represented by $(-1)^{p+1} \cdot \delta(\bar{f}) \cap c \in C_{n-p-1}(M, M \setminus (K \cup L); A)$. \square

PROOF OF THEOREM A.20. Let $P_M(K)$ be the statement that the map PD_K is an isomorphism. Then it is sufficient to verify condition (1), (2) and (3) of the bootstrap lemma. We start by verifying (1). In the case that $K = \{x\}$ is just a point we have already

seen in Lemma A.22 that the statement holds true. For a general compact K with the property of (1) the statement follows from the following commutative diagram:

$$\begin{array}{ccc} \check{H}^p(K; A) & \longrightarrow & H_{n-p}(M, M \setminus K; A) \\ \downarrow \simeq & & \downarrow \simeq \\ \check{H}^p(\{x\}; A) & \xrightarrow{\simeq} & H_{n-p}(M, M \setminus \{x\}; A), \end{array}$$

where the vertical maps are isomorphisms by the homotopy invariance and the bottom row by the observation above. Hence condition (1) is verified.

Condition (2) follows immediately from the five lemma and Lemma A.23.

Let K_i be a sequence of compact subsets such that $P_M(K_i)$ holds for all $i \in \mathbb{N}$. We set $K = \bigcap_{i \in \mathbb{N}} K_i$. It is an exercise in point set topology of manifolds that each K_i has a fundamental system $U_{i,j}$ of open neighbourhoods. Fundamental system means that $U_{i,j} \subseteq U_{i,k}$ if $j < k$ and that for each open set U containing K_i there is a j such that $U_{i,j} \subseteq U$. Another exercise in the point set topology of manifolds shows that one can construct these sets such that $U_{1,j} \supseteq U_{2,j} \supseteq U_{3,j} \supseteq \dots$ for all $j \in \mathbb{N}$. Then $U_{i,j}$ is a fundamental system of open neighbourhoods of K with the order $(i,j) \leq (k,l) \Leftrightarrow i \leq k \wedge j \leq l$. One has the natural isomorphism [Bre97, Appendix D5]:

$$\varinjlim_{i \in \mathbb{N}} \check{H}^p(K_i; A) = \varinjlim_{i \in \mathbb{N}} \varinjlim_{j \in \mathbb{N}} H^p(U_{i,j}; A) \xrightarrow{\simeq} \varinjlim_{i,j \in \mathbb{N}} H^p(U_{i,j}; A) \cong \check{H}^p(K; A).$$

Hence the theorem follows from the commutativity of the diagram:

$$\begin{array}{ccc} \varinjlim_{i \in \mathbb{N}} \check{H}^p(K_i; A) & \longrightarrow & \varinjlim_{i \in \mathbb{N}} H_{n-p}(M, M \setminus K_i; A) \\ \downarrow & & \downarrow \\ \check{H}^p(K; A) & \longrightarrow & H_{n-p}(M, M \setminus K; A). \end{array}$$

□

A.5. PROOF OF THE TWISTED POINCARÉ DUALITY THEOREM

For the reader's convenience we recall the main theorem from the last section. Here, as above, R is a ring and A is an $(R, \mathbb{Z}[\pi])$ -bimodule.

Theorem A.20. *Let M be a compact, oriented, connected n -dimensional manifold. The map $\text{PD}_K: \check{H}^p(K; A) \rightarrow H_{n-p}(M, M \setminus K; A)$ is an isomorphism of left R -modules for all compact subsets $K \subseteq M$ and all $p \in \mathbb{N}_0$.*

Furthermore, we also recall that we need to prove the following theorem.

Theorem A.15. *Let M a compact, oriented, connected n -dimensional manifold. Let S and T be codimension 0 compact submanifolds of ∂M such that $\partial S = \partial T = S \cap T$ and $\partial M = S \cup T$. Let $[M] \in H_n(M, \partial M; \mathbb{Z})$ be the fundamental class of M . The map*

$$- \cap [M]: H^k(M, S; A) \rightarrow H_{n-k}(M, T; A)$$

defined by Lemma A.13 is an isomorphism of left R -modules.

In the remainder of this appendix we will explain how to deduce Theorem A.15 from Theorem A.20. First note that if M is a closed manifold, then we can set $K = M$ in Theorem A.20. Evidently we have $\check{H}^p(M; A) = H(M; A)$. Thus we obtain precisely the statement of Theorem A.15 in the closed case.

Next let M be a compact oriented manifold with nonempty boundary. First we consider the case $R = \emptyset$ and $S = \partial M$. By the Collar Neighbourhood Theorem 2.16 there exists a collar $\partial M \times [0, 2] \subseteq M$ of the boundary such that $\partial M = \partial M \times \{0\}$. We obtain the following chain of isomorphisms:

$$\begin{aligned} H^p(M; A) &\cong H^p(M \setminus (\partial M \times [0, 1]); A) \quad (\text{homotopy}) \\ &\cong \check{H}^p(M \setminus (\partial M \times [0, 1]); A) \quad (\text{follows from considering the open} \\ &\hspace{15em} \text{neighbourhoods } M \setminus (\partial M \times [0, 1 - \frac{1}{n}])) \\ &\cong H_{n-p}(M \setminus \partial M, \partial M \times (0, 1); A) \quad (\text{duality } K = M \setminus (\partial M \times [0, 1])) \\ &\cong H_{n-p}(M, \partial M \times [0, 1]; A) \quad (\text{excision } U = \partial M) \\ &\cong H_{n-p}(M, \partial M; A), \end{aligned}$$

It follows from the definition of the dualising map and naturality of cap product that these isomorphisms are given by capping with a generator $[M] \in H_n(M, \partial M; \mathbb{Z}^{\text{triv}}) \cong H_n(M, \partial M; \mathbb{Z})$ as in the classical case.

The proof of the general case of Theorem A.15 relies on the following lemma.

Lemma A.24. *Let M be a compact, oriented, connected n -dimensional manifold. Let R and S be compact codimension 0 submanifolds of ∂M such that $\partial R = \partial S = R \cap S$ and $\partial M = R \cup S$. For each $p \in \mathbb{N}_0$ the following diagram commutes up to a sign:*

$$\begin{array}{ccccccc} \dots \rightarrow & H^p(M, \partial M; A) & \rightarrow & H^p(M, R; A) & \longrightarrow & H^p(\partial M, R; A) & \xrightarrow{\delta} & H^{p+1}(M, \partial M; A) & \rightarrow & \dots \\ & \downarrow \cap [M] & & \downarrow \cap [M] & & \downarrow \text{Theorem A.17(2)} & & \downarrow \cap [M] & & \\ & & & & & H_{n-p-1}(S, \partial S; A) & & & & \\ & & & & & \downarrow \cap [R] & & & & \\ \dots \rightarrow & H_{n-p}(M; A) & \rightarrow & H_{n-p}(M, S; A) & \xrightarrow{\partial} & H_{n-p-1}(S; A) & \longrightarrow & H_{n-p-1}(M; A) & \rightarrow & \dots \end{array}$$

PROOF. The commutativity is a more or less direct consequence of Lemma A.13 and the observation that $\partial_*[M] = [\partial M]$. More precisely, the proof in the untwisted case is given in detail in [Fri23, p. 2892]. The proof in the twisted case is basically the same. \square

The proof of the general case of Theorem A.15 follows from the previously discussed Poincaré Duality isomorphisms $\cap[M]: H^p(M; A) \xrightarrow{\cong} H_{n-p}(M, \partial M; A)$, $\cap[R]: H^{p+1}(R, A) \xrightarrow{\cong} H_{n-p-1}(R, \partial R; A)$ together with Lemma A.24 and the five lemma.

Bibliography

- [AFW15] Matthias Aschenbrenner, Stefan Friedl, and Henry Wilton. Decision problems for 3-manifolds and their fundamental groups. In *Interactions between low dimensional topology and mapping class groups. Proceedings of the conference, Bonn, Germany, July 1–5, 2013*, pages 201–236. Coventry: Geometry & Topology Publications, 2015.
- [Ale24] James W. Alexander. An example of a simply connected surface bounding a region which is not simply connected. *Proceedings of the National Academy of Sciences*, 10(1):8–10, 1924.
- [AM90] Selman Akbulut and John D. McCarthy. *Casson’s invariant for oriented homology 3-spheres*, volume 36 of *Mathematical Notes*. Princeton University Press, Princeton, NJ, 1990.
- [APS75] Michael F. Atiyah, Vijay K. Patodi, and Isadore M. Singer. Spectral asymmetry and Riemannian geometry. II. *Math. Proc. Cambridge Philos. Soc.*, 78(3):405–432, 1975.
- [Arm70] Mark A. Armstrong. Collars and concordances of topological manifolds. *Comment. Math. Helv.*, 45:119–128, 1970.
- [Ati76] Michael F. Atiyah. Elliptic operators, discrete groups and von Neumann algebras. In *Colloque “Analyse et Topologie” en l’Honneur de Henri Cartan (Orsay, 1974)*, pages 43–72. Astérisque, No. 32–33. Soc. Math. France, Paris, 1976.
- [BCF⁺21] Imre Bokor, Diarmuid Crowley, Stefan Friedl, Fabian Hebestreit, Daniel Kasproski, Markus Land, and Johnny Nicholson. Connected sum decompositions of high-dimensional manifolds. In *2019–20 MATRIX annals*, pages 5–30. Cham: Springer, 2021.
- [BG64a] Morton Brown and Herman Gluck. Stable structures on manifolds. I. Homeomorphisms of S^n . *Ann. of Math. (2)*, 79:1–17, 1964.
- [BG64b] Morton Brown and Herman Gluck. Stable structures on manifolds. II. Stable manifolds. *Ann. of Math. (2)*, 79:18–44, 1964.
- [BG23] Ryan Budney and David Gabai. On the automorphism groups of hyperbolic manifolds. *arXiv e-prints*, 2023.
- [Bin59] R. H. Bing. An alternative proof that 3-manifolds can be triangulated. *Ann. of Math.*, (2) 69:37–65, 1959.
- [Bin83] R. H. Bing. *The geometric topology of 3-manifolds*, 1983. American Mathematical Society.
- [BKK⁺21] Stefan Behrens, Boldizsár Kalmár, Min Hoon Kim, Mark Powell, and Arunima Ray. *The Disc Embedding Theorem*. Oxford University Press, 2021.
- [Boy86] Steven Boyer. Simply-connected 4-manifolds with a given boundary. *Trans. Amer. Math. Soc.*, 298(1):331–357, 1986.
- [Boy93] Steven Boyer. Realization of simply-connected 4-manifolds with a given boundary. *Comment. Math. Helv.*, 68(1):20–47, 1993.
- [Bre97] Glen Bredon. *Topology and geometry*, volume 139 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1997. Corrected third printing of the 1993 original.
- [Bro60] Morton Brown. A proof of the generalized Schoenflies theorem. *Bull. Am. Math. Soc.*, 66:74–76, 1960.
- [Bro62] Morton Brown. Locally flat imbeddings of topological manifolds. *Ann. of Math. (2)*, 75:331–341, 1962.

- [Bro72] William Browder. *Surgery on simply-connected manifolds*. Springer-Verlag, New York-Heidelberg, 1972. Ergebnisse der Mathematik und ihrer Grenzgebiete, Band 65.
- [BV68] J. Michael Boardman and Rainer M. Vogt. Homotopy-everything H -spaces. *Bull. Amer. Math. Soc.*, 74:1117–1122, 1968.
- [BV73] J. Michael Boardman and Rainer M. Vogt. *Homotopy invariant algebraic structures on topological spaces*. Lecture Notes in Mathematics, Vol. 347. Springer-Verlag, Berlin-New York, 1973.
- [Cer68] Jean Cerf. *Sur les difféomorphismes de la sphère de dimension trois $\Gamma_4 = 0$* , volume 53 of *Lect. Notes Math.* Springer, Cham, 1968.
- [CF13] Jae Choon Cha and Stefan Friedl. Twisted torsion invariants and link concordance. *Forum Math.*, 25(3):471–504, 2013.
- [CG78] Andrew Casson and Cameron Gordon. On slice knots in dimension three. In *Algebraic and geometric topology (Proc. Sympos. Pure Math., Stanford Univ., Stanford, Calif., 1976), Part 2*, pages 39–53. Amer. Math. Soc., Providence, R.I., 1978.
- [CG86] Andrew Casson and Cameron Gordon. Cobordism of classical knots. In *À la recherche de la topologie perdue*, pages 181–199. Birkhäuser Boston, Boston, MA, 1986. With an appendix by P. M. Gilmer.
- [Cha74] Thomas A. Chapman. Topological invariance of Whitehead torsion. *Amer. J. Math.*, 96:488–497, 1974.
- [Con71] Robert Connelly. A new proof of brown’s collaring theorem. *Proceedings of the American Mathematical Society*, 27(1):180–182, 1971.
- [COT03] Tim D. Cochran, Kent E. Orr, and Peter Teichner. Knot concordance, Whitney towers and L^2 -signatures. *Ann. of Math. (2)*, 157(2):433–519, 2003.
- [CP23] Anthony Conway and Mark Powell. Embedded surfaces with infinite cyclic knot group. *Geom. Topol.*, 27(2):739–821, 2023.
- [CPP22] Anthony Conway, Lisa Piccirillo, and Mark Powell. 4-manifolds with boundary and fundamental group \mathbb{Z} . *arXiv e-prints*, 2022.
- [CS71] Sylvain E. Cappell and Julius L. Shaneson. On four dimensional surgery and applications. *Comment. Math. Helv.*, 46:500–528, 1971.
- [CS76] Sylvain E. Cappell and Julius L. Shaneson. Some new four-manifolds. *Ann. of Math. (2)*, 104(1):61–72, 1976.
- [CT07] Tim D. Cochran and Peter Teichner. Knot concordance and von Neumann ρ -invariants. *Duke Math. J.*, 137(2):337–379, 2007.
- [CZ90] Donald J. Collins and Heiner Zieschang. Combinatorial group theory and fundamental groups. In *Algebra VII. Combinatorial group theory. Applications to geometry. Transl. from the Russian*. Berlin: Springer-Verlag, 1990.
- [Ded62] P. Dedecker. Variétés linéaires par morceaux et variétés combinatoires. *Ann. Mat. Pura Appl. (4)*, 60:365–383, 1962.
- [Dol95] Albrecht Dold. *Lectures on algebraic topology*. Classics in Mathematics. Springer-Verlag, Berlin, 1995. Reprint of the 1972 edition.
- [Don83] Simon K. Donaldson. An application of gauge theory to four-dimensional topology. *J. Differential Geom.*, 18(2):279–315, 1983.
- [Don87] Simon K. Donaldson. The orientation of Yang-Mills moduli spaces and 4-manifold topology. *J. Differential Geom.*, 26(3):397–428, 1987.
- [Edw84] Robert D. Edwards. The solution of the 4-dimensional annulus conjecture (after Frank Quinn). In *Four-manifold theory (Durham, N.H., 1982)*, volume 35 of *Contemp. Math.*, pages 211–264. Amer. Math. Soc., Providence, RI, 1984.
- [EK71] Robert D. Edwards and Robion C. Kirby. Deformations of spaces of imbeddings. *Ann. Math. (2)*, 93:63–88, 1971.

- [ERW19] Johannes Ebert and Oscar Randal-Williams. Semisimplicial spaces. *Algebr. Geom. Topol.*, 19(4):2099–2150, 2019.
- [Fad65] Edward Fadell. Generalized normal bundles for locally-flat imbeddings. *Trans. Amer. Math. Soc.*, 114:488–513, 1965.
- [Fer77] Steve Ferry. The homeomorphism group of a compact Hilbert cube manifold is an ANR. *Ann. of Math. (2)*, 106(1):101–119, 1977.
- [Fis60] Gordon M. Fisher. On the group of all homeomorphisms of a manifold. *Trans. Amer. Math. Soc.*, 97:193–212, 1960.
- [Flo22] Raphael Floris. Topological manifolds are locally-flat euclidean neighbourhood retracts, 2022. Preprint, available at arXiv:2205.05179.
- [FM66] Ralph H. Fox and John Milnor. Singularities of 2-spheres in 4-space and cobordism of knots. *Osaka J. Math.*, 3:257–267, 1966.
- [FQ90] Michael H. Freedman and Frank Quinn. *Topology of 4-manifolds*, volume 39 of *Princeton Mathematical Series*. Princeton University Press, Princeton, NJ, 1990.
- [FR01] Steve Ferry and Andrew Ranicki. A survey of Wall’s finiteness obstruction. In *Surveys on surgery theory. Vol. 2: Papers dedicated to C. T. C. Wall on the occasion of his 60th birthday*, pages 63–79. Princeton, NJ: Princeton University Press, 2001.
- [Fre82] Michael Hartley Freedman. The topology of four-dimensional manifolds. *J. Differential Geom.*, 17(3):357–453, 1982.
- [Fri12] Greg Friedman. An elementary illustrated introduction to simplicial sets. *Rocky Mountain J. Math.*, 42:353–424, 2012.
- [Fri23] Stefan Friedl. Topology. AMS open notes: # OMN:202307.111368 <https://www.ams.org/open-math-notes/omn-view-listing?listingId=111368>, 2023.
- [FS84] Ronald Fintushel and Ronald J. Stern. A μ -invariant one homology 3-sphere that bounds an orientable rational ball. In *Four-manifold theory (Durham, N.H., 1982)*, volume 35 of *Contemp. Math.*, pages 265–268. Amer. Math. Soc., Providence, RI, 1984.
- [FT95] Michael H. Freedman and Peter Teichner. 4-manifold topology. I. Subexponential groups. *Invent. Math.*, 122(3):509–529, 1995.
- [Fur01] Mikio Furuta. Monopole equation and the $\frac{11}{8}$ -conjecture. *Math. Res. Lett.*, 8(3):279–291, 2001.
- [GG73] Martin Golubitsky and Victor Guillemin. *Stable mappings and their singularities*, volume Vol. 14 of *Graduate Texts in Mathematics*. Springer-Verlag, New York-Heidelberg, 1973.
- [GH81] Marvin J. Greenberg and John R. Harper. Algebraic topology: a first course. Mathematics Lecture Note Series, 58. Reading, Massachusetts, 1981.
- [Gom84] Robert E. Gompf. Stable diffeomorphism of compact 4-manifolds. *Topology Appl.*, 18:115–120, 1984.
- [GS80] David E. Galewski and Ronald J. Stern. Classification of simplicial triangulations of topological manifolds. *Ann. of Math. (2)*, 111(1):1–34, 1980.
- [GS99] Robert E. Gompf and András Stipsicz. *4-manifolds and Kirby calculus*, volume 20 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 1999.
- [Hab82] Nathan Habegger. Une variété de dimension 4 avec forme d’intersection paire et signature -8 . *Comment. Math. Helv.*, 57(1):22–24, 1982.
- [Ham76] Andrew J. S. Hamilton. The triangulation of 3-manifolds. *Quart. J. Math. Oxford Ser. (2)*, 27(105):63–70, 1976.
- [Ham09] Ian Hambleton. Intersection forms, fundamental groups and 4-manifolds. In *Proceedings of Gökova Geometry-Topology Conference 2008*, pages 137–150. Gökova Geometry/Topology Conference (GGT), Gökova, 2009.
- [Han51] Olof Hanner. Some theorems on absolute neighborhood retracts. *Ark. Mat.*, 1:389–408, 1951.

- [Han89] Vagn Lundsgaard Hansen. *Braids and coverings: selected topics*. Cambridge etc.: Cambridge University Press, 1989.
- [Hat83] Allen Hatcher. A proof of the Smale conjecture, $\text{Diff}(S^3) \simeq O(4)$. *Ann. of Math. (2)*, 117(3):553–607, 1983.
- [Hat02] Allen Hatcher. *Algebraic topology*. Cambridge University Press, Cambridge, 2002.
- [Hat13] Allen Hatcher. The Kirby torus trick for surfaces. Available at arXiv:1312.3518, 2013.
- [HH19] Ian Hambleton and Alyson Hildum. Topological 4-manifolds with right-angled Artin fundamental groups. *J. Topol. Anal.*, 11(4):777–821, 2019.
- [Hil12] Jonathan Hillman. *Algebraic invariants of links. 2nd ed.* Singapore: World Scientific, 2nd ed. edition, 2012.
- [Hir66] Morris W. Hirsch. On normal microbundles. *Topology*, 5:229–240, 1966.
- [Hir68] Morris W. Hirsch. On tubular neighborhoods of piecewise linear and topological manifolds. In *Conference on the Topology of Manifolds (Michigan State Univ., E. Lansing, Mich., 1967)*, pages 63–80. Prindle, Weber & Schmidt, Boston, Mass., 1968.
- [Hir94] Morris Hirsch. *Differential topology*, volume 33 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1994. Corrected reprint of the 1976 original.
- [HK88] Ian Hambleton and Matthias Kreck. On the classification of topological 4-manifolds with finite fundamental group. *Math. Ann.*, 280(1):85–104, 1988.
- [HK93a] Ian Hambleton and Matthias Kreck. Cancellation, elliptic surfaces and the topology of certain four-manifolds. *J. Reine Angew. Math.*, 444:79–100, 1993.
- [HK93b] Ian Hambleton and Matthias Kreck. Cancellation of hyperbolic forms and topological four-manifolds. *J. Reine Angew. Math.*, 443:21–47, 1993.
- [HKT94] Ian Hambleton, Matthias Kreck, and Peter Teichner. Nonorientable 4-manifolds with fundamental group of order 2. *Trans. Amer. Math. Soc.*, 344(2):649–665, 1994.
- [HKT09] Ian Hambleton, Matthias Kreck, and Peter Teichner. Topological 4-manifolds with geometrically two-dimensional fundamental groups. *J. Topol. Anal.*, 1(2):123–151, 2009.
- [HLSX22] Michael J. Hopkins, Jianfeng Lin, XiaoLin Danny Shi, and Zhouli Xu. Intersection forms of spin 4-manifolds and the $\text{Pin}(2)$ -equivariant Mahowald invariant. *Comm. Amer. Math. Soc.*, 2:22–132, 2022.
- [HM74] Morris W. Hirsch and Barry Mazur. *Smoothings of piecewise linear manifolds*. Princeton University Press, Princeton, N. J.; University of Tokyo Press, Tokyo, 1974. Annals of Mathematics Studies, No. 80.
- [HS69] Wu-Chung Hsiang and Julius .L. Shaneson. Fake tori, the annulus conjecture, and the conjectures of Kirby. *Proc. Natl. Acad. Sci. USA*, 62:687–691, 1969.
- [HS97] Peter J. Hilton and Urs Stambach. *A course in homological algebra. 2nd ed.* New York, NY: Springer, 2nd ed. edition, 1997.
- [Hud69] John F. P. Hudson. On transversality. *Proc. Cambridge Philos. Soc.*, 66:17–20, 1969.
- [Hus93] Dale Husemoller. *Fibre bundles. Graduate Texts in Mathematics*, 20, 1993. third edition.
- [HW69] Wu-Chung Hsiang and C. Terence C. Wall. On homotopy tori. II. *Bull. London Math. Soc.*, 1:341–342, 1969.
- [Kea75] Cherry Kearton. Cobordism of knots and Blanchfield duality. *J. London Math. Soc. (2)*, 10(4):406–408, 1975.
- [Kir69] Robion C. Kirby. Stable homeomorphisms and the annulus conjecture. *Ann. of Math. (2)*, 89:575–582, 1969.
- [Kir89] Robion C. Kirby. *The topology of 4-manifolds*, volume 1374 of *Lect. Notes Math.* Berlin etc.: Springer-Verlag, 1989.
- [Kis64] James M. Kister. Microbundles are fibre bundles. *Ann. of Math. (2)*, 80:190–199, 1964.

- [KL66] Nicolaas H. Kuiper and Richard Lashof. Microbundles and bundles. II. Semisimplicial theory. *Invent. Math.*, 1:243–259, 1966.
- [KL22] Daniel Kasprowski and Markus Land. Topological 4-manifolds with 4-dimensional fundamental group. *Glasg. Math. J.*, 64(2):454–461, 2022.
- [KLPT17] Daniel Kasprowski, Markus Land, Mark Powell, and Peter Teichner. Stable classification of 4-manifolds with 3-manifold fundamental groups. *J. Topol.*, 10(3):827–881, 2017.
- [KM63] Michel A. Kervaire and John Milnor. Groups of homotopy spheres. I. *Ann. Math. (2)*, 77:504–537, 1963.
- [Kos93] Antoni A. Kosinski. *Differential manifolds*, volume 138 of *Pure and Applied Mathematics*. Academic Press, Inc., Boston, MA, 1993.
- [KPRT22] Daniel Kasprowski, Mark Powell, Arunima Ray, and Peter Teichner. Embedding surfaces in 4-manifolds. *arXiv e-prints*, 2022.
- [KQ00] Vyacheslav S. Krushkal and Frank Quinn. Subexponential groups in 4-manifold topology. *Geom. Topol.*, 4:407–430, 2000.
- [Kre84] Matthias Kreck. Some closed 4-manifolds with exotic differentiable structure. In *Algebraic topology, Aarhus 1982 (Aarhus, 1982)*, volume 1051 of *Lecture Notes in Math.*, pages 246–262. Springer, Berlin, 1984.
- [Kre99] Matthias Kreck. Surgery and duality. *Ann. of Math. (2)*, 149(3):707–754, 1999.
- [KS69] Robion C. Kirby and Laurence C. Siebenmann. On the triangulation of manifolds and the Hauptvermutung. *Bull. Amer. Math. Soc.*, 75:742–749, 1969.
- [KS75] Robion C. Kirby and Laurence C. Siebenmann. Normal bundles for codimension 2 locally flat imbeddings. In *Geometric topology (Proc. Conf., Park City, Utah, 1974)*, pages 310–324. Lecture Notes in Math., Vol. 438. Springer, Berlin, 1975.
- [KS77] Robion C. Kirby and Laurence C. Siebenmann. *Foundational essays on topological manifolds, smoothings, and triangulations*. Princeton University Press, Princeton, N.J.; University of Tokyo Press, Tokyo, 1977. With notes by John Milnor and Michael Atiyah, *Annals of Mathematics Studies*, No. 88.
- [KS18] Sławomir Kwasik and Fang Sun. Local coefficients revisited. *Enseign. Math. (2)*, 64(3-4):283–303, 2018.
- [Lam91] Tsit Y. Lam. *A first course in noncommutative rings*. New York etc.: Springer-Verlag, 1991.
- [Lan02a] Serge Lang. *Algebra. 3rd revised ed.* New York, NY: Springer, 3rd revised ed. edition, 2002.
- [Lan02b] Serge Lang. *Introduction to differentiable manifolds*. Universitext. Springer-Verlag, New York, second edition, 2002.
- [Las70] Richard Lashof. The immersion approach to triangulation. In *Topology of Manifolds (Proc. Inst., Univ. of Georgia, Athens, Ga., 1969)*, pages 52–56. Markham, Chicago, Ill., 1970.
- [Las71] Richard Lashof. The immersion approach to triangulation and smoothing. In *Algebraic topology (Proc. Sympos. Pure Math., Vol. XXII, Univ. Wisconsin, Madison, Wis., 1970)*, pages 131–164. Amer. Math. Soc., Providence, R.I., 1971.
- [Lee69] J. Alexander Lees. Immersions and surgeries of topological manifolds. *Bull. Am. Math. Soc.*, 75:529–534, 1969.
- [Lee11] John Lee. *Introduction to topological manifolds*, volume 202 of *Graduate Texts in Mathematics*. Springer, New York, second edition, 2011.
- [Lev67] Jerome Levine. A method for generating link polynomials. *Am. J. Math.*, 89:69–84, 1967.
- [Lev69] Jerome Levine. Knot cobordism groups in codimension two. *Comment. Math. Helv.*, 44:229–244, 1969.
- [Lev77] Jerome Levine. Knot modules. I. *Trans. Am. Math. Soc.*, 229:1–50, 1977.
- [LM19] Lukas Lewark and Duncan McCoy. On calculating the slice genera of 11- and 12-crossing knots. *Exp. Math.*, 28(1):81–94, 2019.

- [LS71] Richard Lashof and Julius L. Shaneson. Smoothing 4-manifolds. *Invent. Math.*, 14:197–210, 1971.
- [LS03] Wolfgang Lück and Thomas Schick. Various L^2 -signatures and a topological L^2 -signature theorem. In *High-dimensional manifold topology. Proceedings of the school, ICTP, Trieste, Italy, May 21–June 8, 2001*, pages 362–399. River Edge, NJ: World Scientific, 2003.
- [LT84] Richard Lashof and Laurence Taylor. Smoothing theory and Freedman’s work on four-manifolds. In *Algebraic topology, Aarhus 1982 (Aarhus, 1982)*, volume 1051 of *Lecture Notes in Math.*, pages 271–292. Springer, Berlin, 1984.
- [Lüc98] Wolfgang Lück. Dimension theory of arbitrary modules over finite von Neumann algebras and L^2 -Betti numbers. I: Foundations. *J. Reine Angew. Math.*, 495:135–162, 1998.
- [Lüc02] Wolfgang Lück. *L^2 -invariants: theory and applications to geometry and K-theory*, volume 44 of *Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics*. Springer-Verlag, Berlin, 2002.
- [Lüc10] Wolfgang Lück. Survey on aspherical manifolds. In *European congress of mathematics. Proceedings of the 5th ECM congress, Amsterdam, Netherlands, July 14–18, 2008*, pages 53–82. Zürich: European Mathematical Society (EMS), 2010.
- [Man16] Ciprian Manolescu. Pin(2)-equivariant Seiberg-Witten Floer homology and the triangulation conjecture. *J. Amer. Math. Soc.*, 29(1):147–176, 2016.
- [Mar58] Andrei Markov. Unsolvability of certain problems in topology. *Dokl. Akad. Nauk SSSR*, 123:978–980, 1958.
- [Mar77] Alexis Marin. La transversalité topologique. *Ann. of Math. (2)*, 106(2):269–293, 1977.
- [Mat78] Takao Matumoto. Triangulation of manifolds. In *Algebraic and geometric topology (Proc. Sympos. Pure Math., Stanford Univ., Stanford, Calif., 1976), Part 2*, Proc. Sympos. Pure Math., XXXII, pages 3–6. Amer. Math. Soc., Providence, R.I., 1978.
- [Mau80] C. Richard F. Maunder. *Algebraic topology*. Cambridge University Press, 1980.
- [May75] J. Peter May. Classifying spaces and fibrations. *Mem. Amer. Math. Soc.*, 1(1, 155):xiii+98, 1975.
- [Maz59] Barry Mazur. On embeddings of spheres. *Bull. Amer. Math. Soc.*, 65:59–65, 1959.
- [MH73] John Milnor and Dale Husemoller. *Symmetric bilinear forms*. Springer-Verlag, New York-Heidelberg, 1973. Ergebnisse der Mathematik und ihrer Grenzgebiete, Band 73.
- [Mil56] John Milnor. Construction of universal bundles. II. *Ann. Math. (2)*, 63:430–436, 1956.
- [Mil57] John Milnor. The geometric realization of a semi-simplicial complex. *Ann. of Math. (2)*, 65:357–362, 1957.
- [Mil58] John Milnor. On simply connected 4-manifolds. In *Symposium internacional de topología algebraica International symposium on algebraic topology*, pages 122–128. Universidad Nacional Autónoma de México and UNESCO, Mexico City, 1958.
- [Mil61] John Milnor. Microbundles and differentiable structures. *Princeton*, 1961. Mimeographed.
- [Mil63] John Milnor. *Morse theory*. Based on lecture notes by M. Spivak and R. Wells. Annals of Mathematics Studies, No. 51. Princeton University Press, Princeton, N.J., 1963.
- [Mil64] John Milnor. Microbundles: Part I. *Topology*, 3(1):53–80, 1964.
- [Mil72] R. T. Miller. Approximating codimension 3 embeddings. *Ann. Math.*, (2) 95:406–416, 1972.
- [Moi52] Edwin Moise. Affine structures in 3-manifolds. V. The triangulation theorem and Hauptvermutung. *Ann. of Math. (2)*, 56:96–114, 1952.
- [Moi77] Edwin Moise. Geometric topology in dimensions 2 and 3. Graduate Texts in Mathematics. 47., 1977.
- [Mor60] Marston Morse. A reduction of the Schoenflies extension problem. *Bull. Amer. Math. Soc.*, 66:113–115, 1960.
- [MS74] John Milnor and James D. Stasheff. *Characteristic classes*. Princeton University Press, Princeton, N. J., 1974. Annals of Mathematics Studies, No. 76.

- [MS17] Dusa McDuff and Dietmar Salamon. *Introduction to symplectic topology*. Oxford Graduate Texts in Mathematics. Oxford University Press, Oxford, third edition, 2017.
- [MT07] John Morgan and Gang Tian. *Ricci flow and the Poincaré conjecture*, volume 3 of *Clay Mathematics Monographs*. American Mathematical Society, Providence, RI; Clay Mathematics Institute, Cambridge, MA, 2007.
- [Mun59] James Munkres. Obstructions to the smoothing of piecewise-differentiable homeomorphisms. *Bull. Amer. Math. Soc.*, 65:332–334, 1959.
- [Mun60a] James Munkres. Differentiable isotopies on the 2-sphere. *Mich. Math. J.*, 7:193–197, 1960.
- [Mun60b] James Munkres. Obstructions to the smoothing of piecewise-differentiable homeomorphisms. *Ann. of Math.*, (2) 72:521–554, 1960.
- [Mun66] James R. Munkres. *Elementary differential topology*, volume 1961 of *Lectures given at Massachusetts Institute of Technology, Fall*. Princeton University Press, Princeton, N.J., 1966.
- [New66a] Maxwell H. A. Newman. The engulfing theorem for topological manifolds. *Ann. of Math.* (2), 84:555–571, 1966.
- [New66b] Maxwell H. A. Newman. The engulfing theorem for topological manifolds. *Ann. Math.* (2), 84:555–571, 1966.
- [Nic79] Andrew John Nicas. *Induction theorems for groups of homotopy manifold structures*. ProQuest LLC, Ann Arbor, MI, 1979. Thesis (Ph.D.)—Princeton University.
- [Nov64] S. P. Novikov. Homotopically equivalent smooth manifolds. I. *Izv. Akad. Nauk SSSR Ser. Mat.*, 28:365–474, 1964.
- [NP17] Matthias Nagel and Mark Powell. Concordance invariance of Levine-Tristram signatures of links. *Doc. Math.*, 22:25–43, 2017.
- [OP22] Patrick Orson and Mark Powell. Mapping class groups of simply connected 4-manifolds with boundary. *arXiv e-prints*, 2022.
- [Osb00] M.Scott Osborne. *Basic homological algebra*. New York, NY: Springer, 2000.
- [Pap46] Christos Papakyriakopoulos. A new proof for the invariance of the homology groups of a complex. *Bull. Soc. Math. Greece*, 2:1–150, 1946.
- [Pap57] Christos D. Papakyriakopoulos. On solid tori. *Proc. London Math. Soc.* (3), 7:281–299, 1957.
- [Per86] Bernard Perron. Pseudo-isotopies et isotopies en dimension quatre dans la catégorie topologique. *Topology*, 25(4):381–397, 1986.
- [Per02] Grigori Perelman. The entropy formula for the Ricci flow and its geometric applications. Preprint, available at arXiv:0211159, 2002.
- [Per03a] Grigori Perelman. Finite extinction time for the solutions to the Ricci flow on certain three-manifolds. Preprint, available at arXiv:0307245, 2003.
- [Per03b] Grigori Perelman. Ricci flow with surgery on three-manifolds. Preprint, available at arXiv:0303109, 2003.
- [Pes96] Christian Peskine. *An algebraic introduction to complex projective geometry. 1: Commutative algebra.*, volume 47. Cambridge: Cambridge Univ. Press, 1996.
- [PRT19] Mark Powell, Arunima Ray, and Peter Teichner. The 4-dimensional disc embedding theorem and dual spheres. *arXiv e-prints*, 2019.
- [Qui70] Frank Quinn. A geometric formulation of surgery. In *Topology of Manifolds (Proc. Inst., Univ. of Georgia, Athens, Ga., 1969)*, pages 500–511. Markham Publishing Co., Chicago, IL, 1970.
- [Qui82] Frank Quinn. Ends of maps III: Dimensions 4 and 5. *J. Differential Geom.*, 17(3):503–521, 1982.
- [Qui83] Frank Quinn. The stable topology of 4-manifolds. *Topology Appl.*, 15(1):71–77, 1983.
- [Qui84] Frank Quinn. Smooth structures on 4-manifolds. In *Four-manifold theory (Durham, N.H., 1982)*, volume 35 of *Contemp. Math.*, pages 473–479. Amer. Math. Soc., Providence, RI, 1984.
- [Qui86] Frank Quinn. Isotopy of 4-manifolds. *J. Differential Geom.*, 24(3):343–372, 1986.

- [Qui88] Frank Quinn. Topological transversality holds in all dimensions. *Bull. Amer. Math. Soc. (N.S.)*, 18(2):145–148, 1988.
- [Rad26] Tibor Radó. Über den Begriff der Riemannschen Fläche. *Acta Szeged*, 2:101–121, 1926.
- [Ran92] Andrew A. Ranicki. *Algebraic L-theory and topological manifolds*, volume 102 of *Cambridge Tracts in Mathematics*. Cambridge University Press, Cambridge, 1992.
- [Roh52] Vladimir A. Rohlin. New results in the theory of four-dimensional manifolds. *Doklady Akad. Nauk SSSR (N.S.)*, 84:221–224, 1952.
- [Rol90] Dale Rolfsen. *Knots and links*, volume 7 of *Mathematics Lecture Series*. Publish or Perish, Inc., Houston, TX, 1990. Corrected reprint of the 1976 original.
- [Row06] Louis Halle Rowen. *Graduate algebra: commutative view*, volume 73. Providence, RI: American Mathematical Society (AMS), 2006.
- [RS67] Colin P. Rourke and Brian J. Sanderson. An embedding without a normal microbundle. *Invent. Math.*, 3:293–299, 1967.
- [RS68] Colin P. Rourke and Brian J. Sanderson. Block bundles. III. Homotopy theory. *Ann. of Math. (2)*, 87:431–483, 1968.
- [RS72] Colin P. Rourke and Brian J. Sanderson. *Introduction to piecewise-linear topology*. Springer-Verlag, New York-Heidelberg, 1972. *Ergebnisse der Mathematik und ihrer Grenzgebiete, Band 69*.
- [Rub84] Daniel Ruberman. Invariant knots of free involutions of S^4 . *Topology Appl.*, 18(2-3):217–224, 1984.
- [Rud16] Yuli Rudyak. *Piecewise linear structures on topological manifolds*. Hackensack, NJ: World Scientific, 2016.
- [Sak13] Katsuro Sakai. *Geometric aspects of general topology*. Springer Monogr. Math. Tokyo: Springer, 2013.
- [Sav02] Nikolai Saveliev. *Invariants for homology 3-spheres*, volume 140 of *Encycl. Math. Sci.* Berlin: Springer, 2002.
- [Sch68] Horst Schubert. *Topology*. Boston: Allyn and Bacon, Inc. 1968, 358 p. (1968)., 1968.
- [Sch70] James A. Schafer. Topological Pontrjagin classes. *Comment. Math. Helv.*, 45:315–332, 1970.
- [Sha84] Peter Shalen. A “piecewise-linear” method for triangulating 3-manifolds. *Adv. in Math.*, 2:34–80, 1984.
- [Sma62] Stephen Smale. On the structure of manifolds. *Amer. J. Math.*, pages 387–399, 1962.
- [Sta63] John Stallings. On topologically unknotted spheres. *Ann. of Math. (2)*, 77:490–503, 1963.
- [Sta71] James D. Stasheff. H-spaces and classifying spaces: Foundations and recent developments. *Algebraic Topology, Proc. Sympos. Pure Math. 22*, 247–272 (1971)., 1971.
- [Ste62] Norman E. Steenrod. *Cohomology operations*. *Annals of Mathematics Studies*, No. 50. Princeton University Press, Princeton, N.J., 1962. Lectures by N. E. Steenrod written and revised by D. B. A. Epstein.
- [Ste75] Ronald J. Stern. On topological and piecewise linear vector fields. *Topology*, 14:257–269, 1975.
- [Sto93] Richard Stong. Simply-connected 4-manifolds with a given boundary. *Topology Appl.*, 52(2):161–167, 1993.
- [Sto94] Richard Stong. Uniqueness of connected sum decompositions in dimension 4. *Topology and its applications*, 56:277–291, 1994.
- [Sul66] Dennis Parnell Sullivan. *Triangulating homotopy equivalences*. ProQuest LLC, Ann Arbor, MI, 1966. Thesis (Ph.D.)—Princeton University.
- [Sun17] Fang Sun. An elementary proof for Poincare Duality with local coefficients. *arXiv e-prints*, 2017.
- [SW00] Richard Stong and Zhenghan Wang. Self-homeomorphisms of 4-manifolds with fundamental group \mathbf{z} . *Topology Appl.*, 106(1):49–56, 2000.

- [Swi02] Robert M. Switzer. *Algebraic topology—homotopy and homology*. Classics in Mathematics. Springer-Verlag, Berlin, 2002. Reprint of the 1975 original [Springer, New York; MR0385836 (52 #6695)].
- [Tei92] Peter Teichner. *Topological 4-manifolds with finite fundamental group*. PhD thesis, University of Mainz, Germany, 1992. Shaker Verlag, ISBN 3-86111-182-9.
- [Tei93] Peter Teichner. On the signature of four-manifolds with universal covering spin. *Math. Ann.*, 295(4):745–759, 1993.
- [Tei97] Peter Teichner. On the star-construction for topological 4-manifolds. In *Geometric topology (Athens, GA, 1993)*, volume 2 of *AMS/IP Stud. Adv. Math.*, pages 300–312. Amer. Math. Soc., Providence, RI, 1997.
- [Tel84] Nicolae Teleman. The index theorem for topological manifolds. *Acta Math.*, 153:117–152, 1984.
- [Tho52] René Thom. Espaces fibrés en sphères et carrés de Steenrod. *Ann. Sci. Ecole Norm. Sup. (3)*, 69:109–182, 1952.
- [Thu97] William P. Thurston. Three-dimensional geometry and topology. Vol. 1. *Princeton Mathematical Series*, 35, 1997. Edited by Silvio Levy.
- [Tri69] Andrew G. Tristram. Some cobordism invariants for links. *Proc. Cambridge Philos. Soc.*, 66:251–264, 1969.
- [Tur01] Vladimir Turaev. *Introduction to combinatorial torsions*. Lectures in Mathematics ETH Zürich. Birkhäuser Verlag, Basel, 2001. Notes taken by Felix Schlenk.
- [Wal62] C. Terence C. Wall. Classification of $(n-1)$ -connected $2n$ -manifolds. *Ann. Math. (2)*, 75:163–189, 1962.
- [Wal64] C. Terence C. Wall. On simply-connected 4-manifolds. *J. London Math. Soc.*, 39:141–149, 1964.
- [Wal65] C. Terence C. Wall. Finiteness conditions for CW-complexes. I, II. *Ann. Math. (2)*, 81:56–69, 1965.
- [Wal66] C. Terence C. Wall. Finiteness conditions for CW complexes. II. *Proc. Roy. Soc. Ser. A*, 295:129–139, 1966.
- [Wal67] C. Terence C. Wall. Poincaré complexes. I. *Ann. of Math. (2)*, 86:213–245, 1967.
- [Wal69] C. Terence C. Wall. On homotopy tori and the annulus theorem. *Bull. Lond. Math. Soc.*, 1:95–97, 1969.
- [Wal99] C. Terence C. Wall. *Surgery on compact manifolds*, volume 69 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, second edition, 1999. Edited and with a foreword by Andrew A. Ranicki.
- [Wal16] C. Terence C. Wall. *Differential topology*. Cambridge: Cambridge University Press, 2016.
- [Wan95] Zhenghan Wang. Classification of closed nonorientable 4-manifolds with infinite cyclic fundamental group. *Mathematical Research Letters*, 2:339–344, 01 1995.
- [Wes77] James West. Mapping Hilbert cube manifolds to ANR’s: a solution of a conjecture of Borsuk. *Ann. of Math. (2)*, 106(1):1–18, 1977.
- [Whi49] J. H. C. Whitehead. On simply connected, 4-dimensional polyhedra. *Comment. Math. Helv.*, 22:48–92, 1949.
- [Whi57] Hassler Whitney. *Geometric integration theory*. Princeton University Press, Princeton, N. J., 1957.
- [Whi61] J. H. C. Whitehead. Manifolds with transverse fields in euclidean space. *Ann. of Math.*, (2) 73:154–212, 1961.
- [Zee62] E. Christopher Zeeman. Polyhedral n -manifolds. I: Foundations. In *Topology of 3-manifolds and related topics. Proceedings of the University of Georgia Institute 1961.*, pages 57–64. Englewood Cliffs, N.J.: Prentice-Hall, Inc., 1962.
- [Zee63] E. Christopher Zeeman. Unknotting combinatorial balls. *Ann. of Math. (2)*, 78:501–526, 1963.