

*Pacific
Journal of
Mathematics*

**INFINITE HOMOTOPY STABLE CLASS
FOR 4-MANIFOLDS WITH BOUNDARY**

ANTHONY CONWAY, DIARMUID CROWLEY AND MARK POWELL

INFINITE HOMOTOPY STABLE CLASS FOR 4-MANIFOLDS WITH BOUNDARY

ANTHONY CONWAY, DIARMUID CROWLEY AND MARK POWELL

We show that for every odd prime q , there exists an infinite family $\{M_i\}_{i=1}^\infty$ of topological 4-manifolds that are all stably homeomorphic to one another, all the manifolds M_i have isometric rank one equivariant intersection pairings and boundary $L(2q, 1)\#(S^1 \times S^2)$, but they are pairwise not homotopy equivalent via any homotopy equivalence that restricts to a homotopy equivalence of the boundary.

1. Introduction

In what follows a manifold is understood to mean a compact, connected, oriented, topological manifold. Let $W_g := \#_g(S^2 \times S^2)$ be the g -fold connected sum of $S^2 \times S^2$ with itself. Two 4-manifolds M and N with the same Euler characteristic are *stably homeomorphic*, denoted $M \cong_{\text{st}} N$, if there exists a nonnegative integer g and a homeomorphism

$$M \# W_g \cong N \# W_g.$$

Surgery theory suggests two ways to classify 4-manifolds. The classical Browder–Novikov–Sullivan–Wall [Wall 1999] approach is to classify up to homotopy equivalence and then employ the surgery exact sequence. Kreck’s modified surgery approach [1999] seeks to classify up to stable homeomorphism, and then attempt to destabilise. A natural question then arising is to compare the homotopy and stable classifications. To do this precisely for 4-manifolds with boundary we fix a 4-manifold M and define the *homotopy stable class*:

$$\mathcal{S}_h^{\text{st}}(M) := \{N \mid N \cong_{\text{st}} M\} / \text{homotopy equivalence of pairs}.$$

Here, we understand a *homotopy equivalence of pairs* $N_1 \simeq N_2$ to be one that restricts to a homotopy equivalence between the boundaries. When the manifolds are closed, this recovers the usual notion of homotopy equivalence.

Using the equivariant intersection form λ_N of N as an invariant, $\mathcal{S}_h^{\text{st}}(M)$ can be arbitrarily large: for example, one can use Freedman’s work [1982] to realise

MSC2020: 57K40, 57R65.

Keywords: stable homeomorphism, homotopy equivalence, 4-manifold.

distinct positive definite symmetric bilinear forms with the same signature and rank by simply connected closed 4-manifolds with identical Kirby–Siebenmann invariant. For this reason, we study the homotopy stable class one intersection form at a time and set

$$\mathcal{S}_{h,\lambda}^{\text{st}}(M) := \{N \mid N \cong_{\text{st}} M, \lambda_N \cong \lambda_M\} / \text{homotopy equivalence of pairs}.$$

If M is closed and has $\pi_1(M) = 1, \mathbb{Z}$, or \mathbb{Z}/n , or $\pi_1(M)$ is a solvable Baumslag–Solitar group, then $|\mathcal{S}_{h,\lambda}^{\text{st}}(M)| = 1$: stably homeomorphic manifolds with isometric equivariant intersection forms are homeomorphic for $\pi_1 = 1$ by [Freedman 1982], for $\pi_1 \cong \mathbb{Z}$ by [Freedman and Quinn 1990], for $\pi_1 \cong \mathbb{Z}/n$ by [Hambleton and Kreck 1993, Theorem C], and for solvable Baumslag–Solitar group by [Hambleton, Kreck and Teichner 2009, Theorem A]. On the other hand, Kreck and Schafer [1984] found pairs of smooth closed 4-manifolds with finite π_1 and isometric equivariant intersection forms that are stably diffeomorphic but not homotopy equivalent. When the boundary is nonempty and $\pi_1 = 1$, one can use work of Boyer [1993] to produce simply connected 4-manifolds M with boundary and arbitrarily large (but necessarily finite) $\mathcal{S}_{h,\lambda}^{\text{st}}(M)$. Until now however, there have been no examples of 4-manifolds with infinite $\mathcal{S}_{h,\lambda}^{\text{st}}(M)$. For every odd prime q , our main result describes a 4-manifold M with fundamental group \mathbb{Z} and infinite $\mathcal{S}_{h,\lambda_{2q}}^{\text{st}}(M)$, where the fixed Hermitian form is

$$\lambda_{2q} : \mathbb{Z}[t^{\pm 1}] \times \mathbb{Z}[t^{\pm 1}] \rightarrow \mathbb{Z}[t^{\pm 1}]; \quad (x, y) \mapsto 2qx\bar{y}.$$

Theorem 1.1. *For every odd prime q , there exists an infinite family $\{M_i\}_{i=1}^{\infty}$ of 4-manifolds with fundamental group \mathbb{Z} that are all stably homeomorphic, and all the manifolds M_i have equivariant intersection pairing isometric to λ_{2q} and boundary $L(2q, 1)\#(S^1 \times S^2)$, but they are pairwise not homotopy equivalent via any homotopy equivalence that restricts to a homotopy equivalence on the boundary. In other words,*

$$|\mathcal{S}_{h,\lambda_{2q}}^{\text{st}}(M_1)| = \infty.$$

For a fixed odd prime q , the manifolds in Theorem 1.1 all have fundamental group \mathbb{Z} , boundary

$$Y_q := L(2q, 1)\#(S^1 \times S^2),$$

equivariant intersection form isometric to λ_{2q} , and integral intersection form isometric to

$$\lambda_{2q}^{\mathbb{Z}} : \mathbb{Z} \times \mathbb{Z} \rightarrow \mathbb{Z}; \quad (x, y) \mapsto 2qxy,$$

but are distinguished by an invariant, first introduced in [Conway, Piccirillo and Powell 2022] and inspired by [Boyer 1993], related to the Blanchfield form of Y_q . While the manifold M_1 is smooth, we cannot tell whether any of the other M_i admit smooth structures. Their construction uses surgery methods, in particular a recent

realisation result from [Conway, Piccirillo and Powell 2022], which a priori only works in the topological category.

Before giving more details and describing the main steps in the proof of [Theorem 1.1](#), we briefly compare the study of the homotopy stable class in dimension 4 with the situation in higher dimensions.

Remark 1.2. Kreck and Schafer [1984] found pairs of closed smooth $4k$ -manifolds, for $k \geq 1$, that are stably diffeomorphic and have hyperbolic equivariant intersection forms, but are pairwise not homotopy equivalent. In [Conway, Crowley, Powell and Sixt 2023], we gave the first examples of simply connected, closed, smooth $4k$ -manifolds, for $k \geq 2$, with hyperbolic intersection form and arbitrarily large homotopy stable class $\mathcal{S}_{h,\lambda}^{\text{st}}$. In [Conway, Crowley, Powell and Sixt 2021] for $k \geq 2$, we produced smooth closed $4k$ -manifolds with fundamental group \mathbb{Z} , again with hyperbolic intersection form, and such that the homotopy stable class $\mathcal{S}_{h,\lambda}^{\text{st}}$ is infinite. In those papers we were unable to obtain examples in dimension 4. In [Conway, Crowley, Powell and Sixt 2023], in lieu of this we defined a spin^c version of the stable class in dimension 4, and we showed that this spin^c stable class can be arbitrarily large. This article shows that a variation on those methods, with analogous underlying algebra, does produce examples of 4-manifolds with nonempty boundary and fundamental group \mathbb{Z} that have infinite homotopy stable class.

Next we describe the main steps in the proof of [Theorem 1.1](#). Fix an odd prime q . The first observation is that if N_1, N_2 are 4-manifolds with integral intersection forms isometric to $\lambda_{2q}^{\mathbb{Z}}$, then there can be no orientation-reversing homotopy equivalence between N_1 and N_2 . For this reason, and for the purpose of proving our main theorem, we restrict to orientation-preserving homotopy equivalences (o.p. homotopy eq. for short) and therefore consider

$$\mathcal{S}_{h^+,\lambda}^{\text{st}}(M) := \{N \mid N \cong_{\text{st}} M, \lambda_N \cong \lambda_M\} / \text{o.p. homotopy eq. of pairs.}$$

We now restrict to 4-manifolds M with fundamental group \mathbb{Z} such that the inclusion $\partial M \subseteq M$ induces a surjection $\varphi : \pi_1(\partial M) \twoheadrightarrow \pi_1(M) \cong \mathbb{Z}$ (we say that M has *ribbon boundary*) and for which $H_1(\partial M; \mathbb{Z}[t^{\pm 1}])$ is a $\mathbb{Z}[t^{\pm 1}]$ -torsion module. Here and throughout the paper we assume that the fundamental groups of our 4-manifolds are equipped with a preferred isomorphism to \mathbb{Z} ; to indicate this we write $\pi_1(M) = \mathbb{Z}$.

Given two such manifolds N_1 and N_2 , we write $\partial N_1 \cong_B \partial N_2$ if there exists an orientation-preserving homeomorphism $f : \partial N_1 \xrightarrow{\cong} \partial N_2$ that intertwines the inclusion induced epimorphisms $\varphi_i : \pi_1(\partial N_i) \twoheadrightarrow \pi_1(N_i)$ and, in the case that N_1 and N_2 are spin, such that the union $N_1 \cup_f -N_2$ is spin. The terminology \cong_B is motivated by modified surgery theory [Kreck 1999], in which B is the standard notation for the normal 1-type.

Next, if N_1 and N_2 have fundamental group \mathbb{Z} , $\partial N_1 \cong_B \partial N_2$, the same Kirby–Siebenmann invariant, and $\lambda_{N_1} \cong \lambda_{N_2}$, then they are stably homeomorphic. Indeed, N_1 and N_2 must have isometric integral intersection forms (in particular with the same type and the same signature) and the same Kirby–Siebenmann invariant, so [Kreck 1999, Theorem 2] ensures they are stably homeomorphic; see Lemma 4.1 for details.

Put differently, if M is a 4-manifold with infinite cyclic fundamental group, then

$$\mathbb{S}_{h^+, \lambda}^{\text{st}}(M) = \frac{\{N \mid \partial N \cong_B \partial M, \pi_1(N) = \mathbb{Z}, \lambda_N \cong \lambda_M, \text{ks}(N) = \text{ks}(M)\}}{\text{o.p. homotopy eq. of pairs}}.$$

This next step is to recast $\mathbb{S}_{h^+, \lambda}^{\text{st}}(M)$ in terms of the group $\text{Aut}(\text{Bl}_{\partial M})$ of isometries of the Blanchfield form $\text{Bl}_{\partial M}$ (whose definition we recall in Section 3). Firstly, as we recall in Section 5, the group $\text{hAut}_{\varphi}^+(\partial M)$ of orientation-preserving homotopy equivalences $h : \partial M \simeq \partial M$ that intertwine the inclusion induced map $\varphi : \pi_1(\partial M) \rightarrow \pi_1(M) = \mathbb{Z}$ acts on $\text{Aut}(\text{Bl}_{\partial M})$. Secondly, as we also recall in Section 5, the group $\text{Aut}(\lambda_M)$ of isometries of λ_M also acts on $\text{Aut}(\text{Bl}_{\partial M})$, and the two actions commute with one another. Quotienting out by these two actions leads to an orbit set $\text{Aut}(\text{Bl}_{\partial M})/(\text{Aut}(\lambda_M) \times \text{hAut}_{\varphi}^+(\partial M))$. Note that it need not be group.

In order to account for our 4-manifolds being spin, we will in fact need to work with a smaller set of isometries. Namely, if M is spin, then $\text{Bl}_{\partial M}$ admits a quadratic enhancement

$$\mu_{\text{Bl}_{\partial M}} : H_1(\partial M; \mathbb{Z}[t^{\pm 1}]) \rightarrow \frac{\{b \in \mathbb{Q}(t) \mid b = \bar{b}\}}{\{a + \bar{a} \mid a \in \mathbb{Z}[t^{\pm 1}]\}}$$

and we write $\text{Aut}(\text{Bl}_{\partial M}, \mu_{\text{Bl}_{\partial M}}) \subseteq \text{Aut}(\text{Bl}_{\partial M})$ for those isometries of $\text{Bl}_{\partial M}$ that also preserve $\mu_{\text{Bl}_{\partial M}}$. Writing $\text{hAut}_{\varphi}^{+,q}(\partial M)$ for those homotopy equivalences whose induced map on the Alexander module preserves $\mu_{\text{Bl}_{\partial M}}$ leads to the orbit set

$$\text{Aut}(\text{Bl}_{\partial M}, \mu_{\text{Bl}_{\partial M}})/(\text{Aut}(\lambda_M) \times \text{hAut}_{\varphi}^{+,q}(\partial M)).$$

One of the main steps in the proof of Theorem 1.1 is the following partial description of $\mathbb{S}_{h^+, \lambda}^{\text{st}}(M)$ for a large class of 4-manifolds M with infinite cyclic fundamental group and ribbon boundary. As we will explain in Proposition 5.2, this result follows fairly promptly from the machinery developed in [Conway, Piccirillo and Powell 2022]. In the following proposition, and throughout the paper, *spin* refers to a manifold that admits a spin structure compatible with the orientation.

Proposition 1.3. *If M is a spin 4-manifold with ribbon boundary, $\pi_1(M) = \mathbb{Z}$, and nondegenerate equivariant intersection form λ_M , then there is a surjection*

$$b : \mathbb{S}_{h^+, \lambda}^{\text{st}}(M) \twoheadrightarrow \text{Aut}(\text{Bl}_{\partial M}, \mu_{\text{Bl}_{\partial M}})/(\text{Aut}(\lambda_M) \times \text{hAut}_{\varphi}^{+,q}(\partial M)).$$

The surjection is described explicitly in Construction 5.1.

Fix an odd prime q and let $X_{2q}(U)$ denote the $2q$ -trace on the unknot U , i.e. the smooth 4-manifold obtained from D^4 by attaching a $2q$ -framed 2-handle along the unknot. The final part of the proof of [Theorem 1.1](#), which is carried out in [Proposition 6.8](#), consists of proving that for $M = X_{2q}(U) \natural (S^1 \times D^3)$, the set

$$\text{Aut}(\text{Bl}_{\partial M}, \mu_{\text{Bl}_{\partial M}}) / (\text{Aut}(\lambda_M) \times \text{hAut}_{\varphi}^{+,q}(\partial M))$$

is countably infinite. Modulo this statement, we can now conclude the proof of [Theorem 1.1](#), which states that $\mathcal{S}_{h, \lambda_{2q}}^{\text{st}}(M)$ is infinite.

Proof of Theorem 1.1. Fix an odd prime q and consider $M := X_{2q}(U) \natural (S^1 \times D^3)$. This 4-manifold is spin, has ribbon boundary, admits an identification $\pi_1(M) = \mathbb{Z}$, and has nondegenerate equivariant intersection $\lambda_M \cong (2q)$. Since for any two 4-manifolds N_1 and N_2 with integral intersection forms isometric to $\lambda_{2q}^{\mathbb{Z}}$, there is no orientation reversing homotopy equivalence between them, $\mathcal{S}_{h, \lambda_{2q}}^{\text{st}}(M) = \mathcal{S}_{h^+, \lambda_{2q}}^{\text{st}}(M)$. We therefore prove that $\mathcal{S}_{h^+, \lambda_{2q}}^{\text{st}}(M)$ is infinite. To prove this we apply [Proposition 1.3](#), which implies that $\mathcal{S}_{h^+, \lambda_{2q}}^{\text{st}}(M)$ surjects onto the orbit set

$$\text{Aut}(\text{Bl}_{\partial M}, \mu_{\text{Bl}_{\partial M}}) / (\text{Aut}(\lambda_M) \times \text{hAut}_{\varphi}^{+,q}(\partial M))$$

and this latter set is countably infinite by [Proposition 6.8](#). \square

Remark 1.4. The existence of M with infinite

$$\text{Aut}(\text{Bl}_{\partial M}, \mu_{\text{Bl}_{\partial M}}) / (\text{Aut}(\lambda_M) \times \text{hAut}_{\varphi}^{+,q}(\partial M))$$

is what makes it possible for us to obtain an example where the homotopy stable class $\mathcal{S}_{h, \lambda}^{\text{st}}$ is infinite. While an analogue of [Proposition 1.3](#) can be proved in the simply connected case using results of [\[Boyer 1993\]](#), the corresponding algebra always remains finite for trivial fundamental group.

All of the infinite sets we discuss are necessarily countable. Primarily, this has to be the case because there are only countably many compact manifolds [\[Cheeger and Kister 1970\]](#). On the algebraic side it is also evident that the orbit set onto which the homotopy stable class surjects in [Proposition 1.3](#) is countable, essentially because all the homology groups involved are finitely generated over $\mathbb{Z}[t^{\pm 1}]$.

Next we discuss a variation on [Proposition 1.3](#) that may be of independent interest. The surjection in [Proposition 1.3](#) can be improved to a bijection if we require the homotopy equivalences $N_1 \simeq N_2$ to restrict to homeomorphisms on the boundary; i.e. if we consider

$$\mathcal{S}_{h^+, \lambda}^{\text{st}, \partial}(M) := \frac{\{N \mid N \cong_{\text{st}} M, \lambda_N \cong \lambda_M\}}{\text{o.p. homotopy eq. that restricts to a homeo. on the boundary}}$$

and change the target accordingly, i.e. consider

$$\text{Aut}(\text{Bl}_{\partial M}, \mu_{\text{Bl}_{\partial M}}) / (\text{Aut}(\lambda_M) \times \text{Homeo}_{\varphi}^{+,q}(\partial M))$$

instead of

$$\text{Aut}(\text{Bl}_{\partial M}, \mu_{\text{Bl}_{\partial M}}) / (\text{Aut}(\lambda_M) \times \text{hAut}_{\varphi}^{+,q}(\partial M)).$$

In fact, the same result is obtained with

$$\mathcal{S}_{+, \lambda}^{\text{st}}(M) := \frac{\{N \mid N \cong_{\text{st}} M, \lambda_N \cong \lambda_M\}}{\text{o.p. homeomorphism}}.$$

Proposition 1.5. *If M is a spin 4-manifold with $\pi_1(M) = \mathbb{Z}$, ribbon boundary and nondegenerate equivariant intersection form λ_M , then there are bijections*

$$\mathcal{S}_{h^+, \lambda}^{\text{st}, \partial}(M) \xrightarrow{\cong} \text{Aut}(\text{Bl}_{\partial M}, \mu_{\text{Bl}_{\partial M}}) / (\text{Aut}(\lambda_M) \times \text{Homeo}_{\varphi}^{+,q}(\partial M)),$$

$$\mathcal{S}_{+, \lambda}^{\text{st}}(M) \xrightarrow{\cong} \text{Aut}(\text{Bl}_{\partial M}, \mu_{\text{Bl}_{\partial M}}) / (\text{Aut}(\lambda_M) \times \text{Homeo}_{\varphi}^{+,q}(\partial M)).$$

The bijections are induced by the map b that will be introduced in [Construction 5.1](#). For $M := X_{2q}(U) \natural(S^1 \times D^3)$, with q an odd prime, the sets above are countably infinite.

Proof. The surjectivity follows from the same argument that we will use in [Proposition 5.2](#). We prove injectivity. If $b(N_1) = b(N_2)$, then [[Conway, Piccirillo and Powell 2022](#), Theorem 1.1] shows that the manifolds N_1 and N_2 are orientation-preserving homeomorphic. Since the quotient with $\text{Homeo}_{\varphi}^{+,q}(\partial M)$ replaced by $\text{hAut}_{\varphi}^{+,q}(\partial M)$ is infinite, and since $\text{Homeo}_{\varphi}^{+,q}(\partial M) \subseteq \text{hAut}_{\varphi}^{+,q}(\partial M)$, it follows that the sets in the statement are infinite. \square

We now characterise $M := X_{2q}(U) \natural(S^1 \times D^3)$ within $\mathcal{S}_{+, \lambda}^{\text{st}}(M)$ in terms of the knottedness of the sphere $S_l^2 := \{\text{pt}\} \times S^2 \subseteq ((S^1 \times S^2) \setminus \text{Int}(D^3)) \subseteq \partial M$ and the connect sum sphere $S_c^2 \subseteq M$:

Theorem 1.6. *For $M = X_{2q}(U) \natural(S^1 \times D^3)$ and $N \in \mathcal{S}_{+, \lambda}^{\text{st}}(M)$, the following are equivalent:*

- (1) N is homeomorphic to M .
- (2) $S_l^2 \subseteq \partial N$ bounds a locally flat $D^3 \subseteq N$.
- (3) S_c^2 bounds a locally flat $D^3 \subseteq N$.

Proof. The implications (1) \Rightarrow (2) and (1) \Rightarrow (3) are immediate.

We prove the implication (2) \Rightarrow (1). Cut N along the D^3 with boundary S_l^2 to obtain a simply connected 4-manifold with boundary $L(2q, 1)$ and $H_2 = \mathbb{Z}$. Theorem 0.1 of [[Boyer 1986](#)] implies that such a manifold is homeomorphic to $X_{2q}(U)$. Glue back the $D^3 \times [0, 1]$ that we removed to recover N as M .

Finally, we prove the implication (3) \Rightarrow (1). Cut N open along the separating D^3 , resulting in a disjoint union of two 4-manifolds. The first is simply connected with $H_2 = \mathbb{Z}$ and boundary $L(2q, 1)$ and is therefore homeomorphic to $X_{2q}(U)$ [Boyer 1986, Theorem 0.1]. The second has $\pi_1 = \mathbb{Z}$, no H_2 and boundary $S^1 \times S^2$; it is thus homeomorphic to $S^1 \times D^3$ [Freedman and Quinn 1990, §11.6]. Glue back the $D^3 \times [0, 1]$ that we removed to recover N as M . \square

Organisation. In Sections 2 and 3, we review some facts about linking forms and in particular the Blanchfield form. In Section 4 we give a criterion that implies stable homeomorphism of 4-manifolds with fundamental group \mathbb{Z} and nonempty boundary. In Section 5, we prove Proposition 1.3. In Section 6 we show that for $M = X_{2q}(U) \natural (S^1 \times D^3)$, with q an odd prime, the set

$$\text{Aut}(\text{Bl}_{\partial M}, \mu_{\text{Bl}_{\partial M}}) / (\text{Aut}(\lambda_M) \times \text{hAut}_{\varphi}^{+,q}(\partial M))$$

is infinite.

Conventions. We work in the topological category unless otherwise stated. All manifolds are assumed to be compact, connected, based, and oriented. If a manifold has a nonempty boundary, then the basepoint is assumed to be in the boundary. For a 4-manifold M with fundamental group \mathbb{Z} , we fix an identification and write $\pi_1(M) = \mathbb{Z}$. We say that M is spin if M admits a spin structure compatible with the orientation. We write $p \mapsto \bar{p}$ for the involution on $\mathbb{Z}[t^{\pm 1}]$ induced by $t \mapsto t^{-1}$. Given a $\mathbb{Z}[t^{\pm 1}]$ -module H , we write \bar{H} for the $\mathbb{Z}[t^{\pm 1}]$ -module whose underlying abelian group is H but with module structure given by $p \cdot h = \bar{p}h$ for $h \in H$ and $p \in \mathbb{Z}[t^{\pm 1}]$. We write $H^* := \overline{\text{Hom}_{\mathbb{Z}[t^{\pm 1}]}(H, \mathbb{Z}[t^{\pm 1}])}$.

2. Linking forms and unions

Since a large part of this paper is concerned with the Blanchfield form and isometries thereof, we start by recalling terminology related to the underlying algebra. In Section 2.1 we recall symmetric and quadratic linking forms. In Section 2.2 we recall how a Hermitian form has a boundary which is a symmetric linking form, and the boundary of an even form has the additional structure of a quadratic refinement. In Section 2.3 we recall how isometries of these linking forms can be used to glue two linking forms together, and we show that the union of two even forms along an isometry of their boundary quadratic linking forms is again an even form.

2.1. Symmetric and quadratic linking forms. Everything in this subsection is the special case for $\mathbb{Z}[t^{\pm 1}]$ of a general theory for arbitrary rings with involution developed by Ranicki [1981, §3.4].

Definition 2.1. A *symmetric linking form* over $\mathbb{Z}[t^{\pm 1}]$ is a pair (T, ℓ) , where T is a torsion $\mathbb{Z}[t^{\pm 1}]$ -module, and $\ell : T \times T \rightarrow \mathbb{Q}(t)/\mathbb{Z}[t^{\pm 1}]$ is a Hermitian, sesquilinear, nonsingular pairing.

We write $S := \mathbb{Z}[t^{\pm 1}] \setminus \{0\}$, and set

$$\begin{aligned} Q^1(\mathbb{Q}(t)/\mathbb{Z}[t^{\pm 1}]) &:= \frac{\{b \in \mathbb{Q}(t) \mid b - \bar{b} \in \mathbb{Z}[t^{\pm 1}]\}}{\mathbb{Z}[t^{\pm 1}]}, \\ Q_1(\mathbb{Z}[t^{\pm 1}], S) &:= \frac{\{b \in \mathbb{Q}(t) \mid b = \bar{b}\}}{\{a + \bar{a} \mid a \in \mathbb{Z}[t^{\pm 1}]\}}, \\ Q^1(\mathbb{Z}[t^{\pm 1}], S) &:= \frac{\{b \in \mathbb{Q}(t) \mid b - \bar{b} = a - \bar{a} \text{ for some } a \in \mathbb{Z}[t^{\pm 1}]\}}{\mathbb{Z}[t^{\pm 1}]} \\ &\subseteq Q^1(\mathbb{Q}(t)/\mathbb{Z}[t^{\pm 1}]). \end{aligned}$$

For a symmetric linking form (T, ℓ) , we have that $\ell(x, x) \in Q^1(\mathbb{Q}(t)/\mathbb{Z}[t^{\pm 1}])$ for all $x \in T$. The symmetric linking form is called *even* if $\ell(x, x) \in Q^1(\mathbb{Z}[t^{\pm 1}], S)$ for all $x \in T$. We define a map

$$q : Q_1(\mathbb{Z}[t^{\pm 1}], S) \rightarrow Q^1(\mathbb{Z}[t^{\pm 1}], S), \quad [b] \mapsto [b].$$

Definition 2.2. A *quadratic refinement* of an even symmetric linking form (T, ℓ) is a function $\mu : T \rightarrow Q_1(\mathbb{Z}[t^{\pm 1}], S)$ satisfying

- (i) $\mu(rx) = r\mu(x)\bar{r} \in Q_1(\mathbb{Z}[t^{\pm 1}], S)$ for all $x \in T$ and for all $r \in \mathbb{Z}[t^{\pm 1}]$;
- (ii) $\mu(x + y) = \mu(x) + \mu(y) + \ell(x, y) + \overline{\ell(x, y)} \in Q_1(\mathbb{Z}[t^{\pm 1}], S)$ for all $x, y \in T$;
- (iii) $q(\mu(x)) = \ell(x, x) \in Q^1(\mathbb{Z}[t^{\pm 1}], S)$ for all $x \in T$.

A triple (T, ℓ, μ) consisting of a symmetric linking form together with a quadratic refinement is called a *quadratic linking form* over $\mathbb{Z}[t^{\pm 1}]$.

For aficionados of [Ranicki 1981], we emphasise that we are using the nonsplit version of quadratic linking forms.

We will also need to consider isometries and automorphisms of symmetric and quadratic linking forms.

Definition 2.3. Let (T, ℓ) and (T', ℓ') be symmetric linking forms over $\mathbb{Z}[t^{\pm 1}]$ and let $\mu : T \rightarrow Q_1(\mathbb{Z}[t^{\pm 1}], S)$ and $\mu' : T' \rightarrow Q_1(\mathbb{Z}[t^{\pm 1}], S)$ be respective quadratic refinements.

- (1) An isomorphism $f : T \rightarrow T'$ is an *isometry of symmetric linking forms* if

$$\ell'(f(x), f(y)) = \ell(x, y)$$

for every $x, y \in T$.

- (2) The isometry of symmetric linking forms f is moreover an *isometry of quadratic linking forms*, $f : (T, \ell, \mu) \cong (T', \ell', \mu')$ if $\mu'(f(x)) = \mu(x)$ for every $x \in T$.

- (3) If $(T, \ell) = (T', \ell')$, then f as in (1) is an *automorphism of symmetric linking forms*. We write $\text{Aut}(T, \ell)$ for the group of automorphisms.
- (4) If $(T, \ell, \mu) = (T', \ell', \mu')$, then f as in (2) is an *automorphism of quadratic linking forms*. We write $\text{Aut}(T, \ell, \mu)$ for the group of automorphisms.

Remark 2.4. Given a quadratic linking form (T, ℓ, μ) over $\mathbb{Z}[t^{\pm 1}]$ with underlying symmetric linking form (T, ℓ) , we note that $\text{Aut}(T, \ell, \mu) \subseteq \text{Aut}(T, \ell)$. We give an example showing that this can be a proper inclusion: multiplication by 3 induces an isomorphism $\mathbb{Z}[t^{\pm 1}]/8 \rightarrow \mathbb{Z}[t^{\pm 1}]/8$ that preserves the linking form $\ell(x, y) = \frac{1}{8}x\bar{y}$ but does not preserve the quadratic refinement $\mu(x) = \frac{1}{8}x\bar{x}$. Indeed $\mu(3) = \frac{9}{8} \neq \frac{1}{8} = \mu(1) \in \mathcal{Q}_1(\mathbb{Z}[t^{\pm 1}], S)$ because 1 cannot be written as $a + \bar{a}$ with $a \in \mathbb{Z}[t^{\pm 1}]$.

2.2. Boundaries of quadratic forms. We recall some terminology about Hermitian forms. A *Hermitian form* refers to a pair (H, λ) , where H is a free $\mathbb{Z}[t^{\pm 1}]$ -module and $\lambda : H \times H \rightarrow \mathbb{Z}[t^{\pm 1}]$ is a sesquilinear Hermitian pairing. Given a Hermitian form (H, λ) over $\mathbb{Z}[t^{\pm 1}]$, we use $\hat{\lambda} : H \rightarrow H^* := \text{Hom}_{\mathbb{Z}[t^{\pm 1}]}(H, \mathbb{Z}[t^{\pm 1}])$ to denote the linear map defined by $\hat{\lambda}(y)(x) = \lambda(x, y)$. We often refer to $\hat{\lambda}$ as the *adjoint* of λ . We say that λ is *nondegenerate* if $\hat{\lambda}$ is injective and *nonsingular* if $\hat{\lambda}$ is an isomorphism. We also recall that a Hermitian form (H, λ) is called *even* if for all $x \in H$, there exists $a \in \mathbb{Z}[t^{\pm 1}]$ such that $\lambda(x, x) = a + \bar{a}$.

We describe how a nondegenerate even Hermitian form over $\mathbb{Z}[t^{\pm 1}]$ determines a quadratic linking form, following [Ranicki 1981, p. 243].

Definition 2.5. The *boundary symmetric linking form* of a nondegenerate Hermitian form (H, λ) over $\mathbb{Z}[t^{\pm 1}]$ is the symmetric linking form $(\text{coker}(\hat{\lambda}), \partial\lambda)$, where $\partial\lambda$ is defined as

$$\partial\lambda : \text{coker}(\hat{\lambda}) \times \text{coker}(\hat{\lambda}) \rightarrow \mathbb{Q}(t)/\mathbb{Z}[t^{\pm 1}], \quad ([x], [y]) \mapsto \frac{1}{s}(y(z)),$$

where, as $\text{coker}(\hat{\lambda})$ is $\mathbb{Z}[t^{\pm 1}]$ -torsion, there exists $s \in \mathbb{Z}[t^{\pm 1}]$ and $z \in H$ such that $sx = \hat{\lambda}(z)$.

If (H, λ) is additionally assumed to be even, then its *boundary quadratic linking form* is the quadratic linking form $(\text{coker}(\hat{\lambda}), \partial\lambda, \mu_{\partial})$, where the quadratic refinement of $\partial\lambda$ is

$$\mu_{\partial} : \text{coker}(\hat{\lambda}) \rightarrow \mathcal{Q}_1(\mathbb{Z}[t^{\pm 1}], S), \quad [y] \mapsto \frac{1}{s}(y(z)),$$

with $s \in \mathbb{Z}[t^{\pm 1}]$ and $z \in H$ such that $sy = \hat{\lambda}(z)$.

We assert that $\partial\lambda$ is independent of the choices involved, and is nonsingular, sesquilinear, and Hermitian. These can all be verified directly. To enable us to give a reference to existing literature, note that the boundary symmetric linking

form is the linking form on $H^1(C^*)$ associated with the 1-dimensional $\mathbb{Q}(t)$ -acyclic symmetric Poincaré complex (C_*, φ) over $\mathbb{Z}[t^{\pm 1}]$ given by

$$\begin{array}{ccc} C^0 = H & \xrightarrow{\lambda} & C^1 = H^* \\ \downarrow \varphi_0 = \text{Id} & & \downarrow \varphi_0 = \text{Id} \\ C_1 = H & \xrightarrow{\lambda^* = \lambda} & C_0 = H^* \end{array}$$

In [Powell 2016, Propositions 3.3, 3.4, and 3.8] it was shown that such a linking form is well-defined, nonsingular, sesquilinear, and Hermitian.

The next proposition is implicit in [Ranicki 1981, p. 243]. As far as we know such a proof has not appeared in the literature, so we provide the details of the proof.

Proposition 2.6. *The function μ_{∂} is a well-defined function $\text{coker}(\hat{\lambda}) \rightarrow \mathcal{Q}_1(\mathbb{Z}[t^{\pm 1}], S)$, and is a quadratic refinement of the boundary symmetric linking form $\partial \lambda$ of (H, λ) , i.e. μ_{∂} satisfies the requirements of Definition 2.2.*

Proof. First we show it is well-defined. Let y, z , and s be as in Definition 2.5. Express $\hat{\lambda}$ as a Hermitian matrix $A = \bar{A}^T$ over $\mathbb{Z}[t^{\pm 1}]$ with respect to some basis of H . Then A is invertible over $\mathbb{Q}(t)$ because λ is nondegenerate, so $\det(A) \neq 0$. We have $\mu_{\partial}(y) = y^T A^{-1} \bar{y}$. Therefore

$$\overline{\mu_{\partial}(y)} = \overline{\mu_{\partial}(y)}^T = \overline{(y^T A^{-1} \bar{y})}^T = y^T \overline{A^{-1}}^T \bar{y} = y^T (\bar{A}^T)^{-1} \bar{y} = y^T A^{-1} \bar{y} = \mu_{\partial}(y).$$

Hence $\mu_{\partial}(y) \in \mathcal{Q}_1(\mathbb{Z}[t^{\pm 1}], S)$.

Next we show that the choices of z and s do not change $\mu_{\partial}(y)$. Let $y \in H$ and let $y \in \text{coker}(\hat{\lambda})$, $s \in \mathbb{Z}[t^{\pm 1}]$, and $z \in H$ be as in Definition 2.5, with $\hat{\lambda}(z) = sy$. Let $s' \in \mathbb{Z}[t^{\pm 1}]$ and $z' \in H$ be another pair of choices, such that $\hat{\lambda}(z') = s'y$. Since $\mu_{\partial}(y) \in \mathcal{Q}_1(\mathbb{Z}[t^{\pm 1}], S)$, we have

$$\frac{1}{s'} y(z') = \overline{\frac{1}{s'} y(z')}.$$

The difference between the two computations of $\mu_{\partial}(y)$ yields

$$\begin{aligned} \frac{1}{s} y(z) - \frac{1}{s'} y(z') &= \frac{1}{s} y(z) - \overline{\frac{1}{s'} y(z')} = \frac{1}{s} y(z) \frac{\bar{s}'}{s'} - \overline{\frac{1}{s'} y(z') \frac{\bar{s}}{s}} \\ &= \frac{1}{s} (s'y)(z) \frac{1}{s'} - \overline{\frac{1}{s'} (sy)(z') \frac{1}{s}} = \frac{1}{s} \hat{\lambda}(z')(z) \frac{1}{s'} - \overline{\frac{1}{s'} \hat{\lambda}(z)(z') \frac{1}{s}} \\ &= \frac{1}{s} \lambda(z, z') \frac{1}{s'} - \overline{\frac{1}{s'} \overline{\lambda(z, z')} \frac{1}{s}} = 0 \in \mathcal{Q}_1(\mathbb{Z}[t^{\pm 1}], S). \end{aligned}$$

Next we show that μ_{∂} does not depend on the representative y for the class in $\text{coker}(\hat{\lambda})$. Replace $y \in \text{coker}(\hat{\lambda})$ by another representative $y + \hat{\lambda}(u)$, for some $u \in H$. Then $\hat{\lambda}(z + su) = s(y + \hat{\lambda}(u))$. Therefore

$$\mu_{\partial}(y + \hat{\lambda}(u)) = \frac{1}{s} (y + \hat{\lambda}(u))(z + su) = \frac{1}{s} y(z) + \frac{1}{s} \hat{\lambda}(u)(z) + y(u) + \hat{\lambda}(u)(u).$$

We have

$$\hat{\lambda}(u)(z) = \lambda(z, u) = \overline{\lambda(u, z)} = \overline{\hat{\lambda}(z)(u)} = \overline{(sy)(u)} = \overline{y(u)\bar{s}} = s\overline{y(u)}.$$

Substituting, we obtain that

$$\mu_{\partial}(y + \hat{\lambda}(u)) = \frac{1}{s}y(z) + \overline{y(u)} + y(u) + \lambda(u, u).$$

The last term is symmetric over $\mathbb{Z}[t^{\pm 1}]$, which implies it is of the form $a + \bar{a}$. Hence up to terms of the form $a + \bar{a}$, we have

$$\mu_{\partial}(y + \hat{\lambda}(u)) = \frac{1}{s}y(z) = \mu_{\partial}(y),$$

as desired.

Now we know that μ_{∂} is well-defined, we prove that it satisfies the conditions in [Definition 2.2](#) for it to be a quadratic refinement of the boundary symmetric linking form $\partial\lambda$. For (i), let $y \in \text{coker}(\hat{\lambda})$, let $r \in \mathbb{Z}[t^{\pm 1}]$, and let $z \in H$ be such that $\hat{\lambda}(z) = sy$. Then $\hat{\lambda}(rz) = rsy = sry$. Thus

$$\mu_{\partial}(ry) = \frac{1}{s}((ry)(rz)) = \frac{r}{s}(y(z))\bar{r} = r\mu_{\partial}(y)\bar{r},$$

as desired. Next, aiming for (ii), we compute $\mu_{\partial}(x + y)$, for $x, y \in \text{coker}(\hat{\lambda})$. Let $r, s \in \mathbb{Z}[t^{\pm 1}]$ and $w, z \in H$ be such that $\hat{\lambda}(w) = rx$ and $\hat{\lambda}(z) = sy$. Then $\hat{\lambda}(sw + rz) = srw + rzy = rs(x + y)$. Hence

$$\begin{aligned} \mu_{\partial}(x + y) &= \frac{1}{rs}(x + y)(sw + rz) = \frac{1}{r}x(w) + \frac{1}{s}(y)(z) + \frac{1}{r}y(w) + \frac{1}{s}x(z) \\ &= \mu_{\partial}(x) + \mu_{\partial}(y) + \partial\lambda(x, y) + \partial\lambda(y, x). \end{aligned}$$

Since $\partial\lambda$ is Hermitian, this proves (ii). Condition (iii) is immediate from the formulae. \square

Remark 2.7. We note for later use that an isomorphism $F : H_0 \rightarrow H_1$ induces an isomorphism $F^{-*} := (F^*)^{-1} : H_0^* \rightarrow H_1^*$ and that if additionally the isomorphism F is an isometry, then F^{-*} descends to an isomorphism

$$\partial F := F^{-*} : \text{coker}(\hat{\lambda}_0) \rightarrow \text{coker}(\hat{\lambda}_1)$$

which determines an isometry of quadratic linking forms. Hence $\text{Aut}(\lambda)$ acts both on $\text{Aut}(\text{coker}(\hat{\lambda}), \partial\lambda)$ and on the subset $\text{Aut}(\text{coker}(\hat{\lambda}), \partial\lambda, \partial\mu_{\partial})$ by $F \cdot h = h \circ \partial F^{-1}$.

2.3. Algebraic unions. We recall the definition of the union of two Hermitian forms along an isometry of their boundary linking forms. The definition appears for the ring \mathbb{Z} in [\[Crowley 2002, Lemma 3.6\]](#) and was generalised to the ring $\mathbb{Z}[t^{\pm 1}]$ in [\[Conway and Powell 2023, Construction 2.7\]](#). The goal of this section is to prove that if the isometry preserves the quadratic refinements, then the union is an even form.

Construction 2.8. Let (H_0, λ_0) and (H_1, λ_1) be nondegenerate Hermitian forms over $\mathbb{Z}[t^{\pm 1}]$, and let $h : (\text{coker}(\hat{\lambda}_0), \partial\lambda_0) \rightarrow (\text{coker}(\hat{\lambda}_1), \partial\lambda_1)$ be an isometry of their boundary symmetric linking forms. Consider the pair $(H_0 \cup_h H_1, \lambda_0 \cup_h -\lambda_1)$ with

$$H_0 \cup_h H_1 := \ker(h\pi_0 - \pi_1 : H_0^* \oplus H_1^* \rightarrow \text{coker}(\hat{\lambda}_1))$$

$$\lambda_0 \cup_h -\lambda_1 \left(\begin{pmatrix} x_0 \\ x_1 \end{pmatrix}, \begin{pmatrix} y_0 \\ y_1 \end{pmatrix} \right) = \frac{1}{s_0} y_0(z_0) - \frac{1}{s_1} y_1(z_1) \in \mathbb{Q}(t),$$

where, since $\text{coker}(\hat{\lambda}_i)$ is torsion, there exists $s_i \in \mathbb{Z}[t^{\pm 1}]$ and $z_i \in H_i$ such that $s_i x_i = \hat{\lambda}_i(z_i)$. Since the Hermitian forms λ_0 and λ_1 are nondegenerate, it is not difficult to prove that the pairing $\lambda_0 \cup_h -\lambda_1$ does not depend on the choice of s_0, s_1, z_0, z_1 . One verifies that $\lambda_0 \cup_h -\lambda_1$ is a sesquilinear, Hermitian form and takes values in $\mathbb{Z}[t^{\pm 1}]$; see [Conway and Powell 2023, Proposition 2.8]. This pairing will be referred to as the *algebraic union* of λ_0 and λ_1 .

Lemma 2.9. Let (H_0, λ_0) , (H_1, λ_1) and (H, λ) be nondegenerate Hermitian forms over $\mathbb{Z}[t^{\pm 1}]$, and let $h : (\text{coker}(\hat{\lambda}_0), \partial\lambda_0) \rightarrow (\text{coker}(\hat{\lambda}_1), \partial\lambda_1)$ be an isometry of the boundary linking forms. If $F : \lambda_0 \cong \lambda_1$ is an isometry, then there is an isometry

$$\lambda_0 \cup_h -\lambda_1 \cong \lambda_1 \cup_{h \circ \partial F^{-1}} -\lambda_1.$$

Proof. See [Conway and Powell 2023, Proposition 2.8]. □

Lemma 2.10. Let (H_0, λ_0) and (H_1, λ_1) be two nondegenerate even Hermitian forms over $\mathbb{Z}[t^{\pm 1}]$. Suppose that $F : (H_0, \lambda_0) \cong (H_1, \lambda_1)$ is an isometry and that

$$h : (\text{coker}(\hat{\lambda}_0), \partial\lambda_0, (\mu_{\partial})_0) \rightarrow (\text{coker}(\hat{\lambda}_1), \partial\lambda_1, (\mu_{\partial})_1)$$

is an isometry of quadratic linking forms. Then the algebraic union $\lambda_0 \cup_h -\lambda_1$ is even.

Proof. Using Lemma 2.9 we can assume without loss of generality that $H_0 = H_1$ and $\lambda_0 = \lambda_1$ and $(\mu_{\partial})_0 = (\mu_{\partial})_1$. Write them both as $(H, \lambda, \mu_{\partial})$. This means composing h with ∂F , but since ∂F is an isometry of quadratic linking forms too, this does not affect the argument. We abuse notation and without loss of generality use h to denote the new isometry of boundary quadratic linking forms.

It thus suffices to check that for every $x_0, x_1 \in H^*$ such that $h \circ \pi_0(x_0) = \pi_1(x_1)$, the self-intersection $\lambda \cup_h -\lambda((x_0, x_1), (x_0, x_1))$ is of the form $a + \bar{a}$ for some $a \in \mathbb{Z}[t^{\pm 1}]$.

Pick $z_0, z_1 \in H$ and $s_0, s_1 \in \mathbb{Z}[t^{\pm 1}]$ such that $s_0 x_0 = \hat{\lambda}(z_0)$ and $s_1 x_1 = \hat{\lambda}(z_1)$. This implies both that

$$\mu_{\partial}(\pi_i(x_i)) = \frac{1}{s_i} (x_i(z_i)) \in \mathcal{Q}_1(\mathbb{Z}[t^{\pm 1}], S)$$

and that

$$\lambda \cup_h -\lambda((x_0, x_1), (x_0, x_1)) = \frac{1}{s_0} x_0(z_0) - \frac{1}{s_1} x_1(z_1) \in \mathbb{Z}[t^{\pm 1}].$$

Passing to $Q_1(\mathbb{Z}[t^{\pm 1}], S)$, by the definition of μ_{∂} this equals

$$\mu_{\partial}(\pi_0(x_0)) - \mu_{\partial}(\pi_1(x_1)) = \mu_{\partial}(\pi_0(x_0)) - \mu_{\partial}(h \circ \pi_0(x_0)) = 0.$$

The first equality used $h \circ \pi_0(x_0) = \pi_1(x_1)$. The second equality used that h is an isometry of quadratic linking forms.

Now we just have to note that the indeterminacy in $Q_1(\mathbb{Z}[t^{\pm 1}], S)$ consists entirely of even elements $\{a + \bar{a} \mid a \in \mathbb{Z}[t^{\pm 1}]\}$, and therefore $\lambda \cup_h -\lambda((x_0, x_1), (x_0, x_1)) \subseteq \{a + \bar{a} \mid a \in \mathbb{Z}[t^{\pm 1}]\}$. It follows that $\lambda \cup_h -\lambda$ is even, as desired. \square

3. The Blanchfield form

In [Section 3.1](#) we review the definition of the Blanchfield form and how it is related to the equivariant intersection form of a 4-manifold with fundamental group \mathbb{Z} . Then in [Section 3.2](#) we review isometries of Blanchfield forms. In [Section 3.3](#) we prove promised the spin gluing result, which demonstrates how isometries of the Blanchfield form together with a quadratic refinement can be used to ensure that the union of two spin 4-manifolds is again spin.

3.1. The Blanchfield form. We recall the definition of the Blanchfield form Bl_Y of a closed 3-manifold Y equipped with an epimorphism $\varphi : \pi_1(Y) \rightarrow \mathbb{Z}$ and how, if M is a 4-manifold with ribbon boundary, then $\text{Bl}_{\partial M}$ is related to the equivariant intersection form λ_M of M .

Construction 3.1. Let Y be a closed 3-manifold and let $\varphi : \pi_1(Y) \rightarrow \mathbb{Z}$ be an epimorphism. Assume that the Alexander module $H_1(Y; \mathbb{Z}[t^{\pm 1}])$ is torsion so that, in particular, the Bockstein homomorphism $\text{BS} : H^1(Y; \mathbb{Q}(t)/\mathbb{Z}[t^{\pm 1}]) \rightarrow H^2(Y; \mathbb{Z}[t^{\pm 1}])$ is an isomorphism. Consider the composition of Poincaré duality, the inverse Bockstein homomorphism and the evaluation homomorphism:

$$\begin{aligned} \Phi : H_1(Y; \mathbb{Z}[t^{\pm 1}]) &\xrightarrow{\text{PD}, \cong} H^2(Y; \mathbb{Z}[t^{\pm 1}]) \xrightarrow{\text{BS}^{-1}, \cong} H^1(Y; \mathbb{Q}(t)/\mathbb{Z}[t^{\pm 1}]) \\ &\xrightarrow{\text{ev}} \overline{\text{Hom}(H_1(Y; \mathbb{Z}[t^{\pm 1}]), \mathbb{Q}(t)/\mathbb{Z}[t^{\pm 1}])}. \end{aligned}$$

Using the universal coefficient spectral sequence, one can check that the evaluation map is an isomorphism, and thus so is Φ . Thus the pairing $(x, y) \mapsto \Phi(y)(x)$ is nonsingular. It is straightforward to see that this pairing is sesquilinear. It is also Hermitian; see e.g. [\[Powell 2016\]](#).

Definition 3.2. Let Y be a closed 3-manifold and let $\varphi : \pi_1(Y) \rightarrow \mathbb{Z}$ be an epimorphism such that the Alexander module $H_1(Y; \mathbb{Z}[t^{\pm 1}])$ is torsion. The *Blanchfield form*

$$\text{Bl}_Y : H_1(Y; \mathbb{Z}[t^{\pm 1}]) \times H_1(Y; \mathbb{Z}[t^{\pm 1}]) \rightarrow \mathbb{Q}(t)/\mathbb{Z}[t^{\pm 1}]$$

is the sesquilinear, nonsingular, Hermitian form defined by $\text{Bl}_Y(x, y) = \Phi(y)(x)$.

Let M be a 4-manifold with $\pi_1(M) = \mathbb{Z}$, nondegenerate equivariant intersection form λ_M and ribbon boundary (meaning that the inclusion induced map $\pi_1(\partial M) \rightarrow \pi_1(M)$ is surjective). We now outline why the symmetric boundary linking form $\partial\lambda_M$ (that was described in [Section 2.2](#)) is isometric to $-\text{Bl}_{\partial M}$.

As explained in [\[Conway and Powell 2023, Remark 3.3\]](#), the connecting homomorphism δ in the long exact sequence of the pair $(M, \partial M)$, together with Poincaré duality and the evaluation map, determines an isomorphism

$$D_M : \text{coker}(\hat{\lambda}_M) \xrightarrow{\cong} H_1(\partial M; \mathbb{Z}[t^{\pm 1}]), \quad [x] \mapsto \delta \circ \text{PD} \circ \text{ev}^{-1}(x)$$

that fits into the following commutative diagram:

$$(1) \quad \begin{array}{ccccccc} 0 & \longrightarrow & H_2(M; \mathbb{Z}[t^{\pm 1}]) & \xrightarrow{\hat{\lambda}_M} & H_2(M; \mathbb{Z}[t^{\pm 1}])^* & \longrightarrow & \text{coker}(\hat{\lambda}_M) \longrightarrow 0 \\ & & \downarrow = & & \downarrow \cong \text{PD} \circ \text{ev}^{-1} & & \downarrow \cong D_M \\ 0 & \longrightarrow & H_2(M; \mathbb{Z}[t^{\pm 1}]) & \xrightarrow{j} & H_2(M, \partial M; \mathbb{Z}[t^{\pm 1}]) & \xrightarrow{\delta} & H_1(\partial M; \mathbb{Z}[t^{\pm 1}]) \longrightarrow 0 \end{array}$$

Proposition 3.3 [\[Conway and Powell 2023, Proposition 3.5\]](#). *Let M be a 4-manifold with $\pi_1(M) = \mathbb{Z}$, whose boundary is ribbon and with torsion Alexander module. The isomorphism D_M induces an isometry of symmetric linking forms*

$$D_M : \partial(H_2(M; \mathbb{Z}[t^{\pm 1}]), \lambda_M) = (\text{coker}(\hat{\lambda}_M), \partial\lambda_M) \xrightarrow{\cong} (H_1(\partial M; \mathbb{Z}[t^{\pm 1}]), -\text{Bl}_{\partial M}).$$

Construction 3.4. Suppose M is a spin 4-manifold with $\pi_1(M) = \mathbb{Z}$, whose boundary is ribbon and with torsion Alexander module. Since M is spin, the equivariant intersection form of M is even. Then $\text{Bl}_{\partial M}$ admits a preferred quadratic refinement

$$\mu_{\text{Bl}_{\partial M}} : H_1(\partial M; \mathbb{Z}[t^{\pm 1}]) \rightarrow \mathcal{Q}_1(\mathbb{Z}[t^{\pm 1}], S), \quad x \mapsto \mu_{\partial}(D_M^{-1}(x)).$$

Here recall from [Definition 2.5](#) that μ_{∂} refers to the quadratic refinement of the symmetric linking form $\partial\lambda_M$; it exists because λ_M is even.

By construction

$$D_M : (\text{coker}(\hat{\lambda}_M), \partial\lambda_M, \mu_{\partial}) \xrightarrow{\cong} (H_1(\partial M; \mathbb{Z}[t^{\pm 1}]), -\text{Bl}_{\partial M}, -\mu_{\text{Bl}_{\partial M}})$$

is an isometry of quadratic linking forms.

3.2. Homotopy equivalences and isometries of the Blanchfield form. Given 3-manifolds Y_0, Y_1 equipped with epimorphisms $\varphi_i : \pi_1(Y_i) \rightarrow \mathbb{Z}$, we recall when homotopy equivalences of 3-manifolds induce isometries of the corresponding Blanchfield forms. We then apply these considerations to boundaries of 4-manifolds with $\pi_1 = \mathbb{Z}$.

Proposition 3.5 [\[Conway and Powell 2023, Proposition 3.7\]](#). *Let Y_0, Y_1 be 3-manifolds equipped with epimorphisms $\varphi_i : \pi_1(Y_i) \rightarrow \mathbb{Z}$ and assume that the resulting*

Alexander modules are torsion for $i = 0, 1$. If an orientation-preserving homotopy equivalence $f : Y_0 \rightarrow Y_1$ satisfies $\varphi_1 \circ f_* = \varphi_0$ on $\pi_1(Y_0)$, then it induces an isometry between the Blanchfield forms

$$\tilde{f}_* : H_1(Y_0; \mathbb{Z}[t^{\pm 1}]) \rightarrow H_1(Y_1; \mathbb{Z}[t^{\pm 1}]).$$

The proof of [Proposition 3.5](#) is fairly straightforward: the condition that

$$\varphi_1 \circ f_* = \varphi_0$$

ensures that f lifts to the infinite cyclic covers, and the required isometry is then obtained by taking the induced map on $H_1(-; \mathbb{Z}[t^{\pm 1}])$.

Remark 3.6. Let M and N be 4-manifolds with fundamental group \mathbb{Z} , whose boundaries are ribbon and with torsion Alexander modules.

- A consequence of [Proposition 3.5](#) is that an orientation-preserving homotopy equivalence $f : \partial M \rightarrow \partial N$ that intertwines the epimorphisms $\pi_1(\partial M) \twoheadrightarrow \pi_1(M)$ and $\pi_1(\partial N) \twoheadrightarrow \pi_1(N)$ induces an isometry $\tilde{f}_* : \text{Bl}_{\partial M} \cong \text{Bl}_{\partial N}$. However, in general \tilde{f}_* need not preserve the boundary quadratic refinements.
- Consider the group $\text{hAut}_{\varphi}^+(\partial M)$ of orientation-preserving homotopy equivalences $f : \partial M \rightarrow \partial M$ that satisfy $\varphi \circ f_* = f_* \circ \pi_1(\partial M) \rightarrow \mathbb{Z}$. We write

$$\text{hAut}_{\varphi}^{+,q}(\partial M) \subseteq \text{hAut}_{\varphi}^+(\partial M)$$

for the subset consisting of homotopy equivalences such that \tilde{f}_* preserves $\mu_{\text{Bl}_{\partial M}}$. [Proposition 3.5](#) implies that $\text{hAut}_{\varphi}^+(\partial M)$ acts on $\text{Aut}(\text{Bl}_{\partial M})$ and that $\text{hAut}_{\varphi}^{+,q}(\partial M)$ acts on $\text{Aut}(\text{Bl}_{\partial M}, \mu_{\text{Bl}_{\partial M}}) \subseteq \text{Aut}(\text{Bl}_{\partial M})$.

- In fact $\text{Aut}(\text{Bl}_{\partial M})$ and $\text{Aut}(\text{Bl}_{\partial M}, \mu_{\text{Bl}_{\partial M}})$ also admit actions of $\text{Aut}(\lambda_M)$. In more detail, we can use D_M to transport the action on $\text{Aut}(\partial\lambda_M)$ (recall [Remark 2.7](#)) to an action on $\text{Aut}(\text{Bl}_{\partial M})$: the action of F on $h \in \text{Aut}(\text{Bl}_{\partial M})$ is by $F \cdot h := h \circ (D_M \circ \partial F \circ D_M^{-1})$. Since ∂F also preserves μ_{∂} it follows that $\text{Aut}(\partial\lambda_M)$ also acts on $\text{Aut}(\text{Bl}_{\partial M}, \mu_{\text{Bl}_{\partial M}}) \subseteq \text{Aut}(\text{Bl}_{\partial M})$.
- [Remark 2.4](#) leads to an example where $\mu_{\text{Bl}_{\partial M}}$, the quadratic refinement of the Blanchfield form, depends on the choice of coboundary M . Consider the 4-manifold $M := (S^2 \widetilde{\times}_8 D^2) \natural (S^1 \times D^3)$, where $S^2 \widetilde{\times}_8 D^2$ denotes the total space of the D^2 -bundle over S^2 with Euler number 8. Multiplication by 3 induces an automorphism of the symmetric linking form of $\partial M = L(8, 1) \# (S^1 \times S^2)$ that is not induced by $\text{Aut}(\lambda_M)$ nor by $\text{Homeo}_{\varphi}^+(\partial M)$. Hence [\[Conway, Piccirillo and Powell 2022\]](#) implies there is a nonhomeomorphic 4-manifold M' with the same boundary and intersection form as M . The quadratic refinements of $\text{Bl}_{\partial M}$ induced by M and M' do not even lie in the same $\text{Homeo}_{\varphi}^+(\partial M)$ orbits, and so depend strongly on the choice of coboundary.

- It is an interesting question whether it is possible to define the quadratic refinement on $\text{Bl}_{\partial M}$ intrinsically, using only the spin structure M induces on ∂M . An analogue of this result is known for the standard linking form on ∂M ; see [Deloup and Massuyeau 2005, §2.4 & 2.5]. We leave this problem for later work.

3.3. Unions of spin 4-manifolds. We describe how the union construction from Section 2.3 and quadratic isometries of the Blanchfield form can be used to ensure that certain unions of spin 4-manifolds with fundamental group \mathbb{Z} remain spin.

In order to cut down on notation, we identify $(H_1(\partial M; \mathbb{Z}[t^{\pm 1}]), -\text{Bl}_{\partial M}, -\mu_{\text{Bl}_{\partial M}})$ with $(\text{coker}(\hat{\lambda}_M), \partial\lambda_M, \mu_\partial)$ i.e. we temporarily omit the isometry D_M mentioned in Proposition 3.3 from the notation. In particular, given an isometry $h : \text{Bl}_{\partial M} \cong \text{Bl}_{\partial N}$, we allow ourselves to write $\lambda_M \cup_h -\lambda_N$.

The next proposition recalls how the algebraic union can be used to understand the equivariant intersection form of a union of two 4-manifolds with fundamental group \mathbb{Z} .

Lemma 3.7 [Conway and Powell 2023, Proposition 3.9]. *Let M and N be two 4-manifolds with fundamental group \mathbb{Z} , nondegenerate equivariant intersection forms, and whose boundaries are ribbon. If there is an orientation-preserving homeomorphism $f : \partial M_0 \cong \partial M_1$ that intertwines the inclusion-induced epimorphisms $\pi_1(\partial M_0) \twoheadrightarrow \pi_1(M_0)$ and $\pi_1(\partial M_1) \twoheadrightarrow \pi_1(M_1)$, then there is an isometry*

$$\lambda_{M_0} \cup_{\tilde{f}_*} -\lambda_{M_1} \cong \lambda_{M_0 \cup_f - M_1}$$

Note that the intersection form being nondegenerate implies that the Alexander modules are $\mathbb{Z}[t^{\pm 1}]$ -torsion, via the long exact sequences of the pairs $(M, \partial M)$ and $(N, \partial N)$ with $\mathbb{Z}[t^{\pm 1}]$ coefficients.

Proposition 3.8. *Let M and N be spin 4-manifolds with fundamental group \mathbb{Z} , nondegenerate equivariant intersection forms, and whose boundaries are ribbon. If $\lambda_M \cong \lambda_N$ and $f : \partial M \cong \partial N$ is a homeomorphism that induces an isometry between the boundary quadratic linking forms of ∂M and ∂N , then $M \cup_f - N$ is spin.*

Proof. Pick an isometry $F : \lambda_M \cong \lambda_N$. Applying successively Lemma 3.7 and Lemma 2.9 we obtain

$$\lambda_{M \cup_f - N} \cong \lambda_M \cup_{\tilde{f}_*} -\lambda_N \cong \lambda_N \cup_{\tilde{f}_* \circ \partial F^{-1}} -\lambda_N.$$

Since \tilde{f}_* and ∂F preserve the quadratic linking forms, so does $\tilde{f}_* \circ \partial F^{-1}$.

As N is spin, it follows that λ_N is even. Lemma 2.10 implies that so is $\lambda_N \cup_{\tilde{f}_* \circ \partial F^{-1}} -\lambda_N$. Therefore $\lambda_{M \cup_f - N}$ is even. Since the fundamental group of $M \cup_f - N$ is \mathbb{Z} (see e.g. [Conway and Powell 2023, Proposition 3.8]) and therefore has no 2-torsion, this implies that $M \cup_f - N$ is spin, as required. \square

4. Stable homeomorphism

In this section, we collect some facts about stable homeomorphism of 4-manifolds with fundamental group \mathbb{Z} and nonempty boundary. We then focus on the case of 4-manifolds whose boundary is ribbon and has torsion Alexander module.

Given two 4-manifolds M and N with fundamental group $\pi_1(M) = \mathbb{Z} = \pi_1(N)$ that are either both spin or both nonspin, we call a homeomorphism $f : \partial M \xrightarrow{\cong} \partial N$ a *B-compatible* homeomorphism if the diagram

$$\begin{array}{ccc} \pi_1(\partial M) & \xrightarrow{f_*} & \pi_1(\partial N) \\ \downarrow & & \downarrow \\ \pi_1(M) & \xrightarrow{=} \mathbb{Z} \xleftarrow{=} & \pi_1(N) \end{array}$$

commutes, and if, in the case that M and N are spin, the union $M \cup_f -N$ is spin. We write $\partial M \cong_B \partial N$ if such a homeomorphism exists, and say that ∂M and ∂N are *B-homeomorphic*.

The terminology “*B-compatible*” and “*B-homeomorphic*” is motivated by our use of modified surgery below, where B is shorthand for the normal 1-type of the manifolds involved.

Lemma 4.1. *Let M and N be two 4-manifolds with fundamental group \mathbb{Z} and nonempty B -homeomorphic boundaries. Suppose that M and N have the same Kirby–Siebenmann invariant and isometric equivariant intersection forms. Then M and N are stably homeomorphic.*

In particular, there is an equality

$$\mathbb{S}_{h^+, \lambda}^{\text{st}}(M) = \frac{\{N \mid \partial N \cong_B \partial M, \pi_1(N) = \mathbb{Z}, \lambda_N \cong \lambda_M, \text{ks}(N) = \text{ks}(M)\}}{\text{o.p. homotopy eq. of pairs}}.$$

Proof. Theorem 2 of [Kreck 1999] ensures that M and N are stably homeomorphic if and only if there is a B -compatible homeomorphism $f : \partial M \xrightarrow{\cong} \partial N$ such that the union $M \cup_f -N$ vanishes in the bordism group $\Omega_4^{\text{STOP}}(S^1) \cong \mathbb{Z} \oplus \mathbb{Z}_2$ (when M and N are nonspin) or $\Omega_4^{\text{TOPSpin}}(S^1) \cong \mathbb{Z}$ (when M and N are spin). In the nonspin case, this bordism group is detected by the signature and Kirby–Siebenmann invariant whereas in the spin case, it is detected by the signature alone. Since $\lambda_M \cong \lambda_N$, we deduce that M and N have the same signature and Wall additivity implies that the union $M \cup_f -N$ has vanishing signature. The fact that $M \cup_f -N$ has vanishing Kirby–Siebenmann invariant follows from the additivity of this invariant; see e.g. [Friedl, Nagel, Orson and Powell 2019, Theorem 8.2]. \square

Given a 4-manifold M with $\pi_1(M) = \mathbb{Z}$, nondegenerate equivariant intersection form and ribbon boundary, as we will describe below, the methods of [Conway, Piccirillo and Powell 2022] produce 4-manifolds N with $\pi_1(N) = \mathbb{Z}$, ribbon boundary, $\lambda_M \cong \lambda_N$, $\text{ks}(M) = \text{ks}(N)$ and a preferred identification $g : \partial M \cong \partial N$. Since

$\lambda_M \cong \lambda_N$ and $\pi_1(M) = \mathbb{Z} = \pi_1(N)$ has no 2-torsion, either M and N are both spin or both nonspin. By [Lemma 4.1](#), in order to prove that M and N are stably homeomorphic, it therefore suffices to know that in the spin case, the union $M \cup_g N$ is spin, for which we use [Proposition 3.8](#).

Proposition 4.2. *Let M and N be spin 4-manifolds with fundamental group \mathbb{Z} , nondegenerate equivariant intersection forms, and whose boundaries are ribbon. If $\lambda_M \cong \lambda_N$, and $g : \partial M \cong \partial N$ is a homeomorphism that induces an isometry between the boundary quadratic linking forms of ∂M and ∂N , then M and N are stably homeomorphic.*

Proof. [Proposition 3.8](#) implies that the homeomorphism $\partial M \cong \partial N$ is B -compatible and the result therefore follows from [Lemma 4.1](#). \square

5. From stable homeomorphism to isometries of the Blanchfield form

In this section, M denotes a spin 4-manifold with an identification $\pi_1(M) = \mathbb{Z}$, ribbon boundary (the inclusion induced map $\varphi : \pi_1(\partial M) \rightarrow \pi_1(M)$ is surjective), and nondegenerate equivariant intersection form λ_M . As we recalled in [Section 3](#), since λ_M is nondegenerate, the Alexander module $H_1(\partial M; \mathbb{Z}[t^{\pm 1}])$ is torsion and supports the Blanchfield form,

$$\text{Bl}_{\partial M} : H_1(\partial M; \mathbb{Z}[t^{\pm 1}]) \times H_1(\partial M; \mathbb{Z}[t^{\pm 1}]) \rightarrow \mathbb{Q}(t)/\mathbb{Z}[t^{\pm 1}],$$

which is nonsingular, sesquilinear, and Hermitian. The goal of this section is to describe in more detail the set $\text{Aut}(\text{Bl}_{\partial M}, \mu_{\text{Bl}_{\partial M}})/(\text{Aut}(\lambda_M) \times \text{hAut}_{\varphi}^{+,q}(\partial M))$ that was mentioned in the introduction and then to prove [Proposition 1.3](#).

We start by recalling the aforementioned actions of $\text{hAut}_{\varphi}^{+,q}(\partial M)$ and $\text{Aut}(\lambda_M)$ on the group $\text{Aut}(\text{Bl}_{\partial M}, \mu_{\text{Bl}_{\partial M}})$ of isometries of the quadratic refinement of the Blanchfield form of ∂M induced by the even form λ_M . Here recall that $\text{hAut}_{\varphi}^{+,q}(\partial M)$ denotes the group of orientation-preserving homotopy equivalences $f : \partial M \rightarrow \partial M$ that satisfy $\varphi \circ f_* = f_* : \pi_1(\partial M) \rightarrow \mathbb{Z}$ and preserve $\mu_{\text{Bl}_{\partial M}}$, while $\text{Aut}(\lambda_M)$ denotes the set of isometries of λ_M .

- We recall the action of $\text{hAut}_{\varphi}^{+,q}(\partial M)$ on $\text{Aut}(\text{Bl}_{\partial M}, \mu_{\text{Bl}_{\partial M}})$. Since any homotopy equivalence $f \in \text{hAut}_{\varphi}^{+,q}(\partial M)$ satisfies $\varphi \circ f_* = f_*$, it lifts to a homotopy equivalence \tilde{f} on the \mathbb{Z} -covers that induces a $\mathbb{Z}[t^{\pm 1}]$ -linear map on homology; we denote this map by \tilde{f}_* . Since f is orientation-preserving, so is \tilde{f} and it follows that \tilde{f}_* is an isometry of the Blanchfield form. The action of f on $h \in \text{Aut}(\text{Bl}_{\partial M}, \mu_{\text{Bl}_{\partial M}})$ is then by $f \cdot h = \tilde{f}_* \circ h$.
- We describe the action of $\text{Aut}(\lambda_M)$ on $\text{Aut}(\text{Bl}_{\partial M}, \mu_{\text{Bl}_{\partial M}})$. Recall from [Section 3](#) that the even Hermitian form λ_M determines an adjoint map

$$\hat{\lambda}_M : H_2(M; \mathbb{Z}[t^{\pm 1}]) \rightarrow H_2(M; \mathbb{Z}[t^{\pm 1}])^*,$$

and a $\mathbb{Q}(t)/\mathbb{Z}[t^{\pm 1}]$ -valued quadratic linking form

$$(2) \quad \begin{aligned} \partial\lambda_M : \text{coker}(\hat{\lambda}_M) \times \text{coker}(\hat{\lambda}_M) &\rightarrow \mathbb{Q}(t)/\mathbb{Z}[t^{\pm 1}]; & \partial\lambda_M([x], [y]) &= y(z)/p, \\ \mu_\partial : \text{coker}(\hat{\lambda}_M) &\rightarrow \mathcal{Q}_+(\mathbb{Z}[t^{\pm 1}], S); & \mu_\partial([x]) &= x(z)/p. \end{aligned}$$

Here $p \in \mathbb{Z}[t^{\pm 1}]$ and $z \in H_2(M; \mathbb{Z}[t^{\pm 1}])$ satisfy $px = \hat{\lambda}_M(z)$. In [Section 3](#) we also recalled the definition of the isometry

$$D_M : -\partial\lambda_M \cong \text{Bl}_{\partial M},$$

and that an isometry $F \in \text{Aut}(\lambda_M)$ induces an isometries $\partial F : -\partial\lambda_M \cong -\text{Bl}_{\partial M}$ and $\partial F : \mu_\partial \cong -\mu_{\text{Bl}_{\partial M}}$ by noting that the isomorphism $(F^*)^{-1}$ descends to an isometry on the cokernels. The action of F on $h \in \text{Aut}(\text{Bl}_{\partial M}, \mu_{\text{Bl}_{\partial M}})$ is then by $F \cdot h := h \circ (D_M \circ \partial F \circ D_M^{-1})$.

Note that the actions commute, because one acts by precomposition and the other acts by postcomposition. We obtain an action of the product $\text{Aut}(\lambda_M) \times \text{hAut}_\varphi^{+,q}(\partial M)$ on $\text{Aut}(\text{Bl}_{\partial M}, \mu_{\text{Bl}_{\partial M}})$. Now that we have made sense of the orbit set

$$\text{Aut}(\text{Bl}_{\partial M}, \mu_{\text{Bl}_{\partial M}}) / (\text{Aut}(\lambda_M) \times \text{hAut}_\varphi^{+,q}(\partial M)),$$

we describe its relation to the homotopy stable class of M . Recall from [Lemma 4.1](#) that

$$(3) \quad \mathcal{S}_{h^+, \lambda}^{\text{st}}(M) = \frac{\{N \mid \partial N \cong_B \partial M, \pi_1(N) = \mathbb{Z}, \lambda_N \cong \lambda_M, \text{ks}(N) = \text{ks}(M)\}}{\text{o.p. hom. equiv. of pairs}}.$$

In order to relate $\mathcal{S}_{h^+, \lambda}^{\text{st}}(M)$ to the orbit set

$$\text{Aut}(\text{Bl}_{\partial M}, \mu_{\text{Bl}_{\partial M}}) / (\text{Aut}(\lambda_M) \times \text{hAut}_\varphi^{+,q}(\partial M)),$$

we recall a construction from [\[Conway, Piccirillo and Powell 2022, Construction 1\]](#) which has its origins in [\[Boyer 1993\]](#).

Construction 5.1. We describe a map

$$b : \mathcal{S}_{h^+, \lambda}^{\text{st}}(M) \rightarrow \text{Aut}(\text{Bl}_{\partial M}) / (\text{Aut}(\lambda_M) \times \text{hAut}_\varphi^+(\partial M)).$$

Given a 4-manifold $N \in \mathcal{S}_{h^+, \lambda}^{\text{st}}(M)$, pick a homeomorphism $g : \partial N \cong \partial M$ and an isometry $F : \lambda_M \cong \lambda_N$. Since $N \in \mathcal{S}_{h^+, \lambda}^{\text{st}}(M)$, it also has ribbon boundary and torsion Alexander module, thus ensuring that the isometry $D_N : -\partial\lambda_N \cong \text{Bl}_{\partial N}$ is defined. Now set

$$b(N) := g_* \circ D_N \circ \partial F \in \text{Aut}(\text{Bl}_{\partial M}) / (\text{Aut}(\lambda_M) \times \text{hAut}_\varphi^+(\partial M)).$$

One verifies that $b(N)$ is independent of the choices of F , g and the orientation-preserving homotopy equivalence class of $(N, \partial N)$. See [\[Conway, Piccirillo and Powell 2022\]](#) for details.

Suppose that in addition we assume the homeomorphism g is such that the induces homomorphism $g_* : H_1(\partial N; \mathbb{Z}[t^{\pm 1}]) \rightarrow H_1(\partial M; \mathbb{Z}[t^{\pm 1}])$ intertwines the quadratic refinements $\mu_{\text{Bl}_{\partial N}}$ and $\mu_{\text{Bl}_{\partial M}}$. To see that such a homeomorphism exists, restrict a stable homeomorphism $\Phi : M \# W_r \cong N \# W_r$ to the boundaries: this will intertwine the quadratic refinements because stabilising a Hermitian form by $\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$ does not affect the boundary quadratic linking form. Since both D_N and ∂F preserve the quadratic refinements, it follows that b in fact defines a map

$$b : \mathcal{S}_{h^+, \lambda}^{\text{st}}(M) \rightarrow \text{Aut}(\text{Bl}_{\partial M}, \mu_{\text{Bl}_{\partial M}}) / (\text{Aut}(\lambda_M) \times \text{hAut}_{\varphi}^{+,q}(\partial M)).$$

The next proposition proves [Proposition 1.3](#) from the introduction.

Proposition 5.2. *If M is a spin 4-manifold with $\pi_1(M) = \mathbb{Z}$, ribbon boundary and nondegenerate equivariant intersection form λ_M , then the map b from [Construction 5.1](#) defines a surjection*

$$b : \mathcal{S}_{h^+, \lambda}^{\text{st}}(M) \twoheadrightarrow \text{Aut}(\text{Bl}_{\partial M}, \mu_{\text{Bl}_{\partial M}}) / (\text{Aut}(\lambda_M) \times \text{hAut}_{\varphi}^{+,q}(\partial M)).$$

Proof. According to [\[Conway, Piccirillo and Powell 2022, Theorem 1.15\]](#), after fixing one choice of isometry $\partial \lambda_M \cong -\text{Bl}_{\partial M}$ (we will use D_M) every element $\text{Aut}(\text{Bl}_{\partial M}) / \text{Aut}(\lambda_M)$ is realised by a 4-manifold N with $\pi_1(N) = \mathbb{Z}$, ribbon boundary ∂N homeomorphic to ∂M via a homeomorphism $g : \partial N \cong \partial M$, equivariant intersection form λ_N isometric to λ_M via an isometry $F : \lambda_M \cong \lambda_N$, and $\text{ks}(N) = \text{ks}(M)$. In particular we can realise every element

$$b \in \text{Aut}(\text{Bl}_{\partial M}, \mu_{\text{Bl}_{\partial M}}) / (\text{Aut}(\lambda_M) \times \text{hAut}_{\varphi}^{+,q}(\partial M))$$

by such a manifold N .

As indicated in [\[Conway, Piccirillo and Powell 2022, Proof of Proposition 4.14\]](#), we have $b = g_* \circ D_N \circ \partial F$. Since b , D_N , and ∂F are isometries that preserve quadratic refinements, so does g .

[Proposition 4.2](#) now ensures that M and N are stably homeomorphic. We have therefore produced $N \in \mathcal{S}_{h^+, \lambda}^{\text{st}}(M)$ with $b(N) = b$, as required. \square

6. Infinite automorphism sets

In this section, we conclude the proof of [Theorem 1.1](#) by showing that

$$\text{Aut}(\text{Bl}_{\partial M}, \mu_{\text{Bl}_{\partial M}}) / (\text{Aut}(\lambda_M) \times \text{hAut}_{\varphi}^{+,q}(\partial M))$$

is countably infinite when

$$M = X_{2q}(U) \natural(S^1 \times D^3),$$

with q an odd prime. Here $X_{2q}(U)$ denotes the $2q$ -trace on the unknot U , i.e. the smooth 4-manifold obtained from D^4 by attaching a $2q$ -framed 2-handle along the

unknot. The plan is to first study $\text{Aut}(\text{Bl}_{\partial M}, \mu_{\text{Bl}_{\partial M}}) / \text{Aut}(\lambda_M)$ and then to consider the action by the self-homotopy equivalences of ∂M . In fact we will show in [Lemma 6.3](#) that $\text{Aut}(\text{Bl}_{\partial M}, \mu_{\text{Bl}_{\partial M}}) / \text{Aut}(\lambda_M) = \text{Aut}(\text{Bl}_{\partial M}) / \text{Aut}(\lambda_M)$. So first we will consider the latter set, ignoring quadratic refinements for a moment.

To study $\text{Aut}(\text{Bl}_{\partial M}) / \text{Aut}(\lambda_M)$, recall from [Section 5](#) that $\text{Bl}_{\partial M}$ is isometric to the linking form $\partial\lambda_M$ defined in (2), at the start of [Section 5](#). In particular, the isometry $D_M : -\partial\lambda_M \cong \text{Bl}_{\partial M}$ induces a bijection

$$\text{Aut}(\text{Bl}_{\partial M}) / \text{Aut}(\lambda_M) \cong \text{Aut}(\partial\lambda_M) / \text{Aut}(\lambda_M).$$

For rank one forms (such as $(H_2(M; \mathbb{Z}[t^{\pm 1}]), \lambda_M) = (\mathbb{Z}[t^{\pm 1}], \lambda_{2q})$), this set admits a particularly convenient description.

Given a ring R with involution $x \mapsto \bar{x}$, the group of *unitary units* $U(R)$ refers to those $u \in R$ such that $u\bar{u} = 1$, with the group operation given by restricting the multiplication on R . For example, when $R = \mathbb{Z}[t^{\pm 1}]$, all units are unitary and are of the form $\pm t^k$ with $k \in \mathbb{Z}$. In what follows, we make no distinction between rank one Hermitian forms and symmetric Laurent polynomials. The next lemma follows by unwinding the definition of $\text{Aut}(\partial\lambda)$; see also [[Conway and Powell 2023](#), Remark 1.16; [Conway, Piccirillo and Powell 2022](#), Lemma 7.1].

Lemma 6.1. *If $\lambda \in \mathbb{Z}[t^{\pm 1}]$ is a symmetric Laurent polynomial, then*

$$\text{Aut}(\partial\lambda) / \text{Aut}(\lambda) = U(\mathbb{Z}[t^{\pm 1}]/\lambda) / U(\mathbb{Z}[t^{\pm 1}]).$$

Proposition 6.2. *Given an odd prime q , the map*

$$\Theta : \mathbb{Z} \xrightarrow{\cong} U(\mathbb{Z}[t^{\pm 1}]/2q) / U(\mathbb{Z}[t^{\pm 1}]), \quad n \mapsto (q-1)t^n + q$$

is a group isomorphism.

Proof. One verifies that $(q-1)t^n + q$ is a unitary unit by using that $q(q-1) \equiv 0 \pmod{2q}$ (recall that q is odd). We then check that Θ is a homomorphism:

$$\begin{aligned} n + m \mapsto ((q-1)t^n + q)((q-1)t^m + q) &= (q-1)^2 t^{n+m} + q(q-1)(t^m + t^n) + q^2 \\ &\sim -(q-1)t^{n+m} - q \\ &\sim (q-1)t^{n+m} + q. \end{aligned}$$

Here the penultimate equivalence uses that $(q-1)^2 = q(q-1) - (q-1) \equiv -(q-1) \pmod{2q}$ and $q^2 \equiv q \equiv -q \pmod{2q}$. The last equivalence uses that $-1 \in U(\mathbb{Z}[t^{\pm 1}])$.

Next we show that Θ is injective. If $(q-1)t^n + q$ were trivial, we would have $(q-1)t^n + q = \pm t^k \in \mathbb{Z}[t^{\pm 1}]/2q$ for some k , but this is true only if $n = 0$.

Now we show that Θ is surjective. An explicit verification shows that the following map is an isomorphism:

$$U(\mathbb{Z}[t^{\pm 1}]/2) \times U(\mathbb{Z}[t^{\pm 1}]/q) \rightarrow U(\mathbb{Z}[t^{\pm 1}]/2q), \quad (a, b) \mapsto qa - (q-1)b.$$

To see this one should check that $(qa - (q-1)b)(q\bar{a} - (q-1)\bar{b}) \equiv 1$ when $a\bar{a} = 1 = b\bar{b}$, which implies that the map lands in the claimed target. The inverse is given by $x \mapsto ([x]_2, [x]_q)$, i.e. considering the coefficients modulo 2 and q respectively. Checking that this is the inverse homomorphism implies that the map is an isomorphism as asserted.

The units of $\mathbb{Z}[t^{\pm 1}]/2$ are of the form t^m for $m \in \mathbb{Z}$. On the other hand, since q is an odd prime, the unitary units of $\mathbb{Z}[t^{\pm 1}]/q$ are of the form $\pm t^n$ for $n \in \mathbb{Z}$. It follows that

$$U(\mathbb{Z}[t^{\pm 1}]/2q) \cong \{qt^m + (q-1)\varepsilon t^n \mid n, m \in \mathbb{Z}, \varepsilon \in \{\pm 1\}\}.$$

Passing to the quotient by $U(\mathbb{Z}[t^{\pm 1}])$ yields the required isomorphism, because once we can multiply by $\pm t^k$ for any $k \in \mathbb{Z}$, we have $qt^m + (q-1)\varepsilon t^n \sim (q-1)\varepsilon t^{n-m} + q$. Also

$$-(q-1)t^{n-m} + q \sim -(q-1)t^{n-m} - q \sim (q-1)t^{n-m} + q,$$

so we can ignore the ε , and every element of $U(\mathbb{Z}[t^{\pm 1}]/2q)$ is of the form $(q-1)t^k + q$ for some $k \in \mathbb{Z}$. So Θ is indeed surjective, which completes the proof that Θ is an isomorphism. \square

Lemma 6.3. *Given an odd prime $q \in \mathbb{Z}$, for the Hermitian form $\lambda = 2q \in \mathbb{Z}[t^{\pm 1}]$, one has*

$$\text{Aut}(\partial\lambda, \mu_{\partial}) / \text{Aut}(\lambda) = \text{Aut}(\partial\lambda) / \text{Aut}(\lambda).$$

Proof. The inclusion $\text{Aut}(\partial\lambda, \mu_{\partial}) / \text{Aut}(\lambda) \subseteq \text{Aut}(\partial\lambda) / \text{Aut}(\lambda)$ always holds. So, thanks to [Proposition 6.2](#), the lemma reduces to proving that for every $n \in \mathbb{Z}$ multiplication by $(q-1)t^n + q$ as a map $\mathbb{Z}[t^{\pm 1}]/2q \rightarrow \mathbb{Z}[t^{\pm 1}]/2q$ preserves the quadratic refinement

$$\mu_{\partial}(x) = \frac{1}{2q}x.$$

Writing $q = 2k + 1$, a direct calculation in $Q_1(\mathbb{Z}[t^{\pm 1}], S)$ now shows that

$$\begin{aligned} \frac{1}{2q}((q-1)t^n + q)((q-1)t^{-n} + q) &= \frac{1}{2q}((q-1)^2 + q^2) + \frac{1}{2q}(q(q-1)(t^n + t^{-n})) \\ &= \frac{1}{2q} + 2k + k(t^n + t^{-n}) \equiv \frac{1}{2q}. \end{aligned}$$

This concludes the proof of the lemma. \square

For q an odd prime, the combination of [Lemma 6.1](#), [Proposition 6.2](#), and [Lemma 6.3](#) implies that for $M = X_{2q}(U) \natural(S^1 \times D^3)$ we have

$$\text{Aut}(\text{Bl}_{\partial M}, \mu_{\text{Bl}_{\partial M}}) / \text{Aut}(\lambda_M) = \text{Aut}(\text{Bl}_{\partial M}) / \text{Aut}(\lambda_M) \cong \mathbb{Z}.$$

We now study the effect of factoring out by $\text{hAut}_{\varphi}^{+,q}(\partial M)$.

Remark 6.4. Our strategy will be to determine $\mathrm{hAut}_\varphi^+(\partial M)$ and show that the effect of its action on $\mathrm{Aut}(\mathrm{Bl}_{\partial M})/\mathrm{Aut}(\lambda_M)$ is given by multiplication by elements of the form $\pm t^n$, $n \in \mathbb{Z}$. This will automatically imply that $\mathrm{hAut}_\varphi^{+,q}(\partial M)$ also acts on $\mathrm{Aut}(\mathrm{Bl}_{\partial M}, \mu_{\mathrm{Bl}_{\partial M}})/\mathrm{Aut}(\lambda_M)$ by multiplication by elements of the form $\pm t^n$.

In fact we will make this argument in a slightly more general setting. Consider the 3-manifold $Y := N\#(S^1 \times S^2)$, where N is a 3-manifold with finite fundamental group. We fix an identification $H_1(S^1 \times S^2) = \mathbb{Z}$, an identification $\pi_1(Y) = \pi_1(N) * \mathbb{Z}$, and consider the finite abelian group $A := TH_1(Y) \cong H_1(N)$. Let $\varphi : \pi_1(Y) \rightarrow H_1(Y)/TH_1(Y) = \mathbb{Z}$ be the canonical projection onto the free part of $H_1(Y)$. In what follows, to distinguish $H_1(S^1 \times S^2) = \mathbb{Z}$ from the free \mathbb{Z} -factor of $\pi_1(Y) \cong \pi_1(N) * \mathbb{Z}$, we will exclusively write $H_1(S^1 \times S^2)$ as $\langle t \rangle$.

Summarising the notation, we have

$$\begin{aligned} A &:= TH_1(Y) \cong H_1(N), & \varphi : \pi_1(Y) &\twoheadrightarrow \langle t \rangle, \\ \theta : \pi_1(N) &\xrightarrow{\mathrm{ab}} H_1(\pi_1(N)) = H_1(N) = A. \end{aligned}$$

The example we have in mind is $Y_q = \partial M = L(2q, 1)\#(S^1 \times S^2)$, where $M = X_{2q}(U) \natural S^1 \times D^3$, so that $A \cong \mathbb{Z}/2q$ and $\varphi : \pi_1(Y_q) \twoheadrightarrow \langle t \rangle$ coincides with the inclusion induced map $\pi_1(\partial M) \twoheadrightarrow \pi_1(M) = \mathbb{Z}$.

Returning to the more general setting where $Y = N\#(S^1 \times S^2)$ with N a 3-manifold with $\pi_1(N)$ finite, the epimorphism $\varphi : \pi_1(Y) \twoheadrightarrow \langle t \rangle$ induces an infinite cyclic cover Y^∞ with

$$H_1(Y^\infty) \cong H_1(Y; \mathbb{Z}[t^{\pm 1}]) \cong H_1(\ker(\varphi)).$$

Our goal is now to describe the isomorphism type of this $\mathbb{Z}[t^{\pm 1}]$ -module (this is the content of [Construction 6.5](#) and [Lemma 6.6](#) below) and to then deduce the effect of the action of $\mathrm{hAut}_\varphi^+(Y)$ on $H_1(Y; \mathbb{Z}[t^{\pm 1}])$ in [Proposition 6.7](#).

In what follows, we write $A[t^{\pm 1}]$ for the abelian group of Laurent polynomials with coefficients in the finite abelian group A .

Construction 6.5. We construct a group homomorphism $\Psi : A[t^{\pm 1}] \rightarrow H_1(\ker(\varphi))$. Elements of $A[t^{\pm 1}]$ are of the form $\sum_i a_i t^i$ with $a_i \in A$. As the map $\varphi : \pi_1(N) * \mathbb{Z} \rightarrow \mathbb{Z}$ is surjective, we can write each t^i as $\varphi(g_i) = t^i$ for some $g_i \in \pi_1(N) * \mathbb{Z}$. The abelianisation $\theta : \pi_1(N) \rightarrow A = H_1(\pi_1(N))$ is also surjective, so we can write each $a \in A$ as $a = \theta(p)$ for some $p \in \pi_1(N)$. We can therefore write an element of $A[t^{\pm 1}]$ as $\sum_i \theta(p_i) \varphi(g_i)$. Since $p_i \in \pi_1(N)$, $g_i \in \pi_1(N) * \mathbb{Z}$ and $A \subseteq \ker(\varphi)$, we can consider the element $g_i p_i g_i^{-1}$ as an element of $\ker(\varphi) \subseteq \pi_1(N) * \mathbb{Z}$ and use $[g_i p_i g_i^{-1}] \in H_1(\ker(\varphi))$ to denote its image in the abelianisation. Define the map Ψ as

$$\Psi : A[t^{\pm 1}] \rightarrow H_1(\ker(\varphi)), \quad \sum_i a_i t^i = \sum_i \theta(p_i) \varphi(g_i) \mapsto \sum_i [g_i p_i g_i^{-1}].$$

We show that Ψ does not depend on the choice of the p_i and the g_i . First we argue that the definition of Ψ does not depend on the choice of the p_i . It suffices to show that if $\theta(p) = \theta(p')$, then $\Psi(\theta(p)\varphi(g)) = \Psi(\theta(p')\varphi(g))$ for every $g \in \pi_1(N) * \mathbb{Z}$. Since $\theta(p(p')^{-1}) = 0$, we know that pp'^{-1} lies in the commutator subgroup $\pi_1(N)^{(1)} = [\pi_1(N), \pi_1(N)]$. Therefore, since $\pi_1(N)^{(1)}$ is normal, $gpp'^{-1}g^{-1} = (gpg^{-1})(gp'^{-1}g^{-1}) \in \pi_1(N)^{(1)}$ for all $g \in \pi_1(N) * \mathbb{Z}$. Since $\pi_1(N) \subseteq \ker(\varphi)$, it follows that $\pi_1(N)^{(1)} \subseteq (\ker(\varphi))^{(1)}$, and therefore $(gpg^{-1})(gp'^{-1}g^{-1}) \in (\ker(\varphi))^{(1)}$, from which it follows $(gpg^{-1})(gp'^{-1}g^{-1}) = 0 \in H_1(\ker(\varphi))$. We deduce that $[gpg^{-1}] = [gp'g^{-1}] \in H_1(\ker(\varphi))$ and thus

$$\Psi(\theta(p)\varphi(g)) = [gpg^{-1}] = [gp'g^{-1}] = \Psi(\theta(p')\varphi(g)) \in H_1(\ker(\varphi)).$$

This proves that Ψ does not depend on the choice of the p_i .

Next, we argue that the definition of Ψ does not depend on the choice of the g_i . This time, it suffices to prove that if $\varphi(g) = \varphi(g')$ and $p \in \pi_1(N)$, then $\Psi(\theta(p)\varphi(g)) = \Psi(\theta(p)\varphi(g'))$. This latter equality holds if and only if $[gpg^{-1}g'p^{-1}g'^{-1}] = 0 \in H_1(\ker(\varphi))$, which in turn, by conjugating with g^{-1} , holds if and only if $[pg^{-1}g'p^{-1}g'^{-1}g] = 0 \in H_1(\ker(\varphi))$. But since $pg^{-1}g'p^{-1}g'^{-1}g$ is a commutator of p and $g^{-1}g'$, which both lie in $\ker(\varphi)$, we indeed obtain $[pg^{-1}g'p^{-1}g'^{-1}g] = 0 \in H_1(\ker(\varphi))$.

This concludes the verification that Ψ does not depend on any of the choices we made. One also verifies readily that Ψ is a group homomorphism. This completes [Construction 6.5](#).

As in [Construction 6.5](#), for each $h \in \langle t \rangle$, we fix a $g \in \pi_1(N) * \mathbb{Z}$ such that $\varphi(g) = h$. This choice will be used again in the next lemma which establishes that the map Ψ is an isomorphism.

Lemma 6.6. *The map $\Psi : A[t^{\pm 1}] \rightarrow H_1(\ker(\varphi))$ from [Construction 6.5](#) is an isomorphism.*

Proof. We construct an inverse $\Theta : H_1(\ker(\varphi)) \rightarrow A[t^{\pm 1}]$ to Ψ . A word $w \in \ker(\varphi) \subseteq \pi_1(N) * \mathbb{Z}$ representing an element of $H_1(\ker(\varphi))$ is a product of elements of $\pi_1(N)$ and \mathbb{Z} .

By introducing cancelling pairs of the type $g_i^{-1}g_i$ in between each occurrence of a $p'_k \in \pi_1(N)$ in w , we can arrange that for some elements $\tilde{g}_k \in \pi_1(N) * \mathbb{Z}$ and $p'_k \in \pi_1(N)$, the word w is of the form

$$w = \prod_k \tilde{g}_k p'_k \tilde{g}_k^{-1}.$$

Here it is crucial to use that $w \in \ker(\varphi)$. For example if $w = p'_1 n_1 p'_2 n_2 p'_3 n_3$, for $p_i \in \pi_1(N)$ and $n_j \in \mathbb{Z}$, then since $w \in \ker(\varphi)$ we know that $n_3 = (n_1 n_2)^{-1} = (n_2 n_1)^{-1}$. Therefore we can express w as $w = p'_1 n_1 p'_2 n_1^{-1} (n_1 n_2) p'_3 (n_1 n_2)^{-1}$.

As was mentioned before the lemma, we fixed a preferred $g_j \in \pi_1(N) * \mathbb{Z}$ with $\varphi(g_j) = \varphi(\tilde{g}_k)$. Arguing as in [Construction 6.5](#) (when we showed that the choice of the g_i is immaterial), up to commutators in $[\ker(\varphi), \ker(\varphi)]$, we can replace $\tilde{g}_k p'_k \tilde{g}_k^{-1}$ with $g_j p'_j g_j^{-1}$. Next, working in $H_1(\ker(\varphi)) = \ker(\varphi)_{ab}$ and collecting terms with the same conjugating element g_j , we obtain an element of the form $\sum_j [g_j p_j g_j^{-1}]$, where $p_j = \prod_{\{k|\varphi(\tilde{g}_k)=\varphi(g_j)\}} p'_k$. We can therefore define a map

$$\Theta : H_1(\ker(\varphi)) \rightarrow A[t^{\pm 1}], \quad [w] \mapsto \sum_j \theta(p_j) \varphi(g_j).$$

One checks that the map Θ is a homomorphism and is the inverse to Ψ . Thus Ψ is an isomorphism. \square

We are now able to describe the action of $\text{hAut}_\varphi^+(Y)$ on $H_1(Y; \mathbb{Z}[t^{\pm 1}])$.

Proposition 6.7. *Let $at^\ell \in H_1(Y; \mathbb{Z}[t^{\pm 1}]) \cong A[t^{\pm 1}]$. The action of $f \in \text{hAut}_\varphi^+(Y)$ sends $at^\ell \mapsto a't^{k+\ell}$, for some $k \in \mathbb{Z}$ and for some element $a' \in A$ having the same order as a .*

Proof. As in [Construction 6.5](#), we can represent any element of $A[t^{\pm 1}]$ as a sum of $\theta(p)\varphi(g)$, where $p \in \pi_1(N)$ and $g \in \pi_1(N) * \mathbb{Z}$. We will describe $f_*(\theta(p)\varphi(g))$.

In fact, since we have the commutative diagram of isomorphisms

$$\begin{array}{ccc} A[t^{\pm 1}] & \xrightarrow{\cong} & H_1(\ker(\varphi)) \\ f_* \downarrow \cong & & f_* \downarrow \cong \\ A[t^{\pm 1}] & \xrightarrow{\cong} & H_1(\ker(\varphi)) \end{array}$$

it is equivalent to describe $\Psi^{-1} \circ f_* \circ \Psi(\theta(p)\varphi(g))$. First, the definition of Ψ implies that $\Psi(\theta(p)\varphi(g)) = [gpg^{-1}] \in H_1(\ker(\varphi))$. Applying f_* , we then obtain $[f_*(g)f_*(p)f_*(g)^{-1}] \in H_1(\ker(\varphi))$.

But now, under an isomorphism $\pi_1(N) * \mathbb{Z} \cong \pi_1(N) * \mathbb{Z}$, every element of $\pi_1(N)$ is sent to an element of finite order, since $\pi_1(N)$ is finite. This implies that for every $p \in \pi_1(N)$, we have that $f_*(p) = hp'h^{-1} \in \pi_1(N) * \mathbb{Z}$ for some $p' \in \pi_1(N)$ and some $h \in \pi_1(N) * \mathbb{Z}$. This follows by considering the cyclic subgroup generated by $f_*(p)$ and applying the Kurosh subgroup theorem, which implies that a finite subgroup of a free product of nontrivial groups is a conjugate of a finite subgroup of one of the factors.

Next, since f_* is an isomorphism, $[f_*(p)] = [hp'h^{-1}]$ has the same order as $[p]$ in $H_1(\ker(\varphi))$. Since they are conjugate, in $\pi_1(N) * \mathbb{Z}$, we know that $hp'h^{-1}$ and p' have the same order. We claim that $[hp'h^{-1}]$ has the same order as $[p']$ in $H_1(\ker(\varphi))$.

To prove the claim, suppose that $\text{ord}([p']) = k$. Then $[(p')^k] = 0 \in H_1(\ker(\varphi))$, i.e. $(p')^k \in \ker(\varphi)^{(1)}$. Since $\ker(\varphi)$ is normal, for every $x \in \ker(\varphi)$ we have that

$hxh^{-1} \in \ker(\varphi)$, and therefore since $h[x, y]h^{-1} = [h x h^{-1}, h y h^{-1}]$, for every $z \in \ker(\varphi)^{(1)}$ we have that $h z h^{-1} \in \ker(\varphi)^{(1)}$. Thus $h(p')^k h^{-1} = (h p' h^{-1})^k \in \ker(\varphi)^{(1)}$, and therefore $\text{ord}([h p' h^{-1}]) \leq k = \text{ord}([p'])$. Since p' is also a conjugate of $h p' h^{-1}$, by symmetry we also have $\text{ord}([p']) \leq \text{ord}([h p' h^{-1}])$, and so we have equality. This completes the proof of the claim.

The claim implies that in $H_1(\ker(\varphi))$ we have

$$\text{ord}([p']) = \text{ord}([h p' h^{-1}]) = \text{ord}([f_*(p)]) = \text{ord}([p]).$$

Returning to the main arc of the proof, so far we have

$$f_* \circ \Psi(\theta(p)\varphi(g)) = [f_*(g) f_*(p) f_*(g)^{-1}] = [f_*(g) h p' h^{-1} f_*(g)^{-1}]$$

and it remains to apply Ψ^{-1} . The effect of Ψ^{-1} is $\theta(p')\varphi(h)\varphi(f_*(g)) \in A[t^{\pm 1}]$. Since $f \in \text{hAut}_\varphi^+(Y)$, we have $\varphi \circ f_* = \varphi$ and therefore

$$f_*(\theta(p)\varphi(g)) = \Psi^{-1} \circ f_* \circ \Psi(\theta(p)\varphi(g)) = \theta(p')\varphi(h)\varphi(g) \in A[t^{\pm 1}].$$

We can now calculate $f_*(a t^\ell)$. Pick $g \in \pi_1(N) * \mathbb{Z}$ and $p \in \pi_1(N)$ such that we have $\varphi(g) = t^\ell$ and $\theta(p) = a$. Now $f_*(a t^\ell) = f_*(\theta(p)\varphi(g)) = \theta(p')\varphi(h)t^\ell$, so the lemma follows by writing $\varphi(h) = t^k$ and $a' := \theta(p')$. Then since $[p']$ has the same order as $[p]$, it follows that a' has the same order as a . \square

We can now prove the main result of this section.

Proposition 6.8. *Fix an odd prime q . For $M = X_{2q}(U) \natural(S^1 \times D^3)$, the sets*

$$\begin{aligned} & \text{Aut}(\text{Bl}_{\partial M}) / (\text{Aut}(\lambda_M) \times \text{hAut}_\varphi^+(\partial M)), \\ & \text{Aut}(\text{Bl}_{\partial M}, \mu_{\text{Bl}_{\partial M}}) / (\text{Aut}(\lambda_M) \times \text{hAut}_\varphi^{+,q}(\partial M)) \end{aligned}$$

are countably infinite.

Proof. Fix identifications $\pi_1(M) = \mathbb{Z}$ and $(H_2(M; \mathbb{Z}[t^{\pm 1}]), \lambda_M) = (\mathbb{Z}[t^{\pm 1}], \lambda_{2q})$. [Lemma 6.1](#) implies that $\text{Aut}(\text{Bl}_{\partial M}) / \text{Aut}(\lambda_M) = U(\mathbb{Z}[t^{\pm 1}]/2q) / U(\mathbb{Z}[t^{\pm 1}])$. We know from [Proposition 6.2](#) that $U(\mathbb{Z}[t^{\pm 1}]/2q) / U(\mathbb{Z}[t^{\pm 1}]) \cong \mathbb{Z}$, every element of which is of the form $(q-1)t^n + q$ with $n \in \mathbb{Z}$. We will now show that there is a bijection of sets

$$\text{Aut}(\text{Bl}_{\partial M}) / (\text{Aut}(\lambda_M) \times \text{hAut}_\varphi^+(\partial M)) \cong \mathbb{Z}.$$

In the notation of [Proposition 6.7](#), we have $N = L(2q, 1)$ with $\pi_1(L(2q, 1)) \cong \mathbb{Z}/2q$ as well as $A = H_1(\pi_1(N)) = \pi_1(N) = \mathbb{Z}/2q$.

Using [Proposition 6.7](#), we will argue that any automorphism of the group $H_1(\partial M; \mathbb{Z}[t^{\pm 1}]) \cong (\mathbb{Z}/2q)[t^{\pm 1}]$ induced by a homotopy equivalence $f \in \text{Aut}_\varphi^+(\partial M)$ is of the form $p(t) \mapsto \pm t^k p(t)$, for some $k \in \mathbb{Z}$. To see this, given $p(t) \in (\mathbb{Z}/2q)[t^{\pm 1}]$, by $\mathbb{Z}[t^{\pm 1}]$ -linearity of f_* we have $f_*(p(t)) = p(t) f_*(1)$. By [Proposition 6.7](#),

$f_*(1) = a \cdot t^k$, for some $k \in \mathbb{Z}$ and some $a \in \mathbb{Z}/2q$. We need to show that $a = \pm 1$. Since f_* is an isometry of $\text{Bl}_{\partial M}$, we also know that $a^2 = 1 \in \mathbb{Z}[t^{\pm 1}]/2q$; this holds because

$$\begin{aligned} \frac{-1}{2q} &= \text{Bl}_{\partial M}(1, 1) = \text{Bl}_{\partial M}(f_*(1), f_*(1)) \\ &= \text{Bl}_{\partial M}(a \cdot t^k, a \cdot t^k) = \frac{-a^2}{2q} \in \mathbb{Q}(t)/\mathbb{Z}[t^{\pm 1}], \end{aligned}$$

which implies that $a^2 = 1 \in \mathbb{Z}[t^{\pm 1}]/2q$. Then since $a \in \mathbb{Z}/2q$ we have that $a^2 = 1 \in \mathbb{Z}/2q$. Here we used that $\text{Bl}_{\partial M} \cong -\partial\lambda_{2q}$ to compute the Blanchfield form [Conway and Powell 2023, Proposition 3.5].

However the only elements of $A = \mathbb{Z}/2q$ with $a^2 = 1$ are $\pm 1 \in \mathbb{Z}/2q$. Indeed such an a belongs to $U(\mathbb{Z}/2q) \cong U(\mathbb{Z}/q) \times U(\mathbb{Z}/2)$. However $U(\mathbb{Z}/2)$ is trivial, so in fact $U(\mathbb{Z}/2q) \cong U(\mathbb{Z}/q)$. We will show that $U(\mathbb{Z}/q) = \{\pm 1\}$. To see this, recall that for q an odd prime the units $(\mathbb{Z}/q)^\times$ is a cyclic group of order $q - 1$, and in such a group there is precisely one element of order 2. Taken together with the trivial element there are therefore precisely two solutions to $x^2 = 1 \in (\mathbb{Z}/q)^\times$, namely ± 1 . So we see that $U(\mathbb{Z}/2q) \cong U(\mathbb{Z}/q) = \{\pm 1\}$. It follows that $a = \pm 1$ and

$$f_*(p(t)) = p(t)f_*(1) = \pm t^k p(t),$$

as asserted above. In particular, observe that the action of a homotopy equivalence $f \in \text{hAut}_\varphi^+(\partial M)$ is the same as the action by an element of $\text{Aut}(\lambda_M) \cong U(\mathbb{Z}[t^{\pm 1}])$. We deduce that

$$\text{Aut}(\text{Bl}_{\partial M})/(\text{Aut}(\lambda_M) \times \text{hAut}_\varphi^+(\partial M)) \cong \text{Aut}(\text{Bl}_{\partial M})/\text{Aut}(\lambda_M).$$

But in Proposition 6.2 we computed the latter set to be

$$\text{Aut}(\text{Bl}_{\partial M})/\text{Aut}(\lambda_M) \cong U(\mathbb{Z}[t^{\pm 1}]/2q)/U(\mathbb{Z}[t^{\pm 1}]) \cong \mathbb{Z}.$$

The inverse of these isomorphisms sends $n \in \mathbb{Z}$ to the automorphism given by multiplying by $(q-1)t^n + q$. Since the action of an element of $\text{hAut}_\varphi^+(\partial M)$ is the same as the action by an element $\text{Aut}(\lambda_M) \cong U(\mathbb{Z}[t^{\pm 1}])$, the same can be said for elements of $\text{hAut}_\varphi^{+,q}(\partial M) \subseteq \text{hAut}_\varphi^+(\partial M)$. As we mentioned in Remark 6.4, Lemma 6.3 now implies that

$$\text{Aut}(\text{Bl}_{\partial M}, \mu_{\text{Bl}_{\partial M}})/(\text{Aut}(\lambda_M) \times \text{hAut}_\varphi^{+,q}(\partial M)) \cong \text{Aut}(\text{Bl}_{\partial M})/(\text{Aut}(\lambda_M) \times \text{hAut}_\varphi^+(\partial M)).$$

The second assertion in Proposition 6.8 therefore follows from the first. \square

Acknowledgements

We thank the referee for helpful comments which helped improve the paper. Conway was partially supported by the NSF Standard Grant DMS-2303674. Powell was

partially supported by the EPSRC New Investigator grant EP/T028335/2 and EPSRC New Horizons grant EP/V04821X/2.

References

- [Boyer 1986] S. Boyer, “Simply-connected 4-manifolds with a given boundary”, *Trans. Amer. Math. Soc.* **298**:1 (1986), 331–357. [MR](#) [Zbl](#)
- [Boyer 1993] S. Boyer, “Realization of simply-connected 4-manifolds with a given boundary”, *Comment. Math. Helv.* **68**:1 (1993), 20–47. [MR](#) [Zbl](#)
- [Cheeger and Kister 1970] J. Cheeger and J. M. Kister, “Counting topological manifolds”, *Topology* **9** (1970), 149–151. [MR](#) [Zbl](#)
- [Conway and Powell 2023] A. Conway and M. Powell, “Embedded surfaces with infinite cyclic knot group”, *Geom. Topol.* **27**:2 (2023), 739–821. [MR](#) [Zbl](#)
- [Conway, Crowley, Powell and Sixt 2021] A. Conway, D. Crowley, M. Powell, and J. Sixt, “Stably diffeomorphic manifolds and modified surgery obstructions”, 2021. [arXiv 2109.05632](#)
- [Conway, Crowley, Powell and Sixt 2023] A. Conway, D. Crowley, M. Powell, and J. Sixt, “Simply connected manifolds with large homotopy stable classes”, *J. Aust. Math. Soc.* **115**:2 (2023), 172–203. [MR](#) [Zbl](#)
- [Conway, Piccirillo and Powell 2022] A. Conway, L. Piccirillo, and M. Powell, “4-manifolds with boundary and fundamental group \mathbb{Z} ”, 2022. [arXiv 2205.12774](#)
- [Crowley 2002] D. Crowley, *The classification of highly connected manifolds in dimensions 7 and 15*, Ph.D. thesis, Indiana University, 2002, available at <https://www.proquest.com/docview/305508775>.
- [Deloup and Massuyeau 2005] F. Deloup and G. Massuyeau, “Quadratic functions and complex spin structures on three-manifolds”, *Topology* **44**:3 (2005), 509–555. [MR](#) [Zbl](#)
- [Freedman 1982] M. H. Freedman, “The topology of four-dimensional manifolds”, *J. Differential Geometry* **17**:3 (1982), 357–453. [MR](#) [Zbl](#)
- [Freedman and Quinn 1990] M. H. Freedman and F. Quinn, *Topology of 4-manifolds*, Princeton Mathematical Series **39**, Princeton University Press, 1990. [MR](#) [Zbl](#)
- [Friedl, Nagel, Orson and Powell 2019] S. Friedl, M. Nagel, P. Orson, and M. Powell, “A survey of the foundations of four-manifold theory in the topological category”, 2019. [arXiv 1910.07372](#)
- [Hambleton and Kreck 1993] I. Hambleton and M. Kreck, “Cancellation, elliptic surfaces and the topology of certain four-manifolds”, *J. Reine Angew. Math.* **444** (1993), 79–100. [MR](#) [Zbl](#)
- [Hambleton, Kreck and Teichner 2009] I. Hambleton, M. Kreck, and P. Teichner, “Topological 4-manifolds with geometrically two-dimensional fundamental groups”, *J. Topol. Anal.* **1**:2 (2009), 123–151. [MR](#) [Zbl](#)
- [Kreck 1999] M. Kreck, “Surgery and duality”, *Ann. of Math. (2)* **149**:3 (1999), 707–754. [MR](#) [Zbl](#)
- [Kreck and Schafer 1984] M. Kreck and J. A. Schafer, “Classification and stable classification of manifolds: some examples”, *Comment. Math. Helv.* **59**:1 (1984), 12–38. [MR](#) [Zbl](#)
- [Powell 2016] M. Powell, “Twisted Blanchfield pairings and symmetric chain complexes”, *Q. J. Math.* **67**:4 (2016), 715–742. [MR](#) [Zbl](#)
- [Ranicki 1981] A. Ranicki, *Exact sequences in the algebraic theory of surgery*, Mathematical Notes **26**, Princeton University Press, 1981. [MR](#) [Zbl](#)
- [Wall 1999] C. T. C. Wall, *Surgery on compact manifolds*, 2nd ed., Mathematical Surveys and Monographs **69**, American Mathematical Society, Providence, RI, 1999. [MR](#) [Zbl](#)

Received October 7, 2022. Revised September 18, 2023.

ANTHONY CONWAY
DEPARTMENT OF MATHEMATICS
UNIVERSITY OF TEXAS AT AUSTIN
AUSTIN, TX
UNITED STATES
anthony.conway@austin.utexas.edu

DIARMUID CROWLEY
SCHOOL OF MATHEMATICS & STATISTICS
UNIVERSITY OF MELBOURNE
PARKVILLE, VIC
AUSTRALIA
dcrowley@unimelb.edu.au

MARK POWELL
SCHOOL OF MATHEMATICS AND STATISTICS
UNIVERSITY OF GLASGOW
GLASGOW
UNITED KINGDOM
mark.powell@glasgow.ac.uk

PACIFIC JOURNAL OF MATHEMATICS

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

msp.org/pjm

EDITORS

Don Blasius (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
blasius@math.ucla.edu

Matthias Aschenbrenner
Fakultät für Mathematik
Universität Wien
Vienna, Austria
matthias.aschenbrenner@univie.ac.at

Paul Balmer
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
balmer@math.ucla.edu

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Atsushi Ichino
Department of Mathematics
Kyoto University
Kyoto 606-8502, Japan
atsushi.ichino@gmail.com

Robert Lipshitz
Department of Mathematics
University of Oregon
Eugene, OR 97403
lipshitz@uoregon.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Dimitri Shlyakhtenko
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
shlyakht@ipam.ucla.edu

Paul Yang
Department of Mathematics
Princeton University
Princeton NJ 08544-1000
yang@math.princeton.edu

Ruixiang Zhang
Department of Mathematics
University of California
Berkeley, CA 94720-3840
ruixiang@berkeley.edu

PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org

See inside back cover or msp.org/pjm for submission instructions.

The subscription price for 2023 is US \$605/year for the electronic version, and \$820/year for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by [Mathematical Reviews](#), [Zentralblatt MATH](#), [PASCAL CNRS Index](#), [Referativnyi Zhurnal](#), [Current Mathematical Publications](#) and [Web of Knowledge \(Science Citation Index\)](#).

The Pacific Journal of Mathematics (ISSN 1945-5844 electronic, 0030-8730 printed) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFLOW® from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2023 Mathematical Sciences Publishers

PACIFIC JOURNAL OF MATHEMATICS

Volume 325 No. 2 August 2023

Totally geodesic hyperbolic 3-manifolds in hyperbolic link complements of tori in S^4	191
MICHELLE CHU and ALAN W. REID	
Infinite homotopy stable class for 4-manifolds with boundary	209
ANTHONY CONWAY, DIARMUID CROWLEY and MARK POWELL	
Counterexamples to the nonsimply connected double soul conjecture	239
JASON DEVITO	
The fundamental group of an extension in a Tannakian category and the unipotent radical of the Mumford–Tate group of an open curve	255
PAYMAN ESKANDARI and V. KUMAR MURTY	
The size of semigroup orbits modulo primes	281
WADE HINDES and JOSEPH H. SILVERMAN	
Tropical Lagrangian multisections and toric vector bundles	299
YAT-HIN SUEN	
Some effectivity results for primitive divisors of elliptic divisibility sequences	331
MATTEO VERZOBIO	
Poisson manifolds of strong compact type over 2-tori	353
LUKA ZWAAN	