

# Mathematics 2A—Multivariate Calculus (2013/14)

D. Bourne

November 7, 2013

## Teaching arrangements

- ▶ This section, 2A<sub>1</sub>, meets at 9 a.m. on Tuesday in room 513 Boyd Orr and 9am on Thursday in room 203 Mathematics Building

## Teaching arrangements

- ▶ This section, 2A<sub>1</sub>, meets at 9 a.m. on Tuesday in room 513 Boyd Orr and 9am on Thursday in room 203 Mathematics Building
- ▶ Other sections meets at 10am and 11am.

## Teaching arrangements

- ▶ This section, 2A<sub>1</sub>, meets at 9 a.m. on Tuesday in room 513 Boyd Orr and 9am on Thursday in room 203 Mathematics Building
- ▶ Other sections meets at 10am and 11am.
- ▶ **This week:** Lectures on Tuesday and Thursday  
- no tutorial this week

## Teaching arrangements

- ▶ This section, 2A<sub>1</sub>, meets at 9 a.m. on Tuesday in room 513 Boyd Orr and 9am on Thursday in room 203 Mathematics Building
- ▶ Other sections meets at 10am and 11am.
- ▶ **This week:** Lectures on Tuesday and Thursday  
- no tutorial this week
- ▶ **Other weeks:** Lectures on Tuesday and Thursday and a tutorial on Monday  
- students come to tutorials *every other week*. Go to MyCampus for information on which tutorial group you are in and which weeks you have a tutorial.

## Teaching arrangements

- ▶ This section, 2A<sub>3</sub>, always meets at 11 a.m. on Tuesday in Lecture Theatre 208 Sir Alexander Stone Building and 11am on Thursday in room 224 (Main Lecture Theatre) Graham Kerr Building.

## Teaching arrangements

- ▶ This section, 2A<sub>3</sub>, always meets at 11 a.m. on Tuesday in Lecture Theatre 208 Sir Alexander Stone Building and 11am on Thursday in room 224 (Main Lecture Theatre) Graham Kerr Building.
- ▶ Other section meets at 9am and 10am.

## Teaching arrangements

- ▶ This section, 2A<sub>3</sub>, always meets at 11 a.m. on Tuesday in Lecture Theatre 208 Sir Alexander Stone Building and 11am on Thursday in room 224 (Main Lecture Theatre) Graham Kerr Building.
- ▶ Other section meets at 9am and 10am.
- ▶ **This week:** Lectures on Tuesday and Thursday  
- no tutorial this week



## Teaching arrangements

- ▶ This section, 2A<sub>3</sub>, always meets at 11 a.m. on Tuesday in Lecture Theatre 208 Sir Alexander Stone Building and 11am on Thursday in room 224 (Main Lecture Theatre) Graham Kerr Building.
- ▶ Other section meets at 9am and 10am.
- ▶ **This week:** Lectures on Tuesday and Thursday  
- no tutorial this week
- ▶ **Other weeks:** Lectures on Tuesday and Thursday and a tutorial on Monday  
- students come to tutorials *every other week*. Go to MyCampus for information on which tutorial group you are in and which weeks you have a tutorial.

# Tutorials

- ▶ Tutorial questions on the exercise sheets should be done each week before the tutorial

# Tutorials

- ▶ Tutorial questions on the exercise sheets should be done each week before the tutorial
- ▶ Do this work even if you don't have a tutorial that week

# Tutorials

- ▶ Tutorial questions on the exercise sheets should be done each week before the tutorial
- ▶ Do this work even if you don't have a tutorial that week
- ▶ Tutors will ask to see your tutorial work so make sure you have your attempts with you. If you couldn't do a question make sure to have a note written down about why you couldn't do the question.

# Tutorials

- ▶ Tutorial questions on the exercise sheets should be done each week before the tutorial
- ▶ Do this work even if you don't have a tutorial that week
- ▶ Tutors will ask to see your tutorial work so make sure you have your attempts with you. If you couldn't do a question make sure to have a note written down about why you couldn't do the question.
- ▶ Seek help with these (and other problems) from tutors

# Tutorials

- ▶ Tutorial questions on the exercise sheets should be done each week before the tutorial
- ▶ Do this work even if you don't have a tutorial that week
- ▶ Tutors will ask to see your tutorial work so make sure you have your attempts with you. If you couldn't do a question make sure to have a note written down about why you couldn't do the question.
- ▶ Seek help with these (and other problems) from tutors
- ▶ Bring lecture notes to tutorials!

# Tutorials

- ▶ Tutorial questions on the exercise sheets should be done each week before the tutorial
- ▶ Do this work even if you don't have a tutorial that week
- ▶ Tutors will ask to see your tutorial work so make sure you have your attempts with you. If you couldn't do a question make sure to have a note written down about why you couldn't do the question.
- ▶ Seek help with these (and other problems) from tutors
- ▶ Bring lecture notes to tutorials!
- ▶ Tutorials are an important resource and opportunity for getting feedback. Be proactive and ask tutors to look at your work and ask them questions.

## Other arrangements

- ▶ **Office hours:** Office number 505 Mathematics Building.  
Office hours: Monday 12-1, Tuesday 12-1, Wednesday 1-2 (or by arrangement)



## Other arrangements

- ▶ **Office hours:** Office number 505 Mathematics Building.  
Office hours: Monday 12-1, Tuesday 12-1, Wednesday 1-2 (or by arrangement)
- ▶ **Continuous assessment:**

## Other arrangements

- ▶ **Office hours:** Office number 505 Mathematics Building.  
Office hours: Monday 12-1, Tuesday 12-1, Wednesday 1-2 (or by arrangement)
- ▶ **Continuous assessment:**
  - ▶ weekly online assignment, from WebAssign (due weeks 2-11, Saturday 3am).

## Other arrangements

- ▶ **Office hours:** Office number 505 Mathematics Building.  
Office hours: Monday 12-1, Tuesday 12-1, Wednesday 1-2 (or by arrangement)
- ▶ **Continuous assessment:**
  - ▶ weekly online assignment, from WebAssign (due weeks 2-11, Saturday 3am).
  - ▶ **The 2A Assignment 1 is due 3am Saturday 5th October**

## Other arrangements

- ▶ **Office hours:** Office number 505 Mathematics Building.  
Office hours: Monday 12-1, Tuesday 12-1, Wednesday 1-2 (or by arrangement)
- ▶ **Continuous assessment:**
  - ▶ weekly online assignment, from WebAssign (due weeks 2-11, Saturday 3am).
  - ▶ The 2A Assignment 1 is due 3am Saturday 5th October
  - ▶ together these make up 20% of final grade

## Other arrangements

- ▶ **Office hours:** Office number 505 Mathematics Building.  
Office hours: Monday 12-1, Tuesday 12-1, Wednesday 1-2 (or by arrangement)
- ▶ **Continuous assessment:**
  - ▶ weekly online assignment, from WebAssign (due weeks 2-11, Saturday 3am).
  - ▶ The 2A Assignment 1 is due 3am Saturday 5th October
  - ▶ together these make up 20% of final grade
  - ▶ in calculating your continuous assessment grade we discard your worst TWO submissions

## Other arrangements

- ▶ **Office hours:** Office number 505 Mathematics Building.  
Office hours: Monday 12-1, Tuesday 12-1, Wednesday 1-2 (or by arrangement)
- ▶ **Continuous assessment:**
  - ▶ weekly online assignment, from WebAssign (due weeks 2-11, Saturday 3am).
  - ▶ The 2A Assignment 1 is due 3am Saturday 5th October
  - ▶ together these make up 20% of final grade
  - ▶ in calculating your continuous assessment grade we discard your worst TWO submissions
  - ▶ **course work cannot be reassessed. This means there is no second attempts at assignments**

## Other arrangements

- ▶ **Office hours:** Office number 505 Mathematics Building.  
Office hours: Monday 12-1, Tuesday 12-1, Wednesday 1-2 (or by arrangement)
- ▶ **Continuous assessment:**
  - ▶ weekly online assignment, from WebAssign (due weeks 2-11, Saturday 3am).
  - ▶ The 2A Assignment 1 is due 3am Saturday 5th October
  - ▶ together these make up 20% of final grade
  - ▶ in calculating your continuous assessment grade we discard your worst TWO submissions
  - ▶ course work cannot be reassessed. This means there is no second attempts at assignments
- ▶ **Degree Examination:** End of December (date not set yet)

## Other arrangements

- ▶ **Office hours:** Office number 505 Mathematics Building.  
Office hours: Monday 12-1, Tuesday 12-1, Wednesday 1-2 (or by arrangement)
- ▶ **Continuous assessment:**
  - ▶ weekly online assignment, from WebAssign (due weeks 2-11, Saturday 3am).
  - ▶ The 2A Assignment 1 is due 3am Saturday 5th October
  - ▶ together these make up 20% of final grade
  - ▶ in calculating your continuous assessment grade we discard your worst TWO submissions
  - ▶ course work cannot be reassessed. This means there is no second attempts at assignments
- ▶ **Degree Examination:** End of December (date not set yet)
  - ▶ worth 80% of final assessment



## Other arrangements

- ▶ **Office hours:** Office number 505 Mathematics Building.  
Office hours: Monday 12-1, Tuesday 12-1, Wednesday 1-2 (or by arrangement)
- ▶ **Continuous assessment:**
  - ▶ weekly online assignment, from WebAssign (due weeks 2-11, Saturday 3am).
  - ▶ The 2A Assignment 1 is due 3am Saturday 5th October
  - ▶ together these make up 20% of final grade
  - ▶ in calculating your continuous assessment grade we discard your worst TWO submissions
  - ▶ course work cannot be reassessed. This means there is no second attempts at assignments
- ▶ **Degree Examination:** End of December (date not set yet)
  - ▶ worth 80% of final assessment
- ▶ **Recommended course book:** James Stewart, Multivariable Calculus International Edition, (Seventh Edition), Brooks Cole /Cengage .

# Lecture notes

- ▶ Exercise sheets are available on Moodle at midday on Monday each week, except Exercise Sheet 1 which is available in the first lecture

## Lecture notes

- ▶ Exercise sheets are available on Moodle at midday on Monday each week, except Exercise Sheet 1 which is available in the first lecture
- ▶ Exercises and solutions and printed lecture notes available at <http://moodle.gla.ac.uk/fims/moodle>  
(Also given in Level-2 General Information)

## Lecture notes

- ▶ Exercise sheets are available on Moodle at midday on Monday each week, except Exercise Sheet 1 which is available in the first lecture
- ▶ Exercises and solutions and printed lecture notes available at <http://moodle.gla.ac.uk/fims/moodle>  
(Also given in Level-2 General Information)
- ▶ Solutions to exercise sheets and solutions to examples given in lectures will appear on Moodle once each Chapter has been taught and not before.

## Lecture notes

- ▶ Exercise sheets are available on Moodle at midday on Monday each week, except Exercise Sheet 1 which is available in the first lecture
- ▶ Exercises and solutions and printed lecture notes available at <http://moodle.gla.ac.uk/fims/moodle>  
(Also given in Level-2 General Information)
- ▶ Solutions to exercise sheets and solutions to examples given in lectures will appear on Moodle once each Chapter has been taught and not before.
- ▶ Only the Chapter 1 lecture notes will be given out in class. You need to download the notes for Chapters 2,3 and 4 from Moodle yourself in advance.

# Syllabus of 2A - *Multivariate Calculus*

General theme: differentiation and integration of functions of several variables and its applications.

## Syllabus of 2A - *Multivariate Calculus*

General theme: differentiation and integration of functions of several variables and its applications.

► Chapter 1 - Partial differentiation

## Syllabus of 2A - *Multivariate Calculus*

General theme: differentiation and integration of functions of several variables and its applications.

- ▶ Chapter 1 - Partial differentiation
- ▶ Chapter 2 - Double and triple integration



## Syllabus of 2A - *Multivariate Calculus*

General theme: differentiation and integration of functions of several variables and its applications.

- ▶ Chapter 1 - Partial differentiation
- ▶ Chapter 2 - Double and triple integration
- ▶ Chapter 3 - Differentiation of vectors

## Syllabus of 2A - *Multivariate Calculus*

General theme: differentiation and integration of functions of several variables and its applications.

- ▶ Chapter 1 - Partial differentiation
- ▶ Chapter 2 - Double and triple integration
- ▶ Chapter 3 - Differentiation of vectors
- ▶ Chapter 4 - Line and surface integrals

## Syllabus of 2A - *Multivariate Calculus*

General theme: differentiation and integration of functions of several variables and its applications.

- ▶ Chapter 1 - Partial differentiation
- ▶ Chapter 2 - Double and triple integration
- ▶ Chapter 3 - Differentiation of vectors
- ▶ Chapter 4 - Line and surface integrals

## Syllabus of 2A - *Multivariate Calculus*

General theme: differentiation and integration of functions of several variables and its applications.

- ▶ Chapter 1 - Partial differentiation
- ▶ Chapter 2 - Double and triple integration
- ▶ Chapter 3 - Differentiation of vectors
- ▶ Chapter 4 - Line and surface integrals

# Chapter 1: Partial differentiation

- ▶ Recall ideas for functions of one variable

# Chapter 1: Partial differentiation

- ▶ Recall ideas for functions of one variable
- ▶ extend to functions of two variables,

# Chapter 1: Partial differentiation

- ▶ Recall ideas for functions of one variable
- ▶ extend to functions of two variables,
- ▶ introduce partial derivatives,

# Chapter 1: Partial differentiation

- ▶ Recall ideas for functions of one variable
- ▶ extend to functions of two variables,
- ▶ introduce partial derivatives,
- ▶ chain rule for partial derivatives.



## Functions of one variable

For example, volume  $V$  of a sphere is a function of one variable, its radius  $r$ ,

$$V = \frac{4}{3}\pi r^3.$$

We write  $V = f(r)$ , where the *rule* is  $f(r) = \frac{4}{3}\pi r^3$ .

## Functions of one variable

For example, volume  $V$  of a sphere is a function of one variable, its radius  $r$ ,

$$V = \frac{4}{3}\pi r^3.$$

We write  $V = f(r)$ , where the *rule* is  $f(r) = \frac{4}{3}\pi r^3$ .

### Domain of $f$

- ▶ the set of points  $D$  at which rule is used,

## Functions of one variable

For example, volume  $V$  of a sphere is a function of one variable, its radius  $r$ ,

$$V = \frac{4}{3}\pi r^3.$$

We write  $V = f(r)$ , where the *rule* is  $f(r) = \frac{4}{3}\pi r^3$ .

### Domain of $f$

- ▶ the set of points  $D$  at which rule is used,
- ▶ radius should be real and non-negative, so  $D = [0, \infty)$ ,

## Functions of one variable

For example, volume  $V$  of a sphere is a function of one variable, its radius  $r$ ,

$$V = \frac{4}{3}\pi r^3.$$

We write  $V = f(r)$ , where the *rule* is  $f(r) = \frac{4}{3}\pi r^3$ .

### Domain of $f$

- ▶ the set of points  $D$  at which rule is used,
- ▶ radius should be real and non-negative, so  $D = [0, \infty)$ ,
- ▶ if no domain were given, we use the *maximal domain* - set of points at which the rule could legitimately be used,

## Functions of one variable

For example, volume  $V$  of a sphere is a function of one variable, its radius  $r$ ,

$$V = \frac{4}{3}\pi r^3.$$

We write  $V = f(r)$ , where the *rule* is  $f(r) = \frac{4}{3}\pi r^3$ .

### Domain of $f$

- ▶ the set of points  $D$  at which rule is used,
- ▶ radius should be real and non-negative, so  $D = [0, \infty)$ ,
- ▶ if no domain were given, we use the *maximal domain* - set of points at which the rule could legitimately be used,
- ▶ the maximal domain of  $f$  is  $\mathbb{R}$ .

# Functions of one variable

## Graph

The set of all ordered pairs  $(a, f(a))$  where  $a \in D$ . Usually shown as a curve in the plane.

# Functions of one variable

## Graph

The set of all ordered pairs  $(a, f(a))$  where  $a \in D$ . Usually shown as a curve in the plane.


Present example - the set of point  $(r, \frac{4}{3}\pi r^3)$  for all  $r \geq 0$ .

# Functions of one variable

## Graph

The set of all ordered pairs  $(a, f(a))$  where  $a \in D$ . Usually shown as a curve in the plane.

Present example - the set of point  $(r, \frac{4}{3}\pi r^3)$  for all  $r \geq 0$ .





# Functions of two variables

## Example

Volume  $V$  of a cylinder depends on *two* dimensions, the radius  $r$  and the height  $h$  -  $V = f(r, h)$ , where  $f(r, h) = \pi r^2 h$  defines a *function of two variables*.

# Functions of two variables

## Example

Volume  $V$  of a cylinder depends on *two* dimensions, the radius  $r$  and the height  $h$  -  $V = f(r, h)$ , where  $f(r, h) = \pi r^2 h$  defines a *function of two variables*.

## Domain


Subset  $D$  of  $\mathbb{R}^2$ , i.e., a region in a plane.

If not specified, the maximal domain is assumed.

# Functions of two variables

## Graph

The set of points  $(a, b, c) \in \mathbb{R}^3$  where  $(a, b) \in D$  and  $c = f(a, b)$  - a *surface*.



## Visualisation of surfaces - Spheres

- ▶ Radius  $r$ , centre  $(a, b, c)$  - points  $(x, y, z)$  a distance  $r$  from  $(a, b, c)$ . Pythagoras's theorem  $\implies$

$$(x - a)^2 + (y - b)^2 + (z - c)^2 = r^2,$$

## Visualisation of surfaces - Spheres

- ▶ Radius  $r$ , centre  $(a, b, c)$  - points  $(x, y, z)$  a distance  $r$  from  $(a, b, c)$ . Pythagoras's theorem  $\implies$

$$(x - a)^2 + (y - b)^2 + (z - c)^2 = r^2,$$

- ▶ solving for  $z$

$$z = c \pm \sqrt{r^2 - (x - a)^2 - (y - b)^2},$$

## Visualisation of surfaces - Spheres

- ▶ Radius  $r$ , centre  $(a, b, c)$  - points  $(x, y, z)$  a distance  $r$  from  $(a, b, c)$ . Pythagoras's theorem  $\implies$

$$(x - a)^2 + (y - b)^2 + (z - c)^2 = r^2,$$

- ▶ solving for  $z$

$$z = c \pm \sqrt{r^2 - (x - a)^2 - (y - b)^2},$$

- ▶  $+$  means “northern” hemisphere  
– means “southern” hemisphere.

## Visualisation of surfaces - Spheres

► Given

$$x^2 + y^2 + z^2 + \alpha x + \beta y + \gamma z + \delta = 0,$$

complete the square to write as

## Visualisation of surfaces - Spheres

- ▶ Given

$$x^2 + y^2 + z^2 + \alpha x + \beta y + \gamma z + \delta = 0,$$

complete the square to write as



## Visualisation of surfaces - Spheres

- ▶ Given

$$x^2 + y^2 + z^2 + \alpha x + \beta y + \gamma z + \delta = 0,$$

complete the square to write as

$$(x + \tfrac{1}{2}\alpha)^2 + (y + \tfrac{1}{2}\beta)^2 + (z + \tfrac{1}{2}\gamma)^2 = \tfrac{1}{4}(\alpha^2 + \beta^2 + \gamma^2) - \delta,$$

- ▶ sphere if and only if  $\tfrac{1}{4}(\alpha^2 + \beta^2 + \gamma^2) - \delta > 0$ .

# Visualisation of surfaces - Spheres

## Example 1


Sketch the graph of  $f(x, y) = -\sqrt{1 - 2x - x^2 - y^2}$ .

# Visualisation of surfaces - Spheres

## Example 1


Sketch the graph of  $f(x, y) = -\sqrt{1 - 2x - x^2 - y^2}$ .

Answer




# Visualisation of surfaces - Cross-sections

- For a surface  $z = f(x, y)$  the set of points satisfying,  $f(x, y) = c$ , is a *level curve* or *contour*,



## Visualisation of surfaces - Cross-sections

- ▶ For a surface  $z = f(x, y)$  the set of points satisfying,  $f(x, y) = c$ , is a *level curve* or *contour*,
- ▶ think of  $z = f(x, y)$  as part of the surface of the earth - each level curve represents a particular contour line on its map.



## Visualisation of surfaces - Cross-sections

- ▶ More generally, the intersection of plane  $x = \text{constant}$  or  $y = \text{constant}$  or  $z = \text{constant}$  and surface  $F(x, y, z) = 0$  is called a *cross-section*,

## Visualisation of surfaces - Cross-sections

- ▶ More generally, the intersection of plane  $x = \text{constant}$  or  $y = \text{constant}$  or  $z = \text{constant}$  and surface  $F(x, y, z) = 0$  is called a *cross-section*,
- ▶ level curve  $L_c$  is the set of points  $(x, y)$  in  $D$  for which  $f(x, y) = c$ ,

## Visualisation of surfaces - Cross-sections

- ▶ More generally, the intersection of plane  $x = \text{constant}$  or  $y = \text{constant}$  or  $z = \text{constant}$  and surface  $F(x, y, z) = 0$  is called a *cross-section*,
- ▶ level curve  $L_c$  is the set of points  $(x, y)$  in  $D$  for which  $f(x, y) = c$ ,
- ▶ for different  $c$ ,  $L_c$  may be a curve, a point or points, or the empty set,



## Visualisation of surfaces - Cross-sections

- ▶ More generally, the intersection of plane  $x = \text{constant}$  or  $y = \text{constant}$  or  $z = \text{constant}$  and surface  $F(x, y, z) = 0$  is called a *cross-section*,
- ▶ level curve  $L_c$  is the set of points  $(x, y)$  in  $D$  for which  $f(x, y) = c$ ,
- ▶ for different  $c$ ,  $L_c$  may be a curve, a point or points, or the empty set,
- ▶ each point in  $D$  lies on one level curve.

# Visualisation of surfaces - Cross-sections

## Example 2


By considering the level curves and the cross-sections  $x = 0$  and  $y = 0$ , obtain a sketch of  $z = \sqrt{x^2 + y^2}$ .

# Visualisation of surfaces - Cross-sections

## Example 2

By considering the level curves and the cross-sections  $x = 0$  and  $y = 0$ , obtain a sketch of  $z = \sqrt{x^2 + y^2}$ .

## Answer




# Visualisation of surfaces - Cross-sections

## Example 2

By considering the level curves and the cross-sections  $x = 0$  and  $y = 0$ , obtain a sketch of  $z = \sqrt{x^2 + y^2}$ .

## Answer

$x=0$  cross-section




# Visualisation of surfaces - Cross-sections

## Example 2

By considering the level curves and the cross-sections  $x = 0$  and  $y = 0$ , obtain a sketch of  $z = \sqrt{x^2 + y^2}$ .

## Answer

$y=0$  cross-section




# Visualisation of surfaces - Cross-sections

## Example 2

By considering the level curves and the cross-sections  $x = 0$  and  $y = 0$ , obtain a sketch of  $z = \sqrt{x^2 + y^2}$ .

Answer



## Visulatisation of surfaces - Ellipsoid

- ▶ An **ellipsoid** of radius  $r_1$  in the  $x$ -direction,  $r_2$  in the  $y$ -direction and  $r_3$  in the  $z$ -direction, with centre  $(a, b, c)$  is defined by

$$\frac{(x - a)^2}{(r_1)^2} + \frac{(y - b)^2}{(r_2)^2} + \frac{(z - c)^2}{(r_3)^2} = 1.$$

## Visualisation of surfaces - Ellipsoid

- ▶ An **ellipsoid** of radius  $r_1$  in the  $x$ -direction,  $r_2$  in the  $y$ -direction and  $r_3$  in the  $z$ -direction, with centre  $(a, b, c)$  is defined by

$$\frac{(x - a)^2}{(r_1)^2} + \frac{(y - b)^2}{(r_2)^2} + \frac{(z - c)^2}{(r_3)^2} = 1.$$

- ▶ The cross sections are ellipses, hence the name ellipsoid.



## Visulatisation of surfaces - Ellipsoid

- ▶ An **ellipsoid** of radius  $r_1$  in the  $x$ -direction,  $r_2$  in the  $y$ -direction and  $r_3$  in the  $z$ -direction, with centre  $(a, b, c)$  is defined by

$$\frac{(x - a)^2}{(r_1)^2} + \frac{(y - b)^2}{(r_2)^2} + \frac{(z - c)^2}{(r_3)^2} = 1.$$

- ▶ The cross sections are ellipses, hence the name ellipsoid.
- ▶ When  $r_1 = r_2 = r_3$  we recover the equation for the sphere.

## Visualisation of surfaces - Planes

- ▶ Recall: a plane with normal vector  $\mathbf{n} = (\alpha, \beta, \gamma)$  has equation  $\alpha x + \beta y + \gamma z = \delta$ ,

## Visualisation of surfaces - Planes

- ▶ Recall: a plane with normal vector  $\mathbf{n} = (\alpha, \beta, \gamma)$  has equation  $\alpha x + \beta y + \gamma z = \delta$ ,
- ▶ the graph of  $f(x, y) = ax + by + c$  is the plane  $z = ax + by + c$  with normal  $(a, b, -1)$  passing through the point  $(0, 0, c)$ .

## Visualisation of surfaces - Planes

### Example 3


Sketch the part of the surface  $2x + y + 4z = 1$  where  $x, y, z \geq 0$ .

## Visualisation of surfaces - Planes

### Example 3

Sketch the part of the surface  $2x + y + 4z = 1$  where  $x, y, z \geq 0$ .

Answer



## Visualisation of surfaces - Circular cylinder

- ▶ A circular **cylinder** in  $\mathbb{R}^3$  of radius  $r$  centred at the origin lying parallel to  $z$ -axis, is defined by

$$x^2 + y^2 = r^2.$$

## Visualisation of surfaces - Circular cylinder

- ▶ A circular **cylinder** in  $\mathbb{R}^3$  of radius  $r$  centred at the origin lying parallel to  $z$ -axis, is defined by

$$x^2 + y^2 = r^2.$$

- ▶ This is **NOT** the equation for a circle, because the surface lies in  $\mathbb{R}^3$ .

## Visualisation of surfaces - Circular cylinder

- ▶ A circular **cylinder** in  $\mathbb{R}^3$  of radius  $r$  centred at the origin lying parallel to  $z$ -axis, is defined by

$$x^2 + y^2 = r^2.$$

- ▶ This is **NOT** the equation for a circle, because the surface lies in  $\mathbb{R}^3$ .
- ▶ Generalisable to cylinders centred at  $(a, b, c)$ , cylinders lying parallel to the  $x$  or  $y$  axes and cylinders with ellipses as cross sections.



## Visualisation of surfaces - Paraboloid

### Example 4


Sketch the region bounded by the paraboloid  $z = 4 - x^2 - 2y^2$  and the plane  $z = 2$ .

# Visualisation of surfaces - Paraboloid

## Example 4

Sketch the region bounded by the paraboloid  $z = 4 - x^2 - 2y^2$  and the plane  $z = 2$ .

## Answer




# Visualisation of surfaces - Paraboloid

## Example 4

Sketch the region bounded by the paraboloid  $z = 4 - x^2 - 2y^2$  and the plane  $z = 2$ .

## Answer



# Ordinary derivative

► Recall:

$$\lim_{h \rightarrow 0} \frac{g(a+h) - g(a)}{h}$$

(if it exists) is the *derivative* of  $g$  at  $a$ ,

# Ordinary derivative

- Recall:

$$\lim_{h \rightarrow 0} \frac{g(a+h) - g(a)}{h}$$

(if it exists) is the *derivative* of  $g$  at  $a$ ,

- written as

$$\frac{dg}{dx}(a) \quad \text{or} \quad g'(a),$$

# Ordinary derivative

- Recall:

$$\lim_{h \rightarrow 0} \frac{g(a+h) - g(a)}{h}$$

(if it exists) is the *derivative* of  $g$  at  $a$ ,


- written as

$$\frac{dg}{dx}(a) \quad \text{or} \quad g'(a),$$

- *gradient* of the tangent to the graph of  $g$  at a point  $(a, g(a))$ .

# Ordinary derivative

- *gradient* of the tangent to the graph of  $g$  at a point  $(a, g(a))$ .



## Partial derivatives

- ▶ On surface  $z = f(x, y)$ , there is no single meaning of gradient,



## Partial derivatives


- ▶ On surface  $z = f(x, y)$ , there is no single meaning of gradient,
- ▶ straight down a mountain side gradient may be very large and traversing the mountain the gradient is much less,

## Partial derivatives

- ▶ On surface  $z = f(x, y)$ , there is no single meaning of gradient,
- ▶ straight down a mountain side gradient may be very large and traversing the mountain the gradient is much less,
- ▶ necessary to define *two* gradients on cross-section of the surface in the  $x$  and  $y$  directions.

## Partial derivatives

Taking cross-sections  $x = a$  and  $y = b$  we get the graphs of two functions of *one* variable -  $z = f(x, b) = g(x)$  and  $z = f(a, y) = h(y)$



## Partial derivatives

- ▶ The gradients to  $z = g(x)$  and  $z = h(y)$  are called the *partial  $x$  and  $y$  derivatives of  $f$  at  $(a, b)$*

## Partial derivatives

- ▶ The gradients to  $z = g(x)$  and  $z = h(y)$  are called the *partial  $x$  and  $y$  derivatives of  $f$  at  $(a, b)$*
- ▶ written as

$$\frac{\partial f}{\partial x}(a, b) = \text{derivative w.r.t. } x \text{ with } y \text{ constant} - \text{equals } g'(a),$$

$$\frac{\partial f}{\partial y}(a, b) = \text{derivative w.r.t. } y \text{ with } x \text{ constant} - \text{equals } h'(b),$$

## Partial derivatives

- ▶ The gradients to  $z = g(x)$  and  $z = h(y)$  are called the *partial  $x$  and  $y$  derivatives of  $f$  at  $(a, b)$*
- ▶ written as

$$\frac{\partial f}{\partial x}(a, b) = \text{derivative w.r.t. } x \text{ with } y \text{ constant} - \text{equals } g'(a),$$

$$\frac{\partial f}{\partial y}(a, b) = \text{derivative w.r.t. } y \text{ with } x \text{ constant} - \text{equals } h'(b),$$

- ▶ for a function of  $x_1, x_2, \dots, x_n$

$$\frac{\partial f}{\partial x_i} = \text{derivative w.r.t. } x_i \text{ with all other variables constant.}$$

## Partial derivatives

- Important to distinguish notation used for ordinary and partial derivatives.

Ordinary derivative :  $\frac{df}{dx}$ ,    partial derivative :  $\frac{\partial f}{\partial x}$ ,

## Partial derivatives

- Important to distinguish notation used for ordinary and partial derivatives.

Ordinary derivative :  $\frac{df}{dx}$ ,    partial derivative :  $\frac{\partial f}{\partial x}$ ,

- subscript notation for partial derivatives

$$\frac{\partial f}{\partial x} \equiv f_x, \text{ and } \frac{\partial f}{\partial y} \equiv f_y,$$



## Partial derivatives

### Example 5

Find  $f_x$ ,  $f_y$  and  $z_x$  where

$$(a) f(x, y) = x^3 y^2 + x, \quad (b) z(x, y) = \sin^{-1} \left( \frac{x}{x+y} \right) \text{ and } x, y > 0.$$

## Partial derivatives

### Example 5

Find  $f_x$ ,  $f_y$  and  $z_x$  where

$$(a) f(x, y) = x^3 y^2 + x, \quad (b) z(x, y) = \sin^{-1} \left( \frac{x}{x+y} \right) \text{ and } x, y > 0.$$

[ $\sin^{-1} u$  is the inverse sine function and *not* the reciprocal  $1/\sin u$ .  
Domain of  $\sin^{-1}$  is  $[-1, 1]$  and  $x/(x+y)$  lies in this domain.]

## Partial derivatives

### Example 5

Find  $f_x$ ,  $f_y$  and  $z_x$  where

$$(a) f(x, y) = x^3 y^2 + x, \quad (b) z(x, y) = \sin^{-1} \left( \frac{x}{x+y} \right) \text{ and } x, y > 0.$$

### Answer

$$(a) f_x = 3x^2 y^2 + 1, \quad f_y = 2x^3 y.$$

$$(b) z_x = \frac{y}{x+y} \frac{1}{\sqrt{2xy + y^2}}.$$

## Partial derivatives

### Example 5

Find  $f_x$ ,  $f_y$  and  $z_x$  where

$$(a) f(x, y) = x^3 y^2 + x, \quad (b) z(x, y) = \sin^{-1} \left( \frac{x}{x+y} \right) \text{ and } x, y > 0.$$

### Chain rule

Recall from Level-1:

$$\frac{d}{dx} f(g(x)) = f'(g(x)) g'(x).$$

We used

$$\frac{\partial}{\partial x} f(g(x, y)) = f'(g(x, y)) g_x(x, y).$$

# Partial derivatives

## Example 6

Find  $z_x$  where  $z$  is defined implicitly as a function of  $x$  and  $y$  by the equation

$$x^4 + 2y^2 + z^3 - 2x^2yz = 1$$

## Partial derivatives

### Example 6

Find  $z_x$  where  $z$  is defined implicitly as a function of  $x$  and  $y$  by the equation

$$x^4 + 2y^2 + z^3 - 2x^2yz = 1$$

### Answer

$$z_x = \frac{4x^3 - 4xyz}{2x^2y - 3z^2}$$

## Partial derivatives

### Example 7

For  $r \in \mathbb{R}^+$ , let  $u = f(r)$  where  $r^2 = x^2 + y^2 + z^2$ . Show that

$$xu_x + yu_y + zu_z = rf'(r).$$

## Higher order derivatives

Let  $u$  be a function of  $x, y, \dots$  then  $u_x$  and  $u_y$  are functions of  $x, y, \dots$  and so may define



## Higher order derivatives

Let  $u$  be a function of  $x, y, \dots$  then  $u_x$  and  $u_y$  are functions of  $x, y, \dots$  and so may define

$$\frac{\partial^2 u}{\partial x^2} = \frac{\partial}{\partial x}(u_x) = u_{xx}, \quad \frac{\partial^2 u}{\partial y \partial x} = \frac{\partial}{\partial y}(u_x) = u_{xy},$$

## Higher order derivatives

Let  $u$  be a function of  $x, y, \dots$  then  $u_x$  and  $u_y$  are functions of  $x, y, \dots$  and so may define

$$\frac{\partial^2 u}{\partial x^2} = \frac{\partial}{\partial x}(u_x) = u_{xx}, \quad \frac{\partial^2 u}{\partial y \partial x} = \frac{\partial}{\partial y}(u_x) = u_{xy},$$

$$\frac{\partial^2 u}{\partial x \partial y} = \frac{\partial}{\partial x}(u_y) = u_{yx}, \quad \frac{\partial^2 u}{\partial y^2} = \frac{\partial}{\partial y}(u_y) = u_{yy},$$

etc.

## Higher order derivatives

There is no automatic guarantee that  $u_{xy} = u_{yx}$  but...

## Higher order derivatives

There is no automatic guarantee that  $u_{xy} = u_{yx}$

► **Theorem:**

Let  $u$  be a function of  $x, y$  such  $u_{xy}$  and  $u_{yx}$  exist and are continuous at a point  $(a, b)$ . Then,

$$u_{xy}(a, b) = u_{yx}(a, b).$$

## Higher order derivatives

There is no automatic guarantee that  $u_{xy} = u_{yx}$

► **Theorem:**

Let  $u$  be a function of  $x, y$  such  $u_{xy}$  and  $u_{yx}$  exist and are continuous at a point  $(a, b)$ . Then,

$$u_{xy}(a, b) = u_{yx}(a, b).$$

- Also for functions of more variables and higher order derivatives - e.g. if  $u = u(x, y, z)$  then

$$u_{xyxz} = u_{zyxx} = u_{yxxz} = \cdots = u_{xxyz},$$

## Higher order derivatives

There is no automatic guarantee that  $u_{xy} = u_{yx}$

► **Theorem:**

Let  $u$  be a function of  $x, y$  such  $u_{xy}$  and  $u_{yx}$  exist and are continuous at a point  $(a, b)$ . Then,

$$u_{xy}(a, b) = u_{yx}(a, b).$$

- Also for functions of more variables and higher order derivatives - e.g. if  $u = u(x, y, z)$  then

$$u_{xyxz} = u_{zyxx} = u_{yxxz} = \cdots = u_{xxyz},$$

- in 2A, we assume the order of taking partial derivatives is unimportant.

## Higher order derivatives

### Example 8

Determine all second order derivatives of  $u = \sin xy$  and verify that

$$u_{xy} = u_{yx}.$$

## Higher order derivatives

### Example 8

Determine all second order derivatives of  $u = \sin xy$  and verify that  $u_{xy} = u_{yx}$ .

### Answers

$$u_{xx} = -y^2 \sin xy,$$

$$u_{xy} = \cos xy - yx \sin xy,$$

$$u_{yx} = \cos xy - xy \sin xy,$$

$$u_{yy} = -x^2 \sin xy.$$



## Higher order derivatives

### Example 9

Let  $u = f(x/y)$ , where  $f$  is an arbitrary (twice differentiable, with continuous second derivative) function of one variable. Show that

$$xu_x + yu_y = 0,$$

and *deduce* that

$$x^2 u_{xx} + 2xy u_{xy} + y^2 u_{yy} = 0.$$

## Two variable chain rule

- ▶ Chain rule for functions of one variable - used to find derivative of  $F(x) = f(u(x))$  -

$$\frac{dF}{dx} = \frac{du}{dx} \frac{df}{du} = u'(x)f'(u(x)).$$

## Two variable chain rule

- ▶ Chain rule for functions of one variable - used to find derivative of  $F(x) = f(u(x))$  -

$$\frac{dF}{dx} = \frac{du}{dx} \frac{df}{du} = u'(x)f'(u(x)).$$

- ▶ extend this technique to functions of several variables

## Two variable chain rule

- ▶ Chain rule for functions of one variable - used to find derivative of  $F(x) = f(u(x))$  -

$$\frac{dF}{dx} = \frac{du}{dx} \frac{df}{du} = u'(x) f'(u(x)).$$

- ▶ extend this technique to functions of several variables

- ▶ **Theorem**

Let  $F(x, y) = f(u(x, y), v(x, y))$ . Then

$$\frac{\partial F}{\partial x} = \frac{\partial u}{\partial x} \frac{\partial f}{\partial u} + \frac{\partial v}{\partial x} \frac{\partial f}{\partial v} \quad \text{and} \quad \frac{\partial F}{\partial y} = \frac{\partial u}{\partial y} \frac{\partial f}{\partial u} + \frac{\partial v}{\partial y} \frac{\partial f}{\partial v}.$$

This is called the *chain rule for functions of two variables*.

## Two variable chain rule

- Observe the pattern

$$\frac{\partial F}{\partial x} = \frac{\boxed{\partial u}}{\partial x} \frac{\partial f}{\boxed{\partial u}} + \frac{\boxed{\partial v}}{\partial x} \frac{\partial f}{\boxed{\partial v}},$$

## Two variable chain rule

- Observe the pattern

$$\frac{\partial F}{\partial x} = \frac{\boxed{\partial u}}{\partial x} \frac{\partial f}{\boxed{\partial u}} + \frac{\boxed{\partial v}}{\partial x} \frac{\partial f}{\boxed{\partial v}},$$

- extends in an obvious way to functions of any number of variables - if  $F(x, y, z) = f(u(x, y, z), v(x, y, z), w(x, y, z))$  then

$$\frac{\partial F}{\partial x} = \frac{\partial u}{\partial x} \frac{\partial f}{\partial u} + \frac{\partial v}{\partial x} \frac{\partial f}{\partial v} + \frac{\partial w}{\partial x} \frac{\partial f}{\partial w}.$$

## Two variable chain rule - Special cases

► if  $F(x, y) = f(u(x, y))$  then

$$\frac{\partial F}{\partial x} = \frac{\partial u}{\partial x} \frac{df}{du};$$

## Two variable chain rule - Special cases

- ▶ if  $F(x, y) = f(u(x, y))$  then

$$\frac{\partial F}{\partial x} = \frac{\partial u}{\partial x} \frac{df}{du};$$

- ▶ if  $F(x) = f(u(x), v(x))$  then

$$\frac{dF}{dx} = \frac{du}{dx} \frac{\partial f}{\partial u} + \frac{dv}{dx} \frac{\partial f}{\partial v}.$$



## Two variable chain rule - Special cases

- ▶ if  $F(x, y) = f(u(x, y))$  then

$$\frac{\partial F}{\partial x} = \frac{\partial u}{\partial x} \frac{df}{du};$$

- ▶ if  $F(x) = f(u(x), v(x))$  then

$$\frac{dF}{dx} = \frac{du}{dx} \frac{\partial f}{\partial u} + \frac{dv}{dx} \frac{\partial f}{\partial v}.$$

Partial derivatives are written as ordinary derivatives when used on functions of one variable.

## Two variable chain rule

### Example 10

Let  $w = u^2 + v^2$  where  $u = \sin \theta$  and  $v = \cos \phi$ . Use the chain rule to calculate  $w_\theta$  and  $w_\phi$  in terms of  $\theta$  and  $\phi$ .

## Two variable chain rule

### Example 10

Let  $w = u^2 + v^2$  where  $u = \sin \theta$  and  $v = \cos \phi$ . Use the chain rule to calculate  $w_\theta$  and  $w_\phi$  in terms of  $\theta$  and  $\phi$ .

### Answer

$$w_\theta = \sin 2\theta, \quad w_\phi = -\sin 2\phi.$$

## Examples of ODEs and PDEs

- ▶ An *ordinary differential equation* (ODE) is a relationship between a function of one variable and its derivatives

## Examples of ODEs and PDEs

- ▶ An *ordinary differential equation* (ODE) is a relationship between a function of one variable and its derivatives
- ▶ e.g Newton's law of cooling states that  
“the rate of change of temperature of an object is proportional to the temperature difference between it and its surroundings”

## Examples of ODEs and PDEs

- ▶ An *ordinary differential equation* (ODE) is a relationship between a function of one variable and its derivatives
- ▶ e.g Newton's law of cooling states that  
"the rate of change of temperature of an object is proportional to the temperature difference between it and its surroundings"
- ▶ in mathematical terms this is the differential equation

$$\frac{dT}{dt} = k(T - T_0),$$

where  $T(t)$  is the temperature,  $T_0$  the temperature of the surroundings and  $k$  a constant

## Examples of ODEs and PDEs

- ▶ A *partial differential equation* (PDE) is a relationship between a function of more than one variable and its partial derivatives

## Examples of ODEs and PDEs

- ▶ A *partial differential equation* (PDE) is a relationship between a function of more than one variable and its partial derivatives
- ▶ e.g. the wave equation,

$$\frac{\partial^2 u}{\partial t^2} = c^2 \frac{\partial^2 u}{\partial x^2},$$

where  $u(x, t)$  is the displacement (from a rest position) of the point  $x$  at time  $t$  and  $c$  is the wave speed.



# Definitions

- ▶ The *order* of a differential equation is the order of the highest derivative

# Definitions

- ▶ The *order* of a differential equation is the order of the highest derivative
- ▶ the unknown function is called the *dependent variable*

# Definitions

- ▶ The *order* of a differential equation is the order of the highest derivative
- ▶ the unknown function is called the *dependent variable*
- ▶ the variable or variables on which it depends are the *independent variable(s)*

# Definitions

- ▶ The *order* of a differential equation is the order of the highest derivative
- ▶ the unknown function is called the *dependent variable*
- ▶ the variable or variables on which it depends are the *independent variable(s)*
- ▶ a *solution* is an expression for the dependent variable which satisfies the relation

## Definitions

- ▶ The *order* of a differential equation is the order of the highest derivative
- ▶ the unknown function is called the *dependent variable*
- ▶ the variable or variables on which it depends are the *independent variable(s)*
- ▶ a *solution* is an expression for the dependent variable which satisfies the relation
- ▶ the *general solution* includes all possible solutions—including arbitrary constants (ODE) or arbitrary functions (PDE)

# Definitions

- ▶ The *order* of a differential equation is the order of the highest derivative
- ▶ the unknown function is called the *dependent variable*
- ▶ the variable or variables on which it depends are the *independent variable(s)*
- ▶ a *solution* is an expression for the dependent variable which satisfies the relation
- ▶ the *general solution* includes all possible solutions—includes arbitrary constants (ODE) or arbitrary functions (PDE)
- ▶ a solution without arbitrary constants/functions is called a *particular solution*. This may be found by giving extra conditions in the form of initial or boundary conditions.

# First order PDEs

## Example 11

Find the general solution of the PDE,

$$\frac{\partial f}{\partial x} = x^2 + y + 9,$$

where  $f$  is a function of two independent variables  $x$  and  $y$ .

# First order PDEs

## Example 11

Find the general solution of the PDE,

$$\frac{\partial f}{\partial x} = x^2 + y + 9,$$

where  $f$  is a function of two independent variables  $x$  and  $y$ .

## Answers

Solution is

$$\frac{1}{3}x^3 + xy + 9x + A(y)$$

where  $A$  is an arbitrary function.



# First order PDEs

## Example 12

Find the general solution of the PDE,

$$\frac{\partial^2 f}{\partial x \partial y} = 2x,$$

where  $f$  is a function of two independent variables  $x$  and  $y$ .

# First order PDEs

## Example 12

Find the general solution of the PDE,

$$\frac{\partial^2 f}{\partial x \partial y} = 2x,$$

where  $f$  is a function of two independent variables  $x$  and  $y$ .

## Answers

Solution is

$$x^2 y + A(y) + B(x),$$

where  $A$  and  $B$  are arbitrary functions.

## Solving PDEs using change of variable

- ▶ In this section we solve some first and second order PDEs by using a change of independent variables to write it in a simpler form

## Solving PDEs using change of variable

- ▶ In this section we solve some first and second order PDEs by using a change of independent variables to write it in a simpler form
- ▶ suppose the dependent variable is  $z$  and independent variables are  $x, y$

## Solving PDEs using change of variable

- ▶ In this section we solve some first and second order PDEs by using a change of independent variables to write it in a simpler form
- ▶ suppose the dependent variable is  $z$  and independent variables are  $x, y$
- ▶ if we change from  $x, y$  to  $u, v$  then the chain rule gives

$$\frac{\partial z}{\partial x} = \frac{\partial u}{\partial x} \frac{\partial z}{\partial u} + \frac{\partial v}{\partial x} \frac{\partial z}{\partial v}.$$

## Solving PDEs using change of variable

- ▶ In this section we solve some first and second order PDEs by using a change of independent variables to write it in a simpler form
- ▶ suppose the dependent variable is  $z$  and independent variables are  $x, y$
- ▶ if we change from  $x, y$  to  $u, v$  then the chain rule gives

$$\frac{\partial z}{\partial x} = \frac{\partial u}{\partial x} \frac{\partial z}{\partial u} + \frac{\partial v}{\partial x} \frac{\partial z}{\partial v}.$$

- ▶ in fact, for any expression  $E$  (e.g. a derivative of  $z$ )

$$\frac{\partial}{\partial x}(E) = \frac{\partial u}{\partial x} \frac{\partial}{\partial u}(E) + \frac{\partial v}{\partial x} \frac{\partial}{\partial v}(E) \quad (1)$$

this is used when we consider second order PDEs.

# First order PDEs

## Example 13

By changing variables from  $(x, y)$  to  $(u, v)$ , where  $u = xy$ ,  $v = x/y$ , solve the PDE

$$x \frac{\partial z}{\partial x} + y \frac{\partial z}{\partial y} = 2x^2 \sin(xy).$$

# First order PDEs

## Example 13

By changing variables from  $(x, y)$  to  $(u, v)$ , where  $u = xy$ ,  $v = x/y$ , solve the PDE

$$x \frac{\partial z}{\partial x} + y \frac{\partial z}{\partial y} = 2x^2 \sin(xy).$$

## Answers

Solution is

$$z = -\frac{x}{y} \cos(xy) + A(x/y),$$

where  $A$  is an arbitrary function.



# First order PDEs

## Example 14

By changing variables from  $(x, y)$  to  $(u, v)$ , where  $u = x^3/y$ ,  $v = x$ , find  $\frac{\partial f}{\partial x}$  and  $\frac{\partial f}{\partial y}$  in terms of partial derivatives with respect to  $u$  and  $v$ . Hence, solve the PDE

$$x \frac{\partial f}{\partial x} + 3y \frac{\partial f}{\partial y} = \frac{6x^5}{y}.$$

# First order PDEs

## Example 14

By changing variables from  $(x, y)$  to  $(u, v)$ , where  $u = x^3/y$ ,  $v = x$ , find  $\frac{\partial f}{\partial x}$  and  $\frac{\partial f}{\partial y}$  in terms of partial derivatives with respect to  $u$  and  $v$ . Hence, solve the PDE

$$x \frac{\partial f}{\partial x} + 3y \frac{\partial f}{\partial y} = \frac{6x^5}{y}.$$

## Answers

Solution is

$$f = \frac{3x^5}{y} + A(x^3/y),$$

where  $A$  is an arbitrary function.

## Chapter 2: Double and triple integration

- ▶ Double integration on regular domains,

## Chapter 2: Double and triple integration

- ▶ Double integration on regular domains,
- ▶ Double integration in polar coordinates (and Beta functions),

## Chapter 2: Double and triple integration

- ▶ Double integration on regular domains,
- ▶ Double integration in polar coordinates (and Beta functions),
- ▶ Double integration and general change of variables,

## Chapter 2: Double and triple integration

- ▶ Double integration on regular domains,
- ▶ Double integration in polar coordinates (and Beta functions),
- ▶ Double integration and general change of variables,
- ▶ Triple integration on regular domains,

## Chapter 2: Double and triple integration

- ▶ Double integration on regular domains,
- ▶ Double integration in polar coordinates (and Beta functions),
- ▶ Double integration and general change of variables,
- ▶ Triple integration on regular domains,
- ▶ Triple integration in spherical coordinates,

## Chapter 2: Double and triple integration

- ▶ Double integration on regular domains,
- ▶ Double integration in polar coordinates (and Beta functions),
- ▶ Double integration and general change of variables,
- ▶ Triple integration on regular domains,
- ▶ Triple integration in spherical coordinates,
- ▶ Using integrals to calculate area, volume and mass.




## Area under curves

- ▶ In first year definite integrals arise as “areas under curves”.


## Area under curves

- ▶ In first year definite integrals arise as “areas under curves”.
- ▶ We approximate the area under the curve the sum of areas of rectangles (called a *Riemann sum*)




## Area under curves

- ▶ In first year definite integrals arise as “areas under curves”.
- ▶ We approximate the area under the curve the sum of areas of rectangles (called a *Riemann sum*)



## Area under curves

- ▶ In first year definite integrals arise as “areas under curves”.
- ▶ We approximate the area under the curve the sum of areas of rectangles (called a *Riemann sum*)




$$\int_a^b g(x) dx = \lim_{N \rightarrow \infty} \sum_{i=1}^N g(x_i) \delta x_i.$$

## Double integration on rectangular domains


- ▶ Similarly, the “volume under a surface”  $z = f(x, y)$  on the set  $D \subset \mathbb{R}^2$  is approximated by the sum of the volumes of cuboids.

## Double integration on rectangular domains

- ▶ Similarly, the “volume under a surface”  $z = f(x, y)$  on the set  $D \subset \mathbb{R}^2$  is approximated by the sum of the volumes of cuboids.
- ▶ Divide  $R = [a, b] \times [c, d]$  into subrectangles of area  $\delta A_{ij} = \delta x_i \delta y_j$  and the cuboid above this has height  $f(x_i, y_j)$ .




## Double integration on rectangular domains



- The whole volume is approximated by

$$\iint_R f(x, y) dA = \iint_R f(x, y) dx dy = \lim_{M, N \rightarrow \infty} \sum_{i=1}^N \sum_{j=1}^M f(x_i, y_j) \delta A_{ij}.$$

# Double integration on rectangular domains



- The whole volume is approximated by


$$\iint_R f(x, y) dA = \iint_R f(x, y) dx dy = \lim_{M, N \rightarrow \infty} \sum_{i=1}^N \sum_{j=1}^M f(x_i, y_j) \delta A_{ij}.$$

- If the limit as  $M, N \rightarrow \infty$  exists we say that  $f$  is *integrable* over  $R$  and  $dA = dx dy$  is called the *area element*.




## Double integration on rectangular domains

The solid under the curve is made up of slices with  $y$  fixed



## Double integration on rectangular domains

The solid under the curve is made up of slices with  $y$  fixed




The area under the curve in such a cross section is

$$I(y) = \int_a^b f(x, y) dx,$$

where  $y$  is fixed in the integrand.

## Double integration on rectangular domains

The solid under the curve is made up of slices with  $y$  fixed



The area under the curve in such a cross section is

$$I(y) = \int_a^b f(x, y) dx,$$


where  $y$  is fixed in the integrand. The volume under the surface is then

$$\iint_R f(x, y) dx dy = \int_c^d I(y) dy = \int_c^d \left( \int_a^b f(x, y) dx \right) dy.$$

## Double integration on rectangular domains

Instead, summing the areas of cross sections of the solid with  $x$  fixed, we have

$$\iint_R f(x, y) \, dx dy = \int_a^b \left( \int_c^d f(x, y) \, dy \right) dx.$$




Notation:

## Double integration on rectangular domains

Instead, summing the areas of cross sections of the solid with  $x$  fixed, we have

$$\iint_R f(x, y) \, dx dy = \int_a^b \left( \int_c^d f(x, y) \, dy \right) dx.$$



Notation:

$$\int_a^b dx \int_c^d f(x, y) \, dy \quad \text{for} \quad \int_a^b \left( \int_c^d f(x, y) \, dy \right) dx.$$

# Double integration on rectangular domains

## Example 1

Evaluate  $\iint_R x^2 + y^2 \, dx dy$

where  $R$  is  $[1, 3] \times [2, 4]$ .

# Double integration on rectangular domains

## Example 1

Evaluate  $\iint_R x^2 + y^2 \, dx dy$


where  $R$  is  $[1, 3] \times [2, 4]$ .

Answer

$$\frac{164}{3}$$

## Double integration on regular domains


Consider a more complicated domain  $T$  which is the triangle with vertices  $(0, 0)$ ,  $(1, 0)$  and  $(1, 2)$ .





## Double integration on regular domains


Consider a more complicated domain  $T$  which is the triangle with vertices  $(0, 0)$ ,  $(1, 0)$  and  $(1, 2)$ .



The domain  $T$  is bounded by the lines  $y = 0$ ,  $x = 1$  and  $y = 2x$ .

## Double integration on regular domains


To evaluate a double integral over  $T$  we could split  $T$  into a collection of **vertical slices**,



integrate with respect to  $y$  and then integrate the result with respect to  $x$ .

## Double integration on regular domains

To evaluate a double integral over  $T$  we could split  $T$  into a collection of **vertical slices**,




integrate with respect to  $y$  and then integrate the result with respect to  $x$ .

$$\iint_T f(x, y) \, dx dy = \int_0^1 dx \int_0^{2x} f(x, y) \, dy.$$

Notice that the limits in the first integral *depend on*  $x$ .


## Double integration on regular domains

Alternatively, looking at horizontal slices, with end-points  $x = \frac{1}{2}y$ ,  $x = 1$ , and summing these from  $y = 0$  to  $y = 2$ .



## Double integration on regular domains

Alternatively, looking at horizontal slices, with end-points  $x = \frac{1}{2}y$ ,  $x = 1$ , and summing these from  $y = 0$  to  $y = 2$ .



Thus the integral is also

$$\iint_T f(x, y) \, dx dy = \int_0^2 dy \int_{\frac{1}{2}y}^1 f(x, y) \, dx.$$

# Double integration on regular domains

## Definition

Let  $D$  be a domain in the  $x, y$ -plane.  $D$  is said to be

- ▶ Type I (*y-simple*) if it is bounded by lines  $x = a$ ,  $x = b$  and curves  $y = g(x)$ ,  $y = h(x)$ , the intersection of any vertical line  $x = c$ , where  $c \in [a, b]$ , is an interval or a single point,

# Double integration on regular domains

## Definition

Let  $D$  be a domain in the  $x, y$ -plane.  $D$  is said to be

- ▶ Type I (*y-simple*) if it is bounded by lines  $x = a$ ,  $x = b$  and curves  $y = g(x)$ ,  $y = h(x)$ , the intersection of any vertical line  $x = c$ , where  $c \in [a, b]$ , is an interval or a single point,
- ▶ Type II (*x-simple*) if it is bounded by curves  $x = g(y)$ ,  $x = h(y)$  and lines  $y = a$ ,  $y = b$ , the intersection of any horizontal line  $y = c$ , where  $c \in [a, b]$ , is an interval or a single point,

# Double integration on regular domains

## Definition


Let  $D$  be a domain in the  $x, y$ -plane.  $D$  is said to be

- ▶ Type I (*y-simple*) if it is bounded by lines  $x = a$ ,  $x = b$  and curves  $y = g(x)$ ,  $y = h(x)$ , the intersection of any vertical line  $x = c$ , where  $c \in [a, b]$ , is an interval or a single point,
- ▶ Type II (*x-simple*) if it is bounded by curves  $x = g(y)$ ,  $x = h(y)$  and lines  $y = a$ ,  $y = b$ , the intersection of any horizontal line  $y = c$ , where  $c \in [a, b]$ , is an interval or a single point,
- ▶ *regular* if it the union of finitely many disjoint type I and type II domains. Every type I and type II domain is regular.




# Double integration on regular domains


## Example



Type I and not type II



Type I and type II




neither type I or type II

## Double integration on regular domains

### Example 2

State whether each of the domains shown below are type I and/or type II or regular.


(a)



(b)



(c)




## Double integration on regular domains

### Example 2

State whether each of the domains shown below are type I and/or type II or regular.


(a)



(b)



(c)



### Answers

(a) Both, (b) Type I only, (c) Neither.

## Double integration on regular domains

### Theorem

If  $D$  is the type I domain defined by  $g(x) \leq y \leq h(x)$  where  $a \leq x \leq b$  then

$$\iint_D f(x, y) \, dx dy = \int_a^b dx \int_{g(x)}^{h(x)} f(x, y) \, dy.$$

## Double integration on regular domains

### Theorem

If  $D$  is the type I domain defined by  $g(x) \leq y \leq h(x)$  where  $a \leq x \leq b$  then

$$\iint_D f(x, y) \, dx dy = \int_a^b dx \int_{g(x)}^{h(x)} f(x, y) \, dy.$$

If  $D$  is the type II domain defined by  $g(y) \leq x \leq h(y)$  where  $a \leq y \leq b$  then

$$\iint_D f(x, y) \, dx dy = \int_a^b dy \int_{g(y)}^{h(y)} f(x, y) \, dx.$$

## Double integration on regular domains

### Theorem

If  $D$  is the type I domain defined by  $g(x) \leq y \leq h(x)$  where  $a \leq x \leq b$  then

$$\iint_D f(x, y) \, dx dy = \int_a^b dx \int_{g(x)}^{h(x)} f(x, y) \, dy.$$

If  $D$  is the type II domain defined by  $g(y) \leq x \leq h(y)$  where  $a \leq y \leq b$  then

$$\iint_D f(x, y) \, dx dy = \int_a^b dy \int_{g(y)}^{h(y)} f(x, y) \, dx.$$

The *inner integral* may have a limit depending on the other variable but the *outer integral* has constant limits.

## Double integration on regular domains

### Example 3

Evaluate

$$\iint_D xy^2 \, dx dy,$$

where  $D$  is the region in the first quadrant bounded by the curve  $y = 4x^2$ , the  $x$  axis and the line  $x = 1$ .

## Double integration on regular domains



### Example 3

Evaluate

$$\iint_D xy^2 dx dy,$$

where  $D$  is the region in the first quadrant bounded by the curve  $y = 4x^2$ , the  $x$  axis and the line  $x = 1$ .

Answer





## Double integration on regular domains



### Example 3

Evaluate

$$\iint_D xy^2 dx dy,$$

where  $D$  is the region in the first quadrant bounded by the curve  $y = 4x^2$ , the  $x$  axis and the line  $x = 1$ .

Answer



$$I = \frac{8}{3}$$

## Double integration on regular domains

### Example 4

Evaluate

$$I = \iint_D 3x^2 + y^2 \, dx dy,$$

where  $D$  is the triangle with vertices  $(0, 0)$ ,  $(1, 1)$  and  $(2, 1)$ .

# Double integration on regular domains



## Example 4

Evaluate

$$I = \iint_D 3x^2 + y^2 \, dx \, dy,$$

where  $D$  is the triangle with vertices  $(0,0)$ ,  $(1,1)$  and  $(2,1)$ .

Answer



# Double integration on regular domains



## Example 4

Evaluate

$$I = \iint_D 3x^2 + y^2 \, dx \, dy,$$

where  $D$  is the triangle with vertices  $(0,0)$ ,  $(1,1)$  and  $(2,1)$ .

Answer



$$I = 2.$$

## Double integration on regular domains

### Example 5

Evaluate

$$I = \int_0^1 dx \int_{\sqrt{x}}^1 \frac{e^{y^2}}{\sqrt{x}} dy.$$



# Double integration on regular domains

## Example 5

Evaluate

$$I = \int_0^1 dx \int_{\sqrt{x}}^1 \frac{e^{y^2}}{\sqrt{x}} dy.$$

Answer





# Double integration on regular domains

## Example 5

Evaluate

$$I = \int_0^1 dx \int_{\sqrt{x}}^1 \frac{e^{y^2}}{\sqrt{x}} dy.$$

Answer



$$I = e - 1.$$

## Double integration on regular domains

### Example 6

Find the volume of the tetrahedron  $T$ , bounded by the planes  $x + 2y + z = 2$ ,  $x = 2y$ ,  $x = 0$  and  $z = 0$ .





# Double integration on regular domains

## Example 6

Find the volume of the tetrahedron  $T$ , bounded by the planes  $x + 2y + z = 2$ ,  $x = 2y$ ,  $x = 0$  and  $z = 0$ .

## Answer





# Double integration on regular domains

## Example 6

Find the volume of the tetrahedron  $T$ , bounded by the planes  $x + 2y + z = 2$ ,  $x = 2y$ ,  $x = 0$  and  $z = 0$ .

## Answer



$$T = \frac{1}{3}.$$

## Double integration in polar coordinates

The position of a point  $(x, y)$  on the cartesian plane can be specified by  $r, \theta$  which are

Polar coordinates


$$x = r \cos \theta, \quad y = r \sin \theta, \quad \theta \in [0, 2\pi), \quad r \geq 0.$$

## Double integration in polar coordinates

The position of a point  $(x, y)$  on the cartesian plane can be specified by  $r, \theta$  which are

Polar coordinates

$$x = r \cos \theta, \quad y = r \sin \theta, \quad \theta \in [0, 2\pi), \quad r \geq 0.$$




## Double integration in polar coordinates

In cartesian coordinates, the area of an elementary rectangle using in the Riemann sum is  $\delta A = \delta x \delta y$ . In polar coordinates, the area element has area  $\delta A \approx r \delta r \delta \theta$ .

## Double integration in polar coordinates

In cartesian coordinates, the area of an elementary rectangle using in the Riemann sum is  $\delta A = \delta x \delta y$ . In polar coordinates, the area element has area  $\delta A \approx r \delta r \delta \theta$ .




For this reason in polar coordinates,  $dA = r dr d\theta$ , i.e.,

$$\iint_D f(x, y) dx dy = \iint_D f(r \cos \theta, r \sin \theta) r dr d\theta.$$

## Double integration in polar coordinates

In cartesian coordinates, the area of an elementary rectangle using in the Riemann sum is  $\delta A = \delta x \delta y$ . In polar coordinates, the area element has area  $\delta A \approx r \delta r \delta \theta$ .



For this reason in polar coordinates,  $dA = r dr d\theta$ , i.e.,

$$\iint_D f(x, y) dx dy = \iint_D f(r \cos \theta, r \sin \theta) r dr d\theta.$$

When either the domain is circular or the integrand is written in terms of  $x^2 + y^2 (= r^2)$ , use polar coordinates.

## Double integration in polar coordinates

### Example 7

Use polar coordinates to evaluate

$$I = \iint_D x + y \, dx dy,$$

where  $D$  is part of the annulus between circles of radius 1 and 2, centre  $(0, 0)$  lying in upper half plane.



## Double integration in polar coordinates


### Example 7

Use polar coordinates to evaluate

$$I = \iint_D x + y \, dx dy,$$

where  $D$  is part of the annulus between circles of radius 1 and 2, centre  $(0,0)$  lying in upper half plane.

Answer



## Double integration in polar coordinates


### Example 7

Use polar coordinates to evaluate

$$I = \iint_D x + y \, dx dy,$$

where  $D$  is part of the annulus between circles of radius 1 and 2, centre  $(0,0)$  lying in upper half plane.

Answer



$$I = \frac{14}{3}.$$

## Double integration in polar coordinates

### Example 8

Evaluate

$$I = \iint_D y \, dx dy,$$

where  $D$  is the part of the disk of radius  $a (> 0)$  and centre  $(a, 0)$  lying in the first quadrant.

# Double integration in polar coordinates


## Example 8

Evaluate

$$I = \iint_D y \, dx dy,$$

where  $D$  is the part of the disk of radius  $a$  ( $> 0$ ) and centre  $(a, 0)$  lying in the first quadrant.

Answer



# Double integration in polar coordinates


## Example 8

Evaluate

$$I = \iint_D y \, dx dy,$$

where  $D$  is the part of the disk of radius  $a$  ( $> 0$ ) and centre  $(a, 0)$  lying in the first quadrant.

Answer



$$I = \frac{2a^3}{3}.$$

## Beta and Gamma functions

Beta functions can help us easily integrate functions that involve powers of cosine and sine.

► *Beta function:*

$$B(p, q) = \int_0^1 x^{p-1} (1-x)^{q-1} dx, \quad p > 0 \text{ and } q > 0.$$

A particularly useful form is,

$$B(p, q) = 2 \int_0^{\pi/2} \sin^{2p-1}(y) \cos^{2q-1}(y) dy .$$

This is found by substituting  $x = \sin^2 y$  in the definition of the Beta function.

## Beta and Gamma functions

Beta functions can help us easily integrate functions that involve powers of cosine and sine.

► *Beta function:*

$$B(p, q) = \int_0^1 x^{p-1} (1-x)^{q-1} dx, \quad p > 0 \text{ and } q > 0.$$

A particularly useful form is,

$$B(p, q) = 2 \int_0^{\pi/2} \sin^{2p-1}(y) \cos^{2q-1}(y) dy.$$

This is found by substituting  $x = \sin^2 y$  in the definition of the Beta function.

► *Gamma function:*

$$\Gamma(k) = \int_0^{\infty} x^{k-1} e^{-x} dx, \quad k > 0.$$

## Properties of Beta and Gamma functions

1.  $\Gamma(1) = 1$ ,  $\Gamma(2) = 1$ ,  $\Gamma(3) = 2$  and in general  $\Gamma(n) = (n-1)!$  for every positive integer  $n$ .



## Properties of Beta and Gamma functions

1.  $\Gamma(1) = 1$ ,  $\Gamma(2) = 1$ ,  $\Gamma(3) = 2$  and in general  $\Gamma(n) = (n-1)!$  for every positive integer  $n$ .
2.  $\Gamma(k) = (k-1)\Gamma(k-1)$  for all real numbers  $k > 1$  Repeatedly applying the formula for  $\Gamma(k)$  gives a formula in terms of  $\Gamma(k-p)$ , where  $0 < k-p \leq 1$ , e.g.  $\Gamma(9/4) = \frac{5}{4}\frac{1}{4}\Gamma(\frac{1}{4})$ .

## Properties of Beta and Gamma functions

1.  $\Gamma(1) = 1$ ,  $\Gamma(2) = 1$ ,  $\Gamma(3) = 2$  and in general  $\Gamma(n) = (n-1)!$  for every positive integer  $n$ .
2.  $\Gamma(k) = (k-1)\Gamma(k-1)$  for all real numbers  $k > 1$  Repeatedly applying the formula for  $\Gamma(k)$  gives a formula in terms of  $\Gamma(k-p)$ , where  $0 < k-p \leq 1$ , e.g.  $\Gamma(9/4) = \frac{5}{4}\frac{1}{4}\Gamma(\frac{1}{4})$ .
3. For  $0 < k < 1$ ,  $\Gamma(k)\Gamma(1-k) = \pi / \sin(k\pi)$ .

## Properties of Beta and Gamma functions

1.  $\Gamma(1) = 1$ ,  $\Gamma(2) = 1$ ,  $\Gamma(3) = 2$  and in general  $\Gamma(n) = (n-1)!$  for every positive integer  $n$ .
2.  $\Gamma(k) = (k-1)\Gamma(k-1)$  for all real numbers  $k > 1$  Repeatedly applying the formula for  $\Gamma(k)$  gives a formula in terms of  $\Gamma(k-p)$ , where  $0 < k-p \leq 1$ , e.g.  $\Gamma(9/4) = \frac{5}{4}\frac{1}{4}\Gamma(\frac{1}{4})$ .
3. For  $0 < k < 1$ ,  $\Gamma(k)\Gamma(1-k) = \pi/\sin(k\pi)$ .
4.  $\Gamma(1/2) = \sqrt{\pi}$ .

## Properties of Beta and Gamma functions

1.  $\Gamma(1) = 1$ ,  $\Gamma(2) = 1$ ,  $\Gamma(3) = 2$  and in general  $\Gamma(n) = (n-1)!$  for every positive integer  $n$ .
2.  $\Gamma(k) = (k-1)\Gamma(k-1)$  for all real numbers  $k > 1$  Repeatedly applying the formula for  $\Gamma(k)$  gives a formula in terms of  $\Gamma(k-p)$ , where  $0 < k-p \leq 1$ , e.g.  $\Gamma(9/4) = \frac{5}{4}\frac{1}{4}\Gamma(\frac{1}{4})$ .
3. For  $0 < k < 1$ ,  $\Gamma(k)\Gamma(1-k) = \pi/\sin(k\pi)$ .
4.  $\Gamma(1/2) = \sqrt{\pi}$ .
5.  $B(p, q) = \frac{\Gamma(p)\Gamma(q)}{\Gamma(p+q)}$ .

# Properties of Beta and Gamma functions

## Result

From the properties of Gamma functions we can derive the following result:

### Property of Beta functions

$$\int_0^{\pi/2} \sin^m x \cos^n x \, dx = \frac{(m-1)(m-3)\dots(n-1)(n-3)\dots}{(m+n)(m+n-2)(m+n-4)\dots} K$$

where  $K = 1$  unless  $m$  and  $n$  are both even in which case  $K = \pi/2$ .

# Properties of Beta and Gamma functions

## Result

From the properties of Gamma functions we can derive the following result:

### Property of Beta functions

$$\int_0^{\pi/2} \sin^m x \cos^n x \, dx = \frac{(m-1)(m-3)\dots(n-1)(n-3)\dots}{(m+n)(m+n-2)(m+n-4)\dots} K$$

where  $K = 1$  unless  $m$  and  $n$  are both even in which case  $K = \pi/2$ .

In the special cases,  $m = 0$  or  $m = 1$  none of the numerator factors involving  $m$  appear.

# Properties of Beta and Gamma functions

## Result

From the properties of Gamma functions we can derive the following result:

### Property of Beta functions

$$\int_0^{\pi/2} \sin^m x \cos^n x \, dx = \frac{(m-1)(m-3)\dots(n-1)(n-3)\dots}{(m+n)(m+n-2)(m+n-4)\dots} K$$

where  $K = 1$  unless  $m$  and  $n$  are both even in which case  $K = \pi/2$ .

In the special cases,  $m = 0$  or  $m = 1$  none of the numerator factors involving  $m$  appear.

For example,

$$\int_0^{\pi/2} \sin^3 x \cos^6 x \, dx = \frac{2 \cdot 5 \cdot 3 \cdot 1}{9 \cdot 7 \cdot 5 \cdot 3 \cdot 1} = \frac{2}{63}.$$




## Simplifying sine and cosine integrals

Properties of the graphs of sine and cosine seen in 1S/X simplify the integral before applying Beta functions.






## Simplifying sine and cosine integrals

Properties of the graphs of sine and cosine seen in 1S/X simplify the integral before applying Beta functions.



## Simplifying sine and cosine integrals

Properties of the graphs of sine and cosine seen in 1S/X simplify the integral before applying Beta functions.



We deduce

$$\int_0^\pi \sin x \, dx = 2 \int_0^{\pi/2} \sin x \, dx; \quad \int_0^{2\pi} \sin x \, dx = 0; \quad \int_0^\pi \cos x \, dx = 0;$$

$$\int_0^\pi \sin^2 x \, dx = 2 \int_0^{\pi/2} \sin^2 x \, dx; \quad \int_0^\pi \cos^2 x \, dx = 2 \int_0^{\pi/2} \cos^2 x \, dx \dots \text{et}$$

# Beta functions

## Example 9

Evaluate:

$$(a) I = \int_0^{\pi} \sin^3 x \cos^4 x \, dx, \quad (b) I = \int_0^{\pi} \sin^3 x \cos^5 x \, dx,$$

$$(c) I = \int_0^{2\pi} \sin^2 x \cos^4 x \, dx.$$

# Beta functions

## Example 9

Evaluate:

$$(a) I = \int_0^{\pi} \sin^3 x \cos^4 x \, dx, \quad (b) I = \int_0^{\pi} \sin^3 x \cos^5 x \, dx,$$

$$(c) I = \int_0^{2\pi} \sin^2 x \cos^4 x \, dx.$$

## Answers

$$(a) I = \frac{4}{35}, \quad (b) I = 0, \quad (c) I = \frac{\pi}{8}.$$

# Change of variables in double integration

## Definition

Consider a change of variables  $x, y$  to  $u, v$ . So  $x = x(u, v)$  and  $y = y(u, v)$ . The *Jacobian*  $\frac{\partial(u, v)}{\partial(x, y)}$  is the determinant

$$\begin{vmatrix} u_x & u_y \\ v_x & v_y \end{vmatrix} = u_x v_y - u_y v_x.$$

# Change of variables in double integration

## Definition

Consider a change of variables  $x, y$  to  $u, v$ . So  $x = x(u, v)$  and  $y = y(u, v)$ . The *Jacobian*  $\frac{\partial(u, v)}{\partial(x, y)}$  is the determinant

$$\begin{vmatrix} u_x & u_y \\ v_x & v_y \end{vmatrix} = u_x v_y - u_y v_x.$$

If the change of variables is invertible then the Jacobian is nonzero and

$$\frac{\partial(x, y)}{\partial(u, v)} = 1 \bigg/ \frac{\partial(u, v)}{\partial(x, y)}.$$

# Change of variables in double integration

## Theorem

Let the change of variables  $x, y$  to  $u, v$  be invertible on the domain  $D$ . Then

$$\iint_D f(x, y) \, dx dy = \iint_S f(x(u, v), y(u, v)) \left| \frac{\partial(x, y)}{\partial(u, v)} \right| \, du dv,$$

where  $D$  is the domain in the  $xy$ -plane and  $S$  is the corresponding domain in the  $uv$ -plane.

## Change of variables in double integration

### Theorem

Let the change of variables  $x, y$  to  $u, v$  be invertible on the domain  $D$ . Then

$$\iint_D f(x, y) \, dx \, dy = \iint_S f(x(u, v), y(u, v)) \left| \frac{\partial(x, y)}{\partial(u, v)} \right| \, du \, dv,$$

where  $D$  is the domain in the  $xy$ -plane and  $S$  is the corresponding domain in the  $uv$ -plane.


Often it is convenient to use

$$\frac{\partial(x, y)}{\partial(u, v)} = 1 \bigg/ \frac{\partial(u, v)}{\partial(x, y)}$$




## Change of variables in double integration

- The idea here is to choose variables  $u, v$  in which the domain is simply described, preferably with constant limits, e.g.




## Change of variables in double integration

- The idea here is to choose variables  $u, v$  in which the domain is simply described, preferably with constant limits, e.g.



## Change of variables in double integration

- ▶ The idea here is to choose variables  $u, v$  in which the domain is simply described, preferably with constant limits, e.g.




The lines  $u = u_0$  and  $v = v_0$  in the  $uv$ -plane get mapped to curves  $x = x(u_0, v)$ ,  $y = y(u_0, v)$  and  $x = x(u, v_0)$ ,  $y = y(u, v_0)$  in the  $xy$ -plane.

- ▶ The idea behind the proof of the theorem is to show that the area element is  $\left| \frac{\partial(x, y)}{\partial(u, v)} \right| du dv$ .

## Change of variables in double integration

- ▶ The idea here is to choose variables  $u, v$  in which the domain is simply described, preferably with constant limits, e.g.




The lines  $u = u_0$  and  $v = v_0$  in the  $uv$ -plane get mapped to curves  $x = x(u_0, v)$ ,  $y = y(u_0, v)$  and  $x = x(u, v_0)$ ,  $y = y(u, v_0)$  in the  $xy$ -plane.

- ▶ The idea behind the proof of the theorem is to show that the area element is  $\left| \frac{\partial(x, y)}{\partial(u, v)} \right| du dv$ .

## Change of variables in double integration

- ▶ The idea here is to choose variables  $u, v$  in which the domain is simply described, preferably with constant limits, e.g.



The lines  $u = u_0$  and  $v = v_0$  in the  $uv$ -plane get mapped to curves  $x = x(u_0, v)$ ,  $y = y(u_0, v)$  and  $x = x(u, v_0)$ ,  $y = y(u, v_0)$  in the  $xy$ -plane.

- ▶ The idea behind the proof of the theorem is to show that the area element is  $\left| \frac{\partial(x, y)}{\partial(u, v)} \right| du dv$ .

## Change of variables in double integration

- ▶ Let  $\mathbf{r} = (x, y)$ . The tangent vector at  $(x_0, y_0)$  to the curve  $(x(u, v_0), y(u, v_0))$  is  $\mathbf{r}_u = \left( \frac{\partial x}{\partial u}, \frac{\partial y}{\partial u} \right)$ .

## Change of variables in double integration

- ▶ Let  $\mathbf{r} = (x, y)$ . The tangent vector at  $(x_0, y_0)$  to the curve  $(x(u, v_0), y(u, v_0))$  is  $\mathbf{r}_u = \left( \frac{\partial x}{\partial u}, \frac{\partial y}{\partial u} \right)$ .
- ▶ The tangent vector at  $(x_0, y_0)$  to the curve  $(x(u_0, v), y(u_0, v))$  is  $\mathbf{r}_v = \left( \frac{\partial x}{\partial v}, \frac{\partial y}{\partial v} \right)$ .

## Change of variables in double integration

- ▶ Let  $\mathbf{r} = (x, y)$ . The tangent vector at  $(x_0, y_0)$  to the curve  $(x(u, v_0), y(u, v_0))$  is  $\mathbf{r}_u = \left( \frac{\partial x}{\partial u}, \frac{\partial y}{\partial u} \right)$ .
- ▶ The tangent vector at  $(x_0, y_0)$  to the curve  $(x(u_0, v), y(u_0, v))$  is  $\mathbf{r}_v = \left( \frac{\partial x}{\partial v}, \frac{\partial y}{\partial v} \right)$ .
- ▶ The area of the small element of  $D$  is  $\Delta A$  and is approximated by the area of the parallelogram which is given by the magnitude of the following cross product

$$\Delta u \mathbf{r}_u \times \Delta v \mathbf{r}_v = \Delta u \Delta v (x_u y_v - x_v y_u) \mathbf{k} = \Delta u \Delta v \frac{\partial(x, y)}{\partial(u, v)} \mathbf{k}.$$





## Change of variables in double integration

- ▶ Let  $\mathbf{r} = (x, y)$ . The tangent vector at  $(x_0, y_0)$  to the curve  $(x(u, v_0), y(u, v_0))$  is  $\mathbf{r}_u = \left( \frac{\partial x}{\partial u}, \frac{\partial y}{\partial u} \right)$ .
- ▶ The tangent vector at  $(x_0, y_0)$  to the curve  $(x(u_0, v), y(u_0, v))$  is  $\mathbf{r}_v = \left( \frac{\partial x}{\partial v}, \frac{\partial y}{\partial v} \right)$ .
- ▶ The area of the small element of  $D$  is  $\Delta A$  and is approximated by the area of the parallelogram which is given by the magnitude of the following cross product

$$\Delta u \mathbf{r}_u \times \Delta v \mathbf{r}_v = \Delta u \Delta v (x_u y_v - x_v y_u) \mathbf{k} = \Delta u \Delta v \frac{\partial(x, y)}{\partial(u, v)} \mathbf{k}.$$



## Change of variables in double integration

- ▶ Let  $\mathbf{r} = (x, y)$ . The tangent vector at  $(x_0, y_0)$  to the curve  $(x(u, v_0), y(u, v_0))$  is  $\mathbf{r}_u = \left( \frac{\partial x}{\partial u}, \frac{\partial y}{\partial u} \right)$ .
- ▶ The tangent vector at  $(x_0, y_0)$  to the curve  $(x(u_0, v), y(u_0, v))$  is  $\mathbf{r}_v = \left( \frac{\partial x}{\partial v}, \frac{\partial y}{\partial v} \right)$ .
- ▶ The area of the small element of  $D$  is  $\Delta A$  and is approximated by the area of the parallelogram which is given by the magnitude of the following cross product

$$\Delta u \mathbf{r}_u \times \Delta v \mathbf{r}_v = \Delta u \Delta v (x_u y_v - x_v y_u) \mathbf{k} = \Delta u \Delta v \frac{\partial(x, y)}{\partial(u, v)} \mathbf{k}.$$



## Change of variables in double integration

- Summing the elements that make up the region  $D$

$$\begin{aligned}\iint_D f(x, y) \, dx dy &= \lim_{N, M \rightarrow \infty} \sum_{i=1}^N \sum_{j=1}^M f(x_i, y_j) \Delta A \\ &= \lim_{N, M \rightarrow \infty} \sum_{i=1}^N \sum_{j=1}^M f(x(u_i, v_j), y(u_i, v_j)) \left| \frac{\partial(x, y)}{\partial(u, v)} \right| \Delta u \Delta v\end{aligned}$$

which is the Riemann sum for  $\iint_S f(x(u, v), y(u, v)) |J| \, du dv$ .

## Change of variables in double integration

- ▶ Summing the elements that make up the region  $D$

$$\begin{aligned}\iint_D f(x, y) \, dx dy &= \lim_{N, M \rightarrow \infty} \sum_{i=1}^N \sum_{j=1}^M f(x_i, y_j) \Delta A \\ &= \lim_{N, M \rightarrow \infty} \sum_{i=1}^N \sum_{j=1}^M f(x(u_i, v_j), y(u_i, v_j)) \left| \frac{\partial(x, y)}{\partial(u, v)} \right| \Delta u \Delta v\end{aligned}$$

which is the Riemann sum for  $\iint_S f(x(u, v), y(u, v)) |J| \, du dv$ .

- ▶ For the change to polar coordinates,

$$J = \begin{vmatrix} (r \cos \theta)_r & (r \cos \theta)_\theta \\ (r \sin \theta)_r & (r \sin \theta)_\theta \end{vmatrix} = \begin{vmatrix} \cos \theta & -r \sin \theta \\ \sin \theta & r \cos \theta \end{vmatrix} = r(\cos^2 \theta + \sin^2 \theta) = r,$$

## Change of variables in double integration

- ▶ Summing the elements that make up the region  $D$

$$\begin{aligned}\iint_D f(x, y) \, dx dy &= \lim_{N, M \rightarrow \infty} \sum_{i=1}^N \sum_{j=1}^M f(x_i, y_j) \Delta A \\ &= \lim_{N, M \rightarrow \infty} \sum_{i=1}^N \sum_{j=1}^M f(x(u_i, v_j), y(u_i, v_j)) \left| \frac{\partial(x, y)}{\partial(u, v)} \right| \Delta u \Delta v\end{aligned}$$

which is the Riemann sum for  $\iint_S f(x(u, v), y(u, v)) |J| \, du dv$ .

- ▶ For the change to polar coordinates,

$$J = \begin{vmatrix} (r \cos \theta)_r & (r \cos \theta)_\theta \\ (r \sin \theta)_r & (r \sin \theta)_\theta \end{vmatrix} = \begin{vmatrix} \cos \theta & -r \sin \theta \\ \sin \theta & r \cos \theta \end{vmatrix} = r(\cos^2 \theta + \sin^2 \theta) = r,$$

## Change of variables in double integration

- ▶ Summing the elements that make up the region  $D$

$$\begin{aligned}\iint_D f(x, y) \, dx dy &= \lim_{N, M \rightarrow \infty} \sum_{i=1}^N \sum_{j=1}^M f(x_i, y_j) \Delta A \\ &= \lim_{N, M \rightarrow \infty} \sum_{i=1}^N \sum_{j=1}^M f(x(u_i, v_j), y(u_i, v_j)) \left| \frac{\partial(x, y)}{\partial(u, v)} \right| \Delta u \Delta v\end{aligned}$$

which is the Riemann sum for  $\iint_S f(x(u, v), y(u, v)) |J| \, du dv$ .

- ▶ For the change to polar coordinates,

$$J = \begin{vmatrix} (r \cos \theta)_r & (r \cos \theta)_\theta \\ (r \sin \theta)_r & (r \sin \theta)_\theta \end{vmatrix} = \begin{vmatrix} \cos \theta & -r \sin \theta \\ \sin \theta & r \cos \theta \end{vmatrix} = r(\cos^2 \theta + \sin^2 \theta) = r,$$

giving the result

$$\iint_D f(x, y) \, dx dy = \iint_S f(r \cos \theta, r \sin \theta) r \, dr d\theta.$$

## Change of variables in double integration

### Example 10

By making a suitable change of variables, evaluate

$$\iint_D x + 3y \, dx dy,$$

where  $D$  is the region bounded by the lines

$$y = x - 1, \quad y = x + 1, \quad y = -x - 1, \quad y = -x + 3.$$

# Change of variables in double integration

## Example 10



By making a suitable change of variables, evaluate

$$\iint_D x + 3y \, dx dy,$$

where  $D$  is the region bounded by the lines

$$y = x - 1, \quad y = x + 1, \quad y = -x - 1, \quad y = -x + 3.$$

## Answer





# Change of variables in double integration

## Example 10



By making a suitable change of variables, evaluate

$$\iint_D x + 3y \, dx dy,$$

where  $D$  is the region bounded by the lines

$$y = x - 1, \quad y = x + 1, \quad y = -x - 1, \quad y = -x + 3.$$

## Answer



$$I = 8.$$

# Change of variables in double integration

## Example 11

Find the area bounded by the curves  $y = e^x$ ,  $y = 2e^x$ ,  $y = e^{-x}$  and  $y = 2e^{-x}$ .

# Change of variables in double integration



## Example 11

Find the area bounded by the curves  $y = e^x$ ,  $y = 2e^x$ ,  $y = e^{-x}$  and  $y = 2e^{-x}$ .

The area of a surface  $A \subset \mathbb{R}^2$  is given by the double integral

$$\iint_A 1 \, dx dy.$$

Answer



# Change of variables in double integration



## Example 11

Find the area bounded by the curves  $y = e^x$ ,  $y = 2e^x$ ,  $y = e^{-x}$  and  $y = 2e^{-x}$ .

The area of a surface  $A \subset \mathbb{R}^2$  is given by the double integral

$$\iint_A 1 \, dx dy.$$

Answer



$$\text{Area} = 2(3 - 2\sqrt{2}).$$

# Triple integration

Define triple integrals for functions of three variables.

## Triple integration

Define triple integrals for functions of three variables. Recall the definition of a double integral is given by

$$\iint_R f(x, y) \, dx dy = \lim_{N, M \rightarrow \infty} \sum_{i=1}^N \sum_{j=1}^M f(x_i, y_j) \delta x_i \delta y_j.$$

## Triple integration

Define triple integrals for functions of three variables. Recall the definition of a double integral is given by

$$\iint_R f(x, y) \, dx dy = \lim_{N, M \rightarrow \infty} \sum_{i=1}^N \sum_{j=1}^M f(x_i, y_j) \delta x_i \delta y_j.$$

For a triple integral instead of summing over an area  $\delta A_{ij} = \delta x_i \delta y_j$ , we sum over a volume  $\delta V_{ijk} = \delta x_i \delta y_j \delta z_k$  which leads us to


$$\iiint_V f(x, y, z) \, dx dy dz = \lim_{N, M, L \rightarrow \infty} \sum_{i=1}^N \sum_{j=1}^M \sum_{k=1}^L f(x_i, y_j, z_k) \delta x_i \delta y_j \delta z_k.$$

## Triple integration

- If  $V$  lies between two continuous functions of  $x$  and  $y$  then

$$\iiint_V f(x, y, z) \, dx dy dz = \iint_D \left( \int_{u_1(x, y)}^{u_2(x, y)} f(x, y, z) \, dz \right) dx dy$$

where  $D$  is the projection of  $V$  onto the  $xy$  plane.






## Triple integration

- ▶ If  $V$  lies between two continuous functions of  $x$  and  $y$  then

$$\iiint_V f(x, y, z) \, dx dy dz = \iint_D \left( \int_{u_1(x, y)}^{u_2(x, y)} f(x, y, z) \, dz \right) dx dy$$

where  $D$  is the projection of  $V$  onto the  $xy$  plane.




## Triple integration

- ▶ If  $V$  lies between two continuous functions of  $x$  and  $y$  then

$$\iiint_V f(x, y, z) \, dx dy dz = \iint_D \left( \int_{u_1(x, y)}^{u_2(x, y)} f(x, y, z) \, dz \right) dx dy$$

where  $D$  is the projection of  $V$  onto the  $xy$  plane.



## Triple integration

- ▶ If  $V$  lies between two functions of  $y$  and  $z$  then

$$\iiint_V f(x, y, z) \, dx dy dz = \iint_D \left( \int_{u_1(y, z)}^{u_2(y, z)} f(x, y, z) \, dx \right) dy dz,$$

where  $D$  is the projection of  $V$  onto the  $yz$  plane.

## Triple integration

- ▶ If  $V$  lies between two functions of  $y$  and  $z$  then

$$\iiint_V f(x, y, z) \, dx dy dz = \iint_D \left( \int_{u_1(y, z)}^{u_2(y, z)} f(x, y, z) \, dx \right) dy dz,$$

where  $D$  is the projection of  $V$  onto the  $yz$  plane.

- ▶ Lastly if  $V$  lies between two functions of  $x$  and  $z$  then

$$\iiint_V f(x, y, z) \, dx dy dz = \iint_D \left( \int_{u_1(x, z)}^{u_2(x, z)} f(x, y, z) \, dy \right) dx dz,$$

where  $D$  is the projection of  $V$  onto the  $xz$  plane.

## Triple integration

- ▶ If  $V$  lies between two functions of  $y$  and  $z$  then

$$\iiint_V f(x, y, z) \, dx dy dz = \iint_D \left( \int_{u_1(y, z)}^{u_2(y, z)} f(x, y, z) \, dx \right) dy dz,$$

where  $D$  is the projection of  $V$  onto the  $yz$  plane.

- ▶ Lastly if  $V$  lies between two functions of  $x$  and  $z$  then

$$\iiint_V f(x, y, z) \, dx dy dz = \iint_D \left( \int_{u_1(x, z)}^{u_2(x, z)} f(x, y, z) \, dy \right) dx dz,$$

where  $D$  is the projection of  $V$  onto the  $xz$  plane.

## Triple integration

- ▶ If  $V$  lies between two functions of  $y$  and  $z$  then

$$\iiint_V f(x, y, z) \, dx dy dz = \iint_D \left( \int_{u_1(y, z)}^{u_2(y, z)} f(x, y, z) \, dx \right) dy dz,$$

where  $D$  is the projection of  $V$  onto the  $yz$  plane.

- ▶ Lastly if  $V$  lies between two functions of  $x$  and  $z$  then

$$\iiint_V f(x, y, z) \, dx dy dz = \iint_D \left( \int_{u_1(x, z)}^{u_2(x, z)} f(x, y, z) \, dy \right) dx dz,$$

where  $D$  is the projection of  $V$  onto the  $xz$  plane.

# Triple integration

In general if  $V$  lies between two continuous functions of  $x$  and  $y$  then

Triple integral

$$\iiint_V f(x, y, z) \, dx dy dz = \underbrace{\int_a^b}_{\text{Constants}} dx \underbrace{\int_{h_1(x)}^{h_2(x)}}_{\text{Curves}} dy \underbrace{\int_{g_1(x,y)}^{g_2(x,y)}}_{\text{Surfaces}} f(x, y, z) \, dz.$$

## Triple integration

- ▶ The *volume of a solid*  $V \subset \mathbb{R}^3$  is given by the following triple integral

$$\iiint_V 1 \, dx dy dz.$$



## Triple integration

- ▶ The *volume* of a solid  $V \subset \mathbb{R}^3$  is given by the following triple integral

$$\iiint_V 1 \, dx dy dz.$$

- ▶ The *mass* of the solid  $V$ , where the *density*  $f(x, y, z)$  of the solid varies across  $V$  is

$$\iiint_V f(x, y, z) \, dx dy dz.$$

## Triple integration

- ▶ The *volume* of a solid  $V \subset \mathbb{R}^3$  is given by the following triple integral

$$\iiint_V 1 \, dx dy dz.$$

- ▶ The *mass* of the solid  $V$ , where the *density*  $f(x, y, z)$  of the solid varies across  $V$  is

$$\iiint_V f(x, y, z) \, dx dy dz.$$

## Triple integration

- ▶ The *volume* of a solid  $V \subset \mathbb{R}^3$  is given by the following triple integral

$$\iiint_V 1 \, dx dy dz.$$

- ▶ The *mass* of the solid  $V$ , where the *density*  $f(x, y, z)$  of the solid varies across  $V$  is

$$\iiint_V f(x, y, z) \, dx dy dz.$$

# Triple integration

## Example 12

Evaluate

$$I = \iiint_V z \, dx dy dz,$$

where  $V$  is the solid tetrahedron bounded by the four planes  $x = 0$ ,  $y = 0$ ,  $z = 0$  and  $x + y + z = 1$ .

# Triple integration



## Example 12

Evaluate

$$I = \iiint_V z \, dx \, dy \, dz,$$

where  $V$  is the solid tetrahedron bounded by the four planes  $x = 0$ ,  $y = 0$ ,  $z = 0$  and  $x + y + z = 1$ .

Answer



# Triple integration



## Example 12

Evaluate

$$I = \iiint_V z \, dx \, dy \, dz,$$

where  $V$  is the solid tetrahedron bounded by the four planes  $x = 0$ ,  $y = 0$ ,  $z = 0$  and  $x + y + z = 1$ .

Answer



# Triple integration

## Example 13



Set up (but do not evaluate) the integral for the volume of the solid that lies below the paraboloid  $z = 9 - x^2 - y^2$  and above the plane  $z = 5$ .

# Triple integration

## Example 13

Set up (but do not evaluate) the integral for the volume of the solid that lies below the paraboloid  $z = 9 - x^2 - y^2$  and above the plane  $z = 5$ .

## Answer







# Triple integration

## Example 13

Set up (but do not evaluate) the integral for the volume of the solid that lies below the paraboloid  $z = 9 - x^2 - y^2$  and above the plane  $z = 5$ .

## Answer



## Triple integration in spherical coordinates

The position of a point  $(x, y, z)$  in cartesian coordinates can be specified by  $\rho$ ,  $\theta$ ,  $\phi$  which are

Spherical coordinates

$$x = \rho \sin \phi \cos \theta, \quad y = \rho \sin \phi \sin \theta, \quad z = \rho \cos \phi.$$

$$\theta \in [0, 2\pi), \quad \phi \in [0, \pi), \quad \rho \geq 0.$$


## Triple integration in spherical coordinates

The position of a point  $(x, y, z)$  in cartesian coordinates can be specified by  $\rho$ ,  $\theta$ ,  $\phi$  which are

Spherical coordinates

$$x = \rho \sin \phi \cos \theta, \quad y = \rho \sin \phi \sin \theta, \quad z = \rho \cos \phi.$$

$$\theta \in [0, 2\pi), \quad \phi \in [0, \pi), \quad \rho \geq 0.$$




## Triple integration in spherical coordinates

In cartesian coordinates, the volume of an elementary cuboid used in the Riemann sum is  $\delta V = \delta x \delta y \delta z$ . In spherical coordinates, the volume element is  $\delta V \approx \rho^2 \sin \phi \delta \theta \delta \phi \delta \rho$ .


## Triple integration in spherical coordinates

In cartesian coordinates, the volume of an elementary cuboid used in the Riemann sum is  $\delta V = \delta x \delta y \delta z$ . In spherical coordinates, the volume element is  $\delta V \approx \rho^2 \sin \phi \delta \theta \delta \phi \delta \rho$ .



## Triple integration in spherical coordinates

In cartesian coordinates, the volume of an elementary cuboid used in the Riemann sum is  $\delta V = \delta x \delta y \delta z$ . In spherical coordinates, the volume element is  $\delta V \approx \rho^2 \sin \phi \delta \theta \delta \phi \delta \rho$ .



$$\iiint_V f(x, y, z) dx dy dz =$$

$$\iiint_V f(\rho \sin \phi \cos \theta, \rho \sin \phi \sin \theta, \rho \cos \phi) \rho^2 \sin \phi d\theta d\phi d\rho.$$

When either the domain is spherical or the integrand is written in terms of  $x^2 + y^2 + z^2 (= \rho^2)$ , use spherical coordinates.

## Triple integration in spherical coordinates

### Example 14

Use spherical coordinates to evaluate

$$I = \iiint_B \exp((x^2 + y^2 + z^2)^{3/2}) \, dx \, dy \, dz,$$

where  $B$  is the unit ball,  $B = \{(x, y, z) | x^2 + y^2 + z^2 \leq 1\}$ .

## Triple integration in spherical coordinates

### Example 14

Use spherical coordinates to evaluate

$$I = \iiint_B \exp((x^2 + y^2 + z^2)^{3/2}) \, dx \, dy \, dz,$$

where  $B$  is the unit ball,  $B = \{(x, y, z) | x^2 + y^2 + z^2 \leq 1\}$ .

Answer

$$I = \frac{4}{3}\pi(e - 1).$$



## Triple integration in spherical coordinates

### Example 15


Find the volume of the solid that lies above the cone  $z = \sqrt{x^2 + y^2}$  and below the sphere  $x^2 + y^2 + z^2 = z$ .

# Triple integration in spherical coordinates

## Example 15

Find the volume of the solid that lies above the cone  $z = \sqrt{x^2 + y^2}$  and below the sphere  $x^2 + y^2 + z^2 = z$ .

Answer




# Triple integration in spherical coordinates

## Example 15

Find the volume of the solid that lies above the cone  $z = \sqrt{x^2 + y^2}$  and below the sphere  $x^2 + y^2 + z^2 = z$ .

Answer



$$V = \frac{\pi}{8}.$$

## Chapter 3: Differentiation of vectors

- ▶ Scalar- and vector-valued functions

## Chapter 3: Differentiation of vectors

- ▶ Scalar- and vector-valued functions
- ▶ **vector and scalar fields**

## Chapter 3: Differentiation of vectors

- ▶ Scalar- and vector-valued functions
- ▶ vector and scalar fields
- ▶ types of derivative—grad, div and curl

## Scalar- and vector-valued functions

- ▶ In Chapters 1 and 2 we considered functions of several variables  $f : D \rightarrow \mathbb{R}$  where  $D \subset \mathbb{R}^n$

## Scalar- and vector-valued functions

- ▶ In Chapters 1 and 2 we considered functions of several variables  $f : D \rightarrow \mathbb{R}$  where  $D \subset \mathbb{R}^n$
- ▶ these are *scalar-valued* functions—the result of applying the functions is a real number (a scalar)



## Scalar- and vector-valued functions

- ▶ In Chapters 1 and 2 we considered functions of several variables  $f : D \rightarrow \mathbb{R}$  where  $D \subset \mathbb{R}^n$
- ▶ these are *scalar-valued* functions—the result of applying the functions is a real number (a scalar)
- ▶ next we consider functions  $\mathbf{f} : D \rightarrow \mathbb{R}^m$  where  $m = 2$  or  $3$

## Scalar- and vector-valued functions

- ▶ In Chapters 1 and 2 we considered functions of several variables  $f : D \rightarrow \mathbb{R}$  where  $D \subset \mathbb{R}^n$
- ▶ these are *scalar-valued* functions—the result of applying the functions is a real number (a scalar)
- ▶ next we consider functions  $\mathbf{f} : D \rightarrow \mathbb{R}^m$  where  $m = 2$  or  $3$
- ▶ they are *vector-valued* functions—the result is a 2- or 3-vector

## Scalar- and vector-valued functions

- ▶ In Chapters 1 and 2 we considered functions of several variables  $f : D \rightarrow \mathbb{R}$  where  $D \subset \mathbb{R}^n$
- ▶ these are *scalar-valued* functions—the result of applying the functions is a real number (a scalar)
- ▶ next we consider functions  $\mathbf{f} : D \rightarrow \mathbb{R}^m$  where  $m = 2$  or  $3$
- ▶ they are *vector-valued* functions—the result is a 2- or 3-vector
- ▶ examples include velocity as a function of time and direction of the Earth's magnetic field.

## Parametric equations of curves

- ▶ The simplest vector-valued functions have the form  $\mathbf{f} : I \rightarrow \mathbb{R}^2$ , where  $I \subset \mathbb{R}$ , an interval of the real line

## Parametric equations of curves

- ▶ The simplest vector-valued functions have the form  $\mathbf{f} : I \rightarrow \mathbb{R}^2$ , where  $I \subset \mathbb{R}$ , an interval of the real line
- ▶ for each  $t \in I$ ,  $\mathbf{f}(t)$  is the position vector of a point in the plane; the set of all such points forms a curve.

## Parametric equations of curves

- ▶ The simplest vector-valued functions have the form  $\mathbf{f} : I \rightarrow \mathbb{R}^2$ , where  $I \subset \mathbb{R}$ , an interval of the real line
- ▶ for each  $t \in I$ ,  $\mathbf{f}(t)$  is the position vector of a point in the plane; the set of all such points forms a curve.
- ▶ position as a function of time is one example. We will revisit parametric equations in Chapter 4.

## Scalar and vector fields

- ▶ A function  $f(\mathbf{r})$  where  $\mathbf{r} = (x, y)$  or  $(x, y, z)$  (mapping a vector to a scalar) is called a *scalar field*.

## Scalar and vector fields

- ▶ A function  $f(\mathbf{r})$  where  $\mathbf{r} = (x, y)$  or  $(x, y, z)$  (mapping a vector to a scalar) is called a *scalar field*.
- ▶ A function  $\mathbf{F}(\mathbf{r})$  where  $\mathbf{r} = (x, y)$  or  $(x, y, z)$  (mapping a vector to another vector) is called a *vector field*.




## Scalar and vector fields

- ▶ A function  $f(\mathbf{r})$  where  $\mathbf{r} = (x, y)$  or  $(x, y, z)$  (mapping a vector to a scalar) is called a *scalar field*.
- ▶ A function  $\mathbf{F}(\mathbf{r})$  where  $\mathbf{r} = (x, y)$  or  $(x, y, z)$  (mapping a vector to another vector) is called a *vector field*.

## Scalar and vector fields

- ▶ A function  $f(\mathbf{r})$  where  $\mathbf{r} = (x, y)$  or  $(x, y, z)$  (mapping a vector to a scalar) is called a *scalar field*.
- ▶ A function  $\mathbf{F}(\mathbf{r})$  where  $\mathbf{r} = (x, y)$  or  $(x, y, z)$  (mapping a vector to another vector) is called a *vector field*.

### A typical vector field



e.g. velocity at different points  
in a fluid.

## Different types of derivative

- ▶ We can define several types of derivative of scalar and vector fields, expressed in terms of

$$\nabla = \left( \frac{\partial}{\partial x}, \frac{\partial}{\partial y}, \frac{\partial}{\partial z} \right)$$

## Different types of derivative

- ▶ We can define several types of derivative of scalar and vector fields, expressed in terms of

$$\nabla = \left( \frac{\partial}{\partial x}, \frac{\partial}{\partial y}, \frac{\partial}{\partial z} \right)$$

- ▶ pronounced *del* or *nabla* (not to be confused with  $\Delta$ )

## Different types of derivative

- ▶ We can define several types of derivative of scalar and vector fields, expressed in terms of

$$\nabla = \left( \frac{\partial}{\partial x}, \frac{\partial}{\partial y}, \frac{\partial}{\partial z} \right)$$

- ▶ pronounced *del* or *nabla* (not to be confused with  $\Delta$ )
- ▶ can form “products” of this vector with other vectors and scalars, but because it is an operator, it always has to be the first term if the product is to make sense.

## Different types of derivative

- ▶ We can define several types of derivative of scalar and vector fields, expressed in terms of

$$\nabla = \left( \frac{\partial}{\partial x}, \frac{\partial}{\partial y}, \frac{\partial}{\partial z} \right)$$

- ▶ pronounced *del* or *nabla* (not to be confused with  $\Delta$ )
- ▶ can form “products” of this vector with other vectors and scalars, but because it is an operator, it always has to be the first term if the product is to make sense.

## Different types of derivative

- ▶ We can define several types of derivative of scalar and vector fields, expressed in terms of

$$\nabla = \left( \frac{\partial}{\partial x}, \frac{\partial}{\partial y}, \frac{\partial}{\partial z} \right)$$

- ▶ pronounced *del* or *nabla* (not to be confused with  $\Delta$ )
- ▶ can form “products” of this vector with other vectors and scalars, but because it is an operator, it always has to be the first term if the product is to make sense.

Name of product	Formula	Type of result	Derivative
Scalar multiplication	$\alpha \mathbf{u}$	Vector	$\nabla f$
Scalar or dot product	$\mathbf{u} \cdot \mathbf{v}$	Scalar	$\nabla \cdot \mathbf{F}$
Vector or cross product	$\mathbf{u} \times \mathbf{v}$	Vector	$\nabla \times \mathbf{F}$

## Gradient of a scalar field

The *gradient* of a scalar field  $f$  is

$$\operatorname{grad} f = \nabla f = \left( \frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}, \frac{\partial f}{\partial z} \right).$$



## Gradient of a scalar field

The *gradient* of a scalar field  $f$  is

$$\text{grad } f = \nabla f = \left( \frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}, \frac{\partial f}{\partial z} \right).$$

### Example 1

Find the gradient of the scalar field  $f(x, y, z) = x^2y + x \cosh yz$ .

## Gradient of a scalar field

The *gradient* of a scalar field  $f$  is

$$\text{grad } f = \nabla f = \left( \frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}, \frac{\partial f}{\partial z} \right).$$

### Example 1

Find the gradient of the scalar field  $f(x, y, z) = x^2y + x \cosh yz$ .

Answer

$$\text{grad } f = (2xy + \cosh yz, x^2 + xz \sinh yz, xy \sinh yz).$$

## Gradient of a scalar field

### Example 2

Let  $\mathbf{r} = (x, y, z)$  so that  $r = |\mathbf{r}| = \sqrt{x^2 + y^2 + z^2}$ . Show that

$$\nabla(r^n) = nr^{n-2}\mathbf{r},$$

for any integer  $n$  and deduce the values of  $\text{grad}(r)$ ,  $\text{grad}(r^2)$  and  $\text{grad}(1/r)$ .

## Gradient of a scalar field

### Example 2

Let  $\mathbf{r} = (x, y, z)$  so that  $r = |\mathbf{r}| = \sqrt{x^2 + y^2 + z^2}$ . Show that

$$\nabla(r^n) = nr^{n-2}\mathbf{r},$$

for any integer  $n$  and deduce the values of  $\text{grad}(r)$ ,  $\text{grad}(r^2)$  and  $\text{grad}(1/r)$ .

### Answers

$$\text{grad}(r) = \frac{\mathbf{r}}{r},$$

$$\text{grad}(r^2) = 2\mathbf{r},$$

$$\text{grad}(1/r) = -\frac{\mathbf{r}}{r^3}.$$

## Gradient of a scalar field

### Example 3

Determine  $\text{grad}(\mathbf{c} \cdot \mathbf{r})$ , when  $\mathbf{c}$  is a constant (vector).

## Gradient of a scalar field

### Example 3

Determine  $\text{grad}(\mathbf{c} \cdot \mathbf{r})$ , when  $\mathbf{c}$  is a constant (vector).

### Answer

$$\text{grad}(\mathbf{c} \cdot \mathbf{r}) = \mathbf{c}.$$

## Directional derivative

- ▶ This is the rate of change of a scalar field  $f$  in the direction of a *unit* vector  $\mathbf{u} = (u_1, u_2, u_3)$ .

## Directional derivative

- ▶ This is the rate of change of a scalar field  $f$  in the direction of a *unit* vector  $\mathbf{u} = (u_1, u_2, u_3)$ .
- ▶ defined by the limit of a difference quotient

$$\frac{\partial f}{\partial \mathbf{u}}(\mathbf{p}) = \lim_{h \rightarrow 0^+} \frac{f(\mathbf{p} + h\mathbf{u}) - f(\mathbf{p})}{h}, \quad (2)$$

(if the limit exists)



## Directional derivative

- ▶ This is the rate of change of a scalar field  $f$  in the direction of a *unit* vector  $\mathbf{u} = (u_1, u_2, u_3)$ .
- ▶ defined by the limit of a difference quotient

$$\frac{\partial f}{\partial \mathbf{u}}(\mathbf{p}) = \lim_{h \rightarrow 0^+} \frac{f(\mathbf{p} + h\mathbf{u}) - f(\mathbf{p})}{h}, \quad (2)$$

(if the limit exists)

- ▶ the key formula is:

$$\frac{\partial f}{\partial \mathbf{u}} = \mathbf{u} \cdot \nabla f = u_1 \frac{\partial f}{\partial x} + u_2 \frac{\partial f}{\partial y} + u_3 \frac{\partial f}{\partial z}.$$

## Directional derivative

- ▶ This is the rate of change of a scalar field  $f$  in the direction of a *unit* vector  $\mathbf{u} = (u_1, u_2, u_3)$ .
- ▶ defined by the limit of a difference quotient

$$\frac{\partial f}{\partial \mathbf{u}}(\mathbf{p}) = \lim_{h \rightarrow 0^+} \frac{f(\mathbf{p} + h\mathbf{u}) - f(\mathbf{p})}{h}, \quad (2)$$

(if the limit exists)

- ▶ the key formula is:

$$\boxed{\frac{\partial f}{\partial \mathbf{u}} = \mathbf{u} \cdot \nabla f = u_1 \frac{\partial f}{\partial x} + u_2 \frac{\partial f}{\partial y} + u_3 \frac{\partial f}{\partial z}}.$$

- ▶ Important: only valid for *unit* vectors  $\mathbf{u}$

## Directional derivative

- ▶ This is the rate of change of a scalar field  $f$  in the direction of a *unit* vector  $\mathbf{u} = (u_1, u_2, u_3)$ .
- ▶ defined by the limit of a difference quotient

$$\frac{\partial f}{\partial \mathbf{u}}(\mathbf{p}) = \lim_{h \rightarrow 0^+} \frac{f(\mathbf{p} + h\mathbf{u}) - f(\mathbf{p})}{h}, \quad (2)$$

(if the limit exists)

- ▶ the key formula is:

$$\boxed{\frac{\partial f}{\partial \mathbf{u}} = \mathbf{u} \cdot \nabla f = u_1 \frac{\partial f}{\partial x} + u_2 \frac{\partial f}{\partial y} + u_3 \frac{\partial f}{\partial z}}.$$

- ▶ Important: only valid for *unit* vectors  $\mathbf{u}$
- ▶ Partial derivatives are directional derivatives, e.g.

$$\frac{\partial f}{\partial \mathbf{i}} = \frac{\partial f}{\partial x}.$$

## Directional derivative

### Example 4

Find the directional derivative of  $f = x^2yz^3$  at the point  $P(3, -2, -1)$  in the direction of the vector  $(1, 2, 2)$ .

## Directional derivative

### Example 4

Find the directional derivative of  $f = x^2yz^3$  at the point  $P(3, -2, -1)$  in the direction of the vector  $(1, 2, 2)$ .

Answer

$$\frac{\partial f}{\partial \mathbf{u}}(3, -2, -1) = -38.$$

## Directional derivative

- ▶ If we fix a point  $\mathbf{p}$  and are given a function  $f$ , then by considering all possible directional derivatives of  $f$  at the point  $\mathbf{p}$  we can ask:

## Directional derivative

- ▶ If we fix a point  $\mathbf{p}$  and are given a function  $f$ , then by considering all possible directional derivatives of  $f$  at the point  $\mathbf{p}$  we can ask:
  - ▶ in which direction does  $f$  change fastest?

## Directional derivative

- ▶ If we fix a point  $\mathbf{p}$  and are given a function  $f$ , then by considering all possible directional derivatives of  $f$  at the point  $\mathbf{p}$  we can ask:
  - ▶ in which direction does  $f$  change fastest?
  - ▶ what is the maximal rate of change?



## Directional derivative

- ▶ If we fix a point  $\mathbf{p}$  and are given a function  $f$ , then by considering all possible directional derivatives of  $f$  at the point  $\mathbf{p}$  we can ask:
  - ▶ in which direction does  $f$  change fastest?
  - ▶ what is the maximal rate of change?
- ▶ The following theorem answers these questions.

## Directional derivative

- ▶ If we fix a point  $\mathbf{p}$  and are given a function  $f$ , then by considering all possible directional derivatives of  $f$  at the point  $\mathbf{p}$  we can ask:
  - ▶ in which direction does  $f$  change fastest?
  - ▶ what is the maximal rate of change?
- ▶ The following theorem answers these questions.
- ▶ **Theorem:** Suppose  $f$  is a differentiable function for which  $\nabla f(\mathbf{p}) \neq 0$  then the maximal value of  $\frac{\partial f}{\partial \mathbf{u}}(\mathbf{p})$  is  $|\nabla f(\mathbf{p})|$  and occurs when  $\mathbf{u}$  is in the same direction as  $\nabla f$ .

## Directional derivative

- ▶ If we fix a point  $\mathbf{p}$  and are given a function  $f$ , then by considering all possible directional derivatives of  $f$  at the point  $\mathbf{p}$  we can ask:
  - ▶ in which direction does  $f$  change fastest?
  - ▶ what is the maximal rate of change?
- ▶ The following theorem answers these questions.
- ▶ **Theorem:** Suppose  $f$  is a differentiable function for which  $\nabla f(\mathbf{p}) \neq 0$  then the maximal value of  $\frac{\partial f}{\partial \mathbf{u}}(\mathbf{p})$  is  $|\nabla f(\mathbf{p})|$  and occurs when  $\mathbf{u}$  is in the same direction as  $\nabla f$ .

## Directional derivative

- ▶ If we fix a point  $\mathbf{p}$  and are given a function  $f$ , then by considering all possible directional derivatives of  $f$  at the point  $\mathbf{p}$  we can ask:
  - ▶ in which direction does  $f$  change fastest?
  - ▶ what is the maximal rate of change?
- ▶ The following theorem answers these questions.
- ▶ **Theorem:** Suppose  $f$  is a differentiable function for which  $\nabla f(\mathbf{p}) \neq 0$  then the maximal value of  $\frac{\partial f}{\partial \mathbf{u}}(\mathbf{p})$  is  $|\nabla f(\mathbf{p})|$  and occurs when  $\mathbf{u}$  is in the same direction as  $\nabla f$ .

## Directional derivative

### Example 5

Consider  $f = \ln(xy + z^3)$  at the point  $P(1, 1, 1)$ . In what direction does  $f$  have the maximal rate of change? What is this maximal rate of change?

## Directional derivative

### Example 5

Consider  $f = \ln(xy + z^3)$  at the point  $P(1, 1, 1)$ . In what direction does  $f$  have the maximal rate of change? What is this maximal rate of change?

### Answer

Direction is  $(1/2, 1/2, 3/2)$ . Maximal rate of change is

$$|\nabla f(1, 1, 1)| = \frac{\sqrt{11}}{2}.$$

## Divergence of a vector field

- ▶ The *divergence* of a vector field  $\mathbf{F} = (F_1, F_2, F_3)$  is the *scalar* obtained as the “scalar product” of  $\nabla$  and  $\mathbf{F}$ ,

$$\operatorname{div} \mathbf{F} = \nabla \cdot \mathbf{F} = \frac{\partial F_1}{\partial x} + \frac{\partial F_2}{\partial y} + \frac{\partial F_3}{\partial z},$$

## Divergence of a vector field

- ▶ The *divergence* of a vector field  $\mathbf{F} = (F_1, F_2, F_3)$  is the *scalar* obtained as the “scalar product” of  $\nabla$  and  $\mathbf{F}$ ,

$$\operatorname{div} \mathbf{F} = \nabla \cdot \mathbf{F} = \frac{\partial F_1}{\partial x} + \frac{\partial F_2}{\partial y} + \frac{\partial F_3}{\partial z},$$

- ▶ so called, because it measures the tendency of a vector field to diverge (positive divergence) or converge (negative divergence)



## Divergence of a vector field

- ▶ The *divergence* of a vector field  $\mathbf{F} = (F_1, F_2, F_3)$  is the *scalar* obtained as the “scalar product” of  $\nabla$  and  $\mathbf{F}$ ,

$$\operatorname{div} \mathbf{F} = \nabla \cdot \mathbf{F} = \frac{\partial F_1}{\partial x} + \frac{\partial F_2}{\partial y} + \frac{\partial F_3}{\partial z},$$

- ▶ so called, because it measures the tendency of a vector field to diverge (positive divergence) or converge (negative divergence)
- ▶ a vector field is said to be *incompressible* (or *solenoidal*) if its divergence is zero.

## Divergence of a vector field

- ▶ Consider  $\mathbf{F} = (x, y, 0)$ ,  $\mathbf{G} = (x, -y, 0)$  and  $\mathbf{H} = (-x, -y, 0)$

## Divergence of a vector field


- ▶ Consider  $\mathbf{F} = (x, y, 0)$ ,  $\mathbf{G} = (x, -y, 0)$  and  $\mathbf{H} = (-x, -y, 0)$
- ▶ we have

$$\operatorname{div} \mathbf{F} = \frac{\partial x}{\partial x} + \frac{\partial y}{\partial y} = 2 > 0, \quad \operatorname{div} \mathbf{G} = 0 \quad \text{and} \quad \operatorname{div} \mathbf{H} = -2 < 0.$$


# Divergence of a vector field

- Consider  $\mathbf{F} = (x, y, 0)$ ,  $\mathbf{G} = (x, -y, 0)$  and  $\mathbf{H} = (-x, -y, 0)$
- we have


$$\operatorname{div} \mathbf{F} = \frac{\partial x}{\partial x} + \frac{\partial y}{\partial y} = 2 > 0, \quad \operatorname{div} \mathbf{G} = 0 \quad \text{and} \quad \operatorname{div} \mathbf{H} = -2 < 0.$$



$\mathbf{F}$ , positive divergence



$\mathbf{G}$ , incompressible



$\mathbf{H}$ , negative divergence

## Divergence of a vector field

### Example 6

Show that the divergence of  $\mathbf{F} = (x - y^2, z, z^3)$  is positive at all points in  $\mathbb{R}^3$ .

# Laplacian

- ▶ A particular example of divergence is the *Laplacian* of a scalar field  $f$

# Laplacian

- ▶ A particular example of divergence is the *Laplacian* of a scalar field  $f$
- ▶  $\text{grad } f = \nabla f$  is a vector field and so we can take its divergence

# Laplacian

- ▶ A particular example of divergence is the *Laplacian* of a scalar field  $f$
- ▶  $\text{grad } f = \nabla f$  is a vector field and so we can take its divergence
- ▶ this is the Laplacian of  $f$ , written  $\nabla^2 f$



# Laplacian

- ▶ A particular example of divergence is the *Laplacian* of a scalar field  $f$
- ▶  $\text{grad } f = \nabla f$  is a vector field and so we can take its divergence
- ▶ this is the Laplacian of  $f$ , written  $\nabla^2 f$
- ▶ we have

$$\nabla^2 f = \nabla \cdot (\nabla f) = \frac{\partial^2 f}{\partial x^2} + \frac{\partial^2 f}{\partial y^2} + \frac{\partial^2 f}{\partial z^2} .$$

# Laplacian

- ▶ A particular example of divergence is the *Laplacian* of a scalar field  $f$
- ▶  $\text{grad } f = \nabla f$  is a vector field and so we can take its divergence
- ▶ this is the Laplacian of  $f$ , written  $\nabla^2 f$
- ▶ we have

$$\nabla^2 f = \nabla \cdot (\nabla f) = \frac{\partial^2 f}{\partial x^2} + \frac{\partial^2 f}{\partial y^2} + \frac{\partial^2 f}{\partial z^2} .$$

- ▶ Can be extended in a natural way to the Laplacian of a vector field  $\mathbf{F} = (F_1, F_2, F_3)$ ,

$$\nabla^2 \mathbf{F} = (\nabla^2 F_1, \nabla^2 F_2, \nabla^2 F_3) .$$

# Laplacian

## Example 7

Find the values of  $n$  for which  $\nabla^2(r^n) = 0$ .

# Laplacian

## Example 7

Find the values of  $n$  for which  $\nabla^2(r^n) = 0$ .

## Answer

$\nabla^2(r^n) = 0$  if and only if  $n = 0$  or  $n = -1$ .

## Curl of a vector field

- ▶ The *curl* of a vector field  $\mathbf{F} = (F_1, F_2, F_3)$  is the *vector* obtained as the “vector product” of  $\nabla$  and  $\mathbf{F}$

## Curl of a vector field

- ▶ The *curl* of a vector field  $\mathbf{F} = (F_1, F_2, F_3)$  is the *vector* obtained as the “vector product” of  $\nabla$  and  $\mathbf{F}$
- ▶ the formula is

$$\operatorname{curl} \mathbf{F} = \left( \frac{\partial F_3}{\partial y} - \frac{\partial F_2}{\partial z} \right) \mathbf{i} + \left( \frac{\partial F_1}{\partial z} - \frac{\partial F_3}{\partial x} \right) \mathbf{j} + \left( \frac{\partial F_2}{\partial x} - \frac{\partial F_1}{\partial y} \right) \mathbf{k} .$$

## Curl of a vector field

- ▶ The *curl* of a vector field  $\mathbf{F} = (F_1, F_2, F_3)$  is the *vector* obtained as the “vector product” of  $\nabla$  and  $\mathbf{F}$
- ▶ the formula is

$$\text{curl } \mathbf{F} = \left( \frac{\partial F_3}{\partial y} - \frac{\partial F_2}{\partial z} \right) \mathbf{i} + \left( \frac{\partial F_1}{\partial z} - \frac{\partial F_3}{\partial x} \right) \mathbf{j} + \left( \frac{\partial F_2}{\partial x} - \frac{\partial F_1}{\partial y} \right) \mathbf{k}.$$

- ▶ can be calculated using a  $3 \times 3$  determinant,

$$\text{curl } \mathbf{F} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ F_1 & F_2 & F_3 \end{vmatrix}.$$

## Curl of a vector field

- ▶ Curl of a vector field measures its tendency to rotate



## Curl of a vector field

- ▶ Curl of a vector field measures its tendency to rotate
- ▶ a vector field is said to be *irrotational* if its curl is the zero vector

## Curl of a vector field

- ▶ Curl of a vector field measures its tendency to rotate
- ▶ a vector field is said to be *irrotational* if its curl is the zero vector
- ▶ consider  $\mathbf{F} = (-y, x, 0)$ ,  $\mathbf{G} = (y, x, 0)$  and  $\mathbf{H} = (y, -x, 0)$

## Curl of a vector field


- ▶ Curl of a vector field measures its tendency to rotate
- ▶ a vector field is said to be *irrotational* if its curl is the zero vector
- ▶ consider  $\mathbf{F} = (-y, x, 0)$ ,  $\mathbf{G} = (y, x, 0)$  and  $\mathbf{H} = (y, -x, 0)$
- ▶ we have

$$\text{curl } \mathbf{F} = 2\mathbf{k}, \quad \text{curl } \mathbf{G} = \mathbf{0} \quad \text{and} \quad \text{curl } \mathbf{H} = -2\mathbf{k}$$


# Curl of a vector field

- ▶ Curl of a vector field measures its tendency to rotate
- ▶ a vector field is said to be *irrotational* if its curl is the zero vector
- ▶ consider  $\mathbf{F} = (-y, x, 0)$ ,  $\mathbf{G} = (y, x, 0)$  and  $\mathbf{H} = (y, -x, 0)$
- ▶ we have


$$\text{curl } \mathbf{F} = 2\mathbf{k}, \quad \text{curl } \mathbf{G} = \mathbf{0} \quad \text{and} \quad \text{curl } \mathbf{H} = -2\mathbf{k}$$



$\mathbf{F}$ , anticlockwise rotation



$\mathbf{G}$ , irrotational



$\mathbf{H}$ , clockwise rotation

## Curl of a vector field

### Example 8

Determine  $\text{curl } \mathbf{F}$  when  $\mathbf{F} = (x^2y, xy^2 + z, xy)$ .

## Curl of a vector field

### Example 8

Determine  $\text{curl } \mathbf{F}$  when  $\mathbf{F} = (x^2y, xy^2 + z, xy)$ .

### Answer

$$\text{curl } \mathbf{F} = (x - 1, -y, y^2 - x^2).$$

## Curl of a vector field

### Example 9

If  $\mathbf{c}$  is a constant vector, find  $\text{curl}(\mathbf{c} \times \mathbf{r})$ .

## Curl of a vector field

### Example 9

If  $\mathbf{c}$  is a constant vector, find  $\text{curl}(\mathbf{c} \times \mathbf{r})$ .

### Answer

$$\text{curl}(\mathbf{c} \times \mathbf{r}) = 2\mathbf{c}.$$



## Nabla identities

Analogues involving  $\text{div}$ ,  $\text{grad}$  and  $\text{curl}$  of the elementary rules of differentiation such as linearity  $(f + g)'(x) = f'(x) + g'(x)$  the product rule  $(fg)'(x) = f(x)g'(x) + f'(x)g(x)$ .

## Nabla identities

Analogues involving  $\text{div}$ ,  $\text{grad}$  and  $\text{curl}$  of the elementary rules of differentiation such as linearity  $(f + g)'(x) = f'(x) + g'(x)$  the product rule  $(fg)'(x) = f(x)g'(x) + f'(x)g(x)$ .

$\text{grad}(f + g) = \text{grad } f + \text{grad } g$	$\text{grad}(fg) = f(\text{grad } g) + (\text{grad } f)g,$
$\text{div}(\mathbf{F} + \mathbf{G}) = \text{div } \mathbf{F} + \text{div } \mathbf{G}$	$\text{div}(f\mathbf{F}) = f \text{div } \mathbf{F} + \text{grad } f \cdot \mathbf{F},$
$\text{curl}(\mathbf{F} + \mathbf{G}) = \text{curl } \mathbf{F} + \text{curl } \mathbf{G}$	$\text{curl}(f\mathbf{F}) = f \text{curl } \mathbf{F} + \text{grad } f \times \mathbf{F},$
$\text{curl grad } f = \mathbf{0},$	$\text{div curl } \mathbf{F} = 0.$

## Nabla identities

- Note the special cases

$$\text{grad}(cf) = c \text{ grad } f, \quad \text{div}(c\mathbf{F}) = c \text{ div } \mathbf{F}, \quad \text{curl}(c\mathbf{F}) = c \text{ curl } \mathbf{F},$$

when  $c$  is a (scalar) constant.

## Nabla identities

- Note the special cases

$$\text{grad}(cf) = c \text{ grad } f, \quad \text{div}(c\mathbf{F}) = c \text{ div } \mathbf{F}, \quad \text{curl}(c\mathbf{F}) = c \text{ curl } \mathbf{F},$$

when  $c$  is a (scalar) constant.

- all of the identities are easier to remember if written using  $\nabla$

## Nabla identities

- Note the special cases

$$\text{grad}(cf) = c \text{ grad } f, \quad \text{div}(c\mathbf{F}) = c \text{ div } \mathbf{F}, \quad \text{curl}(c\mathbf{F}) = c \text{ curl } \mathbf{F},$$

when  $c$  is a (scalar) constant.

- all of the identities are easier to remember if written using  $\nabla$

- e.g.

$$\begin{aligned}\text{curl}(f\mathbf{F}) &= \nabla \times (f\mathbf{F}) \\ &= f(\nabla \times \mathbf{F}) + (\nabla f) \times \mathbf{F} \\ &= f \text{ curl } \mathbf{F} + \text{grad } f \times \mathbf{F}.\end{aligned}$$

# Nabla identities

## Example 10

Prove the identities

$$(i) \operatorname{curl} \operatorname{grad} f = 0, \quad (ii) \operatorname{curl}(f \mathbf{F}) = f \operatorname{curl} \mathbf{F} + \operatorname{grad} f \times \mathbf{F}$$

$$(iii) \operatorname{div}(f \mathbf{F}) = f \operatorname{div} \mathbf{F} + (\operatorname{grad} f) \cdot \mathbf{F}.$$

## Nabla identities

### Example 11

Let  $\mathbf{c}$  be a constant vector and  $\mathbf{r} = (x, y, z)$  so that  $r = |\mathbf{r}| = \sqrt{x^2 + y^2 + z^2}$ . Determine

$$(i) \operatorname{div}(r^n(\mathbf{c} \times \mathbf{r})), \quad (ii) \operatorname{curl}(r^n(\mathbf{c} \times \mathbf{r})).$$

## Nabla identities

### Example 11

Let  $\mathbf{c}$  be a constant vector and  $\mathbf{r} = (x, y, z)$  so that  $r = |\mathbf{r}| = \sqrt{x^2 + y^2 + z^2}$ . Determine

$$(i) \operatorname{div}(r^n(\mathbf{c} \times \mathbf{r})), \quad (ii) \operatorname{curl}(r^n(\mathbf{c} \times \mathbf{r})).$$

### Answers

$$(i) \operatorname{div}(r^n(\mathbf{c} \times \mathbf{r})) = 0 \quad ,$$

$$(ii) \operatorname{curl}(r^n(\mathbf{c} \times \mathbf{r})) = (n+2)r^n\mathbf{c} - n(\mathbf{r} \cdot \mathbf{c})r^{n-2}\mathbf{r}.$$



## Chapter 4: Line and surface integrals

- ▶ Line integrals in two dimensions

## Chapter 4: Line and surface integrals

- ▶ Line integrals in two dimensions
- ▶ Green's Theorem

## Chapter 4: Line and surface integrals

- ▶ Line integrals in two dimensions
- ▶ Green's Theorem
- ▶ Line integrals of closed paths - path independence and conservative vector fields.

## Chapter 4: Line and surface integrals

- ▶ Line integrals in two dimensions
- ▶ Green's Theorem
- ▶ Line integrals of closed paths - path independence and conservative vector fields.
- ▶ Surface integrals

## Chapter 4: Line and surface integrals

- ▶ Line integrals in two dimensions
- ▶ Green's Theorem
- ▶ Line integrals of closed paths - path independence and conservative vector fields.
- ▶ Surface integrals
- ▶ Surface integral of a vector field and Gauss' Divergence Theorem

## Line integrals in two dimensions

- ▶ In first year we integrate over an interval  $[a, b]$ , we will now integrate over a curve in  $\mathbb{R}^2$ .

## Line integrals in two dimensions

- ▶ In first year we integrate over an interval  $[a, b]$ , we will now integrate over a curve in  $\mathbb{R}^2$ .
- ▶ Such curves can be described parametrically for each  $t \in I$ , ( $I \subset \mathbb{R}$ , an interval of the real line)  $\mathbf{f}(t)$  is the position vector of a point in the plane; the set of all such points forms a curve.

## Line integrals in two dimensions

- ▶ In first year we integrate over an interval  $[a, b]$ , we will now integrate over a curve in  $\mathbb{R}^2$ .
- ▶ Such curves can be described parametrically for each  $t \in I$ , ( $I \subset \mathbb{R}$ , an interval of the real line)  $\mathbf{f}(t)$  is the position vector of a point in the plane; the set of all such points forms a curve.
- ▶ We first recall some parametric equations from level 1 and then introduce the concept of a line integral.



## Parametric equation of a line

- ▶ Recall: section formula—if  $P$  lies on the line through  $A$  and  $B$  then

$$\mathbf{p} = \frac{\alpha \mathbf{a} + \beta \mathbf{b}}{\alpha + \beta}, \quad \text{some } \alpha, \beta$$

## Parametric equation of a line

- ▶ Recall: section formula—if  $P$  lies on the line through  $A$  and  $B$  then

$$\mathbf{p} = \frac{\alpha \mathbf{a} + \beta \mathbf{b}}{\alpha + \beta}, \quad \text{some } \alpha, \beta$$

where  $\mathbf{a}$ ,  $\mathbf{b}$  and  $\mathbf{p}$  are the position vectors of  $A$ ,  $B$  and  $P$

## Parametric equation of a line

- ▶ Recall: section formula—if  $P$  lies on the line through  $A$  and  $B$  then

$$\mathbf{p} = \frac{\alpha \mathbf{a} + \beta \mathbf{b}}{\alpha + \beta}, \quad \text{some } \alpha, \beta$$

- ▶ can write as

$$\mathbf{p}(t) = (1 - t)\mathbf{a} + t\mathbf{b}.$$

where  $t = \beta/(\alpha + \beta)$

## Parametric equation of a line

- ▶ Recall: section formula—if  $P$  lies on the line through  $A$  and  $B$  then

$$\mathbf{p} = \frac{\alpha \mathbf{a} + \beta \mathbf{b}}{\alpha + \beta}, \quad \text{some } \alpha, \beta$$

- ▶ can write as

$$\mathbf{p}(t) = (1 - t)\mathbf{a} + t\mathbf{b}.$$

where  $t = \beta/(\alpha + \beta)$

- ▶ Note:  $\mathbf{p}(0) = \mathbf{a}$  and  $\mathbf{p}(1) = \mathbf{b}$  and

$$\{\mathbf{p}(t) : t \in [0, 1]\}$$

is the line segment from  $A$  to  $B$ .

## Parametric equation of a line

- ▶ Recall: section formula—if  $P$  lies on the line through  $A$  and  $B$  then

$$\mathbf{p} = \frac{\alpha \mathbf{a} + \beta \mathbf{b}}{\alpha + \beta}, \quad \text{some } \alpha, \beta$$

- ▶ can write as

$$\mathbf{p}(t) = (1 - t)\mathbf{a} + t\mathbf{b}.$$

where  $t = \beta/(\alpha + \beta)$

- ▶ Note:  $\mathbf{p}(0) = \mathbf{a}$  and  $\mathbf{p}(1) = \mathbf{b}$  and

$$\{\mathbf{p}(t) : t \in [0, 1]\}$$

is the line segment from  $A$  to  $B$ .

- ▶ Also,

$$\mathbf{p}(t) = \mathbf{a} + t(\mathbf{b} - \mathbf{a}) = \mathbf{a} + s\mathbf{d},$$

where  $\mathbf{d}$  is a direction vector for the line.

## Parametric equations of curves

- ▶ In general, a curve, in 2D or 3D space, can be represented as

$$\mathbf{r} = \mathbf{f}(t), \quad t \in I.$$

## Parametric equations of curves

- ▶ In general, a curve, in 2D or 3D space, can be represented as

$$\mathbf{r} = \mathbf{f}(t), \quad t \in I.$$

- ▶ called a *parametric* description of the curve and  $t$  is called a *parameter*

## Parametric equations of curves

- ▶ In general, a curve, in 2D or 3D space, can be represented as

$$\mathbf{r} = \mathbf{f}(t), \quad t \in I.$$

- ▶ called a *parametric* description of the curve and  $t$  is called a *parameter*
- ▶ may also be written in component form; if  $\mathbf{r} = (x, y, z)$  and  $\mathbf{f} = (f_1, f_2, f_3)$  then

$$x = f_1(t), \quad y = f_2(t), \quad z = f_3(t), \quad t \in I.$$



## Standard parametric curves—circles

- ▶ Circle  $(x - a)^2 + (y - b)^2 = r^2$  has centre  $(a, b)$  and radius  $r$

## Standard parametric curves—circles

- ▶ Circle  $(x - a)^2 + (y - b)^2 = r^2$  has centre  $(a, b)$  and radius  $r$
- ▶ can be parametrised using polar coordinates as

$$x = a + r \cos \theta, \quad y = b + r \sin \theta, \quad \theta \in [0, 2\pi)$$

## Standard parametric curves—circles

- ▶ Circle  $(x - a)^2 + (y - b)^2 = r^2$  has centre  $(a, b)$  and radius  $r$
- ▶ can be parametrised using polar coordinates as

$$x = a + r \cos \theta, \quad y = b + r \sin \theta, \quad \theta \in [0, 2\pi)$$

- ▶  $\theta$  is the angle measured anti-clockwise from the positive  $x$ -axis.

## Standard parametric curves—circles

- ▶ Circle  $(x - a)^2 + (y - b)^2 = r^2$  has centre  $(a, b)$  and radius  $r$
- ▶ can be parametrised using polar coordinates as

$$x = a + r \cos \theta, \quad y = b + r \sin \theta, \quad \theta \in [0, 2\pi)$$

- ▶  $\theta$  is the angle measured anti-clockwise from the positive  $x$ -axis.
- ▶ As a curve on the  $xy$ -plane in 3D space it is

$$x = a + r \cos \theta, \quad y = b + r \sin \theta, \quad z = 0, \quad \theta \in [0, 2\pi).$$

## Standard parametric curves—circles

- ▶ Circle  $(x - a)^2 + (y - b)^2 = r^2$  has centre  $(a, b)$  and radius  $r$
- ▶ can be parametrised using polar coordinates as

$$x = a + r \cos \theta, \quad y = b + r \sin \theta, \quad \theta \in [0, 2\pi)$$

- ▶  $\theta$  is the angle measured anti-clockwise from the positive  $x$ -axis.
- ▶ As a curve on the  $xy$ -plane in 3D space it is

$$x = a + r \cos \theta, \quad y = b + r \sin \theta, \quad z = 0, \quad \theta \in [0, 2\pi).$$

- ▶ For example,  $x^2 - 4x + y^2 + 1 = 0$  has parametric form

$$x = 2 + \sqrt{3} \cos \theta, \quad y = \sqrt{3} \sin \theta, \quad \theta \in [0, 2\pi)$$

## Standard parametric curves—circles

- ▶ Circle  $(x - a)^2 + (y - b)^2 = r^2$  has centre  $(a, b)$  and radius  $r$
- ▶ can be parametrised using polar coordinates as

$$x = a + r \cos \theta, \quad y = b + r \sin \theta, \quad \theta \in [0, 2\pi)$$

- ▶  $\theta$  is the angle measured anti-clockwise from the positive  $x$ -axis.
- ▶ As a curve on the  $xy$ -plane in 3D space it is

$$x = a + r \cos \theta, \quad y = b + r \sin \theta, \quad z = 0, \quad \theta \in [0, 2\pi).$$

- ▶ For example,  $x^2 - 4x + y^2 + 1 = 0$  has parametric form

$$x = 2 + \sqrt{3} \cos \theta, \quad y = \sqrt{3} \sin \theta, \quad \theta \in [0, 2\pi)$$

- ▶ restricting  $\theta$  to a smaller interval gives part of the circle; e.g.  $[0, \pi]$  give the top semicircle

## Standard parametric curves—ellipse and parabola

- ▶ Similarly, the ellipse

$$\frac{(x - a)^2}{A^2} + \frac{(y - b)^2}{B^2} = 1,$$

has parametric form

$$x = a + A \cos \theta, \quad y = b + B \sin \theta, \quad \theta \in [0, 2\pi).$$

## Standard parametric curves—ellipse and parabola

- ▶ Similarly, the ellipse

$$\frac{(x-a)^2}{A^2} + \frac{(y-b)^2}{B^2} = 1,$$

has parametric form

$$x = a + A \cos \theta, \quad y = b + B \sin \theta, \quad \theta \in [0, 2\pi).$$

- ▶ e.g. the right semi-ellipse is

$$x = a + A \cos \theta, \quad y = b + B \sin \theta, \quad \theta \in [-\pi/2, \pi/2].$$



## Standard parametric curves—ellipse and parabola

- ▶ Similarly, the ellipse

$$\frac{(x-a)^2}{A^2} + \frac{(y-b)^2}{B^2} = 1,$$

has parametric form

$$x = a + A \cos \theta, \quad y = b + B \sin \theta, \quad \theta \in [0, 2\pi).$$

- ▶ e.g. the right semi-ellipse is

$$x = a + A \cos \theta, \quad y = b + B \sin \theta, \quad \theta \in [-\pi/2, \pi/2].$$

- ▶ The parabola  $y^2 = 4ax$  (standard form) has parametric form

$$x = at^2, \quad y = 2at, \quad t \in (-\infty, \infty).$$

## Standard parametric curves—ellipse and parabola

- ▶ Similarly, the ellipse

$$\frac{(x-a)^2}{A^2} + \frac{(y-b)^2}{B^2} = 1,$$

has parametric form

$$x = a + A \cos \theta, \quad y = b + B \sin \theta, \quad \theta \in [0, 2\pi).$$

- ▶ e.g. the right semi-ellipse is

$$x = a + A \cos \theta, \quad y = b + B \sin \theta, \quad \theta \in [-\pi/2, \pi/2].$$

- ▶ The parabola  $y^2 = 4ax$  (standard form) has parametric form


$$x = at^2, \quad y = 2at, \quad t \in (-\infty, \infty).$$

- ▶ For example,  $y = x^2 - x$  may be written in parametric form as

$$x = t + \frac{1}{2}, \quad y = t^2 - \frac{1}{4}, \quad t \in (-\infty, \infty).$$


## Differentiation of vector-valued functions

Consider a curve defined by  $\mathbf{r} = \mathbf{r}(t)$ , the path taken by a particle and  $t$  is time.



## Differentiation of vector-valued functions

Consider a curve defined by  $\mathbf{r} = \mathbf{r}(t)$ , the path taken by a particle and  $t$  is time.



Average velocity in  $[t, t+h]$   
is

$$\frac{\text{displacement}}{\text{length}} = \frac{\mathbf{r}(t+h) - \mathbf{r}(t)}{h}.$$

## Differentiation of vector-valued functions

- In component form,

$$\left( \frac{x(t+h) - x(t)}{h}, \frac{y(t+h) - y(t)}{h}, \frac{z(t+h) - z(t)}{h} \right)$$

## Differentiation of vector-valued functions

- In component form,

$$\left( \frac{x(t+h) - x(t)}{h}, \frac{y(t+h) - y(t)}{h}, \frac{z(t+h) - z(t)}{h} \right)$$

- if scalar functions  $x$ ,  $y$  and  $z$  are differentiable then this has a limit as  $h \rightarrow 0$

$$\left( \frac{dx}{dt}, \frac{dy}{dt}, \frac{dz}{dt} \right) = (\dot{x}, \dot{y}, \dot{z})$$

## Differentiation of vector-valued functions

- In component form,

$$\left( \frac{x(t+h) - x(t)}{h}, \frac{y(t+h) - y(t)}{h}, \frac{z(t+h) - z(t)}{h} \right)$$

- if scalar functions  $x$ ,  $y$  and  $z$  are differentiable then this has a limit as  $h \rightarrow 0$

$$\left( \frac{dx}{dt}, \frac{dy}{dt}, \frac{dz}{dt} \right) = (\dot{x}, \dot{y}, \dot{z})$$

- this is the *instantaneous velocity* of the particle,

$$\mathbf{v}(t) = \lim_{h \rightarrow 0} \frac{\mathbf{r}(t+h) - \mathbf{r}(t)}{h} = \frac{d}{dt} \mathbf{r}(t) = \dot{\mathbf{r}}(t).$$

## Line integrals in two dimensions

- ▶ One of simplest examples of line integrals arises from considering the work done by a force acting on particle.



## Line integrals in two dimensions

- ▶ One of simplest examples of line integrals arises from considering the work done by a force acting on particle.
- ▶ We can describe the work done by force. On an interval this gives

$$\text{Work} = \text{Force} \times \text{distance} = \sum_x f(x) \delta x = \int_a^b f(x) dx$$


## Line integrals in two dimensions

- ▶ One of simplest examples of line integrals arises from considering the work done by a force acting on particle.
- ▶ We can describe the work done by force. On an interval this gives

$$\text{Work} = \text{Force} \times \text{distance} = \sum_x f(x) \delta x = \int_a^b f(x) dx$$

- ▶ Generalising this to the work done to move a particle along a curve  $C$  gives a line integral.

$$W = |\mathbf{D}| |\mathbf{F}| \cos \theta = \mathbf{F} \cdot \mathbf{D},$$



## Line integrals in two dimensions

- ▶ Let  $\mathbf{r}(t) = (x(t), y(t))$  describe the parameterised curve  $C$ ,  $d\mathbf{r} = (dx, dy)$  is small step along that curve. Then if  $\mathbf{F} = (P(x, y), Q(x, y))$  is the force used to move the particle along  $C$  then

$$\text{Work done} = \int_C \mathbf{F} \cdot d\mathbf{r} = \int_C P(x, y)dx + Q(x, y)dy ,$$

## Line integrals in two dimensions

- ▶ Let  $\mathbf{r}(t) = (x(t), y(t))$  describe the parameterised curve  $C$ ,  $d\mathbf{r} = (dx, dy)$  is small step along that curve. Then if  $\mathbf{F} = (P(x, y), Q(x, y))$  is the force used to move the particle along  $C$  then

$$\text{Work done} = \int_C \mathbf{F} \cdot d\mathbf{r} = \int_C P(x, y)dx + Q(x, y)dy ,$$

- ▶ Parameterise the curve  $C$  by

$$x = x(t), \quad y = y(t) \quad a \leq t \leq b$$


then  $d\mathbf{r} = \frac{d\mathbf{r}}{dt}dt$  so this gives

$$\int_a^b \mathbf{F}(\mathbf{r}(t)) \frac{d\mathbf{r}}{dt} dt = \int_a^b P(x(t), y(t)) \frac{dx}{dt} dt + \int_a^b Q(x(t), y(t)) \frac{dy}{dt} dt .$$

## Line integrals in two dimensions

### Example 1


Find the work done by the force  $\mathbf{F}(x, y) = x^2\mathbf{i} - xy\mathbf{j}$  in moving a particle along the curve which runs from  $(1, 0)$  to  $(0, 1)$  along the unit circle and then from  $(0, 1)$  to  $(0, 0)$  along the  $y$ -axis.



## Line integrals in two dimensions

### Example 1

Find the work done by the force  $\mathbf{F}(x, y) = x^2\mathbf{i} - xy\mathbf{j}$  in moving a particle along the curve which runs from  $(1, 0)$  to  $(0, 1)$  along the unit circle and then from  $(0, 1)$  to  $(0, 0)$  along the  $y$ -axis.



## Line integrals in two dimensions

### Example 2


Evaluate the line integral  $\int_C (y^2)dx + (x)dy$ , where  $C$  is the arc of the parabola  $x = 4 - y^2$  from  $(-5, -3)$  to  $(0, 2)$ .

## Line integrals in two dimensions

### Example 2

Evaluate the line integral  $\int_C (y^2)dx + (x)dy$ , where  $C$  is the arc of the parabola  $x = 4 - y^2$  from  $(-5, -3)$  to  $(0, 2)$ .

### Answers






## Line integrals in two dimensions

### Example 2

Evaluate the line integral  $\int_C (y^2)dx + (x)dy$ , where  $C$  is the arc of the parabola  $x = 4 - y^2$  from  $(-5, -3)$  to  $(0, 2)$ .

### Answers



245/6.

## Line integrals in two dimensions

- ▶ If  $C$  is something simple like a straight line then it's often easier **not** to parameterise  $C$ .

## Line integrals in two dimensions

- ▶ If  $C$  is something simple like a straight line then it's often easier **not** to parameterise  $C$ .
- ▶ Instead use

$$\int_C \mathbf{F} \cdot d\mathbf{r} = \int_C P(x, y)dx + Q(x, y)dy ,$$

directly.

## Line integrals in two dimensions

### Example 3

Evaluate the line integral,  $\int_C (x^2 + y^2)dx + (4x + y^2)dy$ , where  $C$  is the straight line segment from  $(6, 3)$  to  $(6, 0)$ .

## Line integrals in two dimensions

### Example 3

Evaluate the line integral,  $\int_C (x^2 + y^2)dx + (4x + y^2)dy$ , where  $C$  is the straight line segment from  $(6, 3)$  to  $(6, 0)$ .


### Answers

-81.


# Green's Theorem

Let  $C$  be a positively oriented simple closed curve in the plane and let  $D$  be the region bounded by  $C$ . If  $P(x, y)$  and  $Q(x, y)$  have continuous partial derivatives on an open region that contains  $D$ , then

$$\int_C P(x, y)dx + Q(x, y)dy = \iint_D \left( \frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} \right) dx dy .$$



(a) Positive orientation



(b) Negative orientation

## Green's Theorem

### Example 4

Use Green's Theorem to evaluate

$$\int_C (3y - e^{\sin x})dx + (7x + \sqrt{y^4 + 1})dy, \text{ where } C \text{ is the circle } x^2 + y^2 = 9.$$

# Green's Theorem

## Example 4

Use Green's Theorem to evaluate

$\int_C (3y - e^{\sin x})dx + (7x + \sqrt{y^4 + 1})dy$ , where  $C$  is the circle  $x^2 + y^2 = 9$ .

## Answers

$36\pi$



## Green's Theorem

### Example 5

Evaluate  $\int_C (3x - 5y)dx + (x - 6y)dy$ , where  $C$  is the ellipse  $\frac{x^2}{4} + y^2 = 1$  in the anticlockwise direction. Evaluate the integral by (i) Green's Theorem, (ii) directly.

## Green's Theorem

### Example 5

Evaluate  $\int_C (3x - 5y)dx + (x - 6y)dy$ , where  $C$  is the ellipse  $\frac{x^2}{4} + y^2 = 1$  in the anticlockwise direction. Evaluate the integral by (i) Green's Theorem, (ii) directly.

### Answers

$12\pi$

## Path independence and conservative vector fields

- ▶ We considered line integrals  $\int_C \mathbf{F} \cdot d\mathbf{r}$  where  $C$  was a curve in  $\mathbb{R}^2$  now consider  $C$  as a curve in  $\mathbb{R}^3$ .

## Path independence and conservative vector fields

- ▶ We considered line integrals  $\int_C \mathbf{F} \cdot d\mathbf{r}$  where  $C$  was a curve in  $\mathbb{R}^2$  now consider  $C$  as a curve in  $\mathbb{R}^3$ .
- ▶ Two curves  $C_1$  and  $C_2$  with the same initial point  $A$  and the same end point  $B$  are *path independent* if

$$\int_{C_1} \mathbf{F} \cdot d\mathbf{r} = \int_{C_2} \mathbf{F} \cdot d\mathbf{r}$$

## Path independence and conservative vector fields

- ▶ We considered line integrals  $\int_C \mathbf{F} \cdot d\mathbf{r}$  where  $C$  was a curve in  $\mathbb{R}^2$  now consider  $C$  as a curve in  $\mathbb{R}^3$ .
- ▶ Two curves  $C_1$  and  $C_2$  with the same initial point  $A$  and the same end point  $B$  are *path independent* if

$$\int_{C_1} \mathbf{F} \cdot d\mathbf{r} = \int_{C_2} \mathbf{F} \cdot d\mathbf{r}$$

- ▶ We have path independence when we can write

$$\mathbf{F} = \nabla \phi$$

for some continuous scalar-valued function  $\phi$ .

## Path independence and conservative vector fields

- ▶ When we can find a scalar-valued function  $\phi$  such that  $\mathbf{F} = \nabla\phi$  we say that  $\mathbf{F}$  is a *conservative vector field*.

## Path independence and conservative vector fields

- ▶ When we can find a scalar-valued function  $\phi$  such that  $\mathbf{F} = \nabla\phi$  we say that  $\mathbf{F}$  is a *conservative vector field*.
- ▶ We denote  $\phi$  as the *potential function*.

## Path independence and conservative vector fields

- ▶ When we can find a scalar-valued function  $\phi$  such that  $\mathbf{F} = \nabla\phi$  we say that  $\mathbf{F}$  is a *conservative vector field*.
- ▶ We denote  $\phi$  as the *potential function*.
- ▶  $\mathbf{F}$  is conservative ensures the independence of path and gives an integral that is related to the Fundamental Theorem of Calculus which states  $\int_a^b F'(x)dx = F(b) - F(a)$ .



## Path independence and conservative vector fields

- ▶ When we can find a scalar-valued function  $\phi$  such that  $\mathbf{F} = \nabla\phi$  we say that  $\mathbf{F}$  is a *conservative vector field*.
- ▶ We denote  $\phi$  as the *potential function*.
- ▶  $\mathbf{F}$  is conservative ensures the independence of path and gives an integral that is related to the Fundamental Theorem of Calculus which states  $\int_a^b F'(x)dx = F(b) - F(a)$ .



$$\int_C \mathbf{F} \cdot d\mathbf{r} = \int_C \nabla\phi \, d\mathbf{r} = \phi(\mathbf{r}(b)) - \phi(\mathbf{r}(a)) = \phi(B) - \phi(A)$$

where  $\mathbf{r}(t)$  is the parameterised curve and the parameter  $t$  satisfies  $a \leq t \leq b$ .

## Conservative vector fields

- ▶ The grad of every smooth scalar field is a vector field. It is natural to ask whether *all* vector fields are the grad of some scalar field. In general, the answer is “no”.

## Conservative vector fields

- ▶ The grad of every smooth scalar field is a vector field. It is natural to ask whether *all* vector fields are the grad of some scalar field. In general, the answer is “no”.
- ▶ (See Example 3.10) that

$$\text{curl grad } \phi = \mathbf{0}$$

for all smooth scalar fields  $\phi$ .

## Conservative vector fields

- ▶ The grad of every smooth scalar field is a vector field. It is natural to ask whether *all* vector fields are the grad of some scalar field. In general, the answer is “no”.
- ▶ (See Example 3.10) that

$$\operatorname{curl} \operatorname{grad} \phi = \mathbf{0}$$

for all smooth scalar fields  $\phi$ .

- ▶ This means that if  $\mathbf{F} = \operatorname{grad} \phi$  for some  $\phi$  then

$$\operatorname{curl} \mathbf{F} = \mathbf{0}$$

This is a *necessary* and *sufficient* condition for  $\mathbf{F}$  to be conservative.

## Path independence

### Example 6

Vector fields  $\mathbf{V}$  and  $\mathbf{W}$  are defined by

$$\mathbf{V} = (2x - 3y + z, -3x - y + 4z, 4y + z)$$

$$\mathbf{W} = (2x - 4y - 5z, -4x + 2y, -5x + 6z).$$

One of these is conservative while the other is not. Determine which is conservative and denote it by  $\mathbf{F}$ . Find a potential function  $\phi$  for  $\mathbf{F}$  and evaluate

$$\int_C \mathbf{F} \cdot d\mathbf{r},$$

where  $C$  is the curve from  $A(1,0,0)$  to  $B(0,0,1)$  in which the plane  $x + z = 1$  cuts the hemisphere given by  $x^2 + y^2 + z^2 = 1$ ,  $y \geq 0$ .

# Path independence

## Answers

2.

## Surface integrals



Instead formulating a double integral by integrating a function  $f$  over a domain in the  $xy$ -plane we could integrate over a domain given by a surface.

# Surface integrals

Instead formulating a double integral by integrating a function  $f$  over a domain in the  $xy$ -plane we could integrate over a domain given by a surface.

Consider a crop growing on a hillside  $S$  and  $f(x, y, z)$  is the yeild per unit surface area at the point  $(x, y, z)$ . The *surface integrals* gives the total yeild of the entire crop as follows:

$$\int \int_S f(x, y, z) dS .$$





# Surface integrals

- ▶ Relate  $\delta S$  to the area of an element at the base  $\delta x \delta y$ .

# Surface integrals

- ▶ Relate  $\delta S$  to the area of an element at the base  $\delta x \delta y$ .
- ▶ The special case where the surface  $S$  can be expressed as  $\mathbf{r} = (x, y, z(x, y))$  we can use the tangent plane to approximate a small piece of surface.

## Surface integrals

- ▶ Relate  $\delta S$  to the area of an element at the base  $\delta x \delta y$ .
- ▶ The special case where the surface  $S$  can be expressed as  $\mathbf{r} = (x, y, z(x, y))$  we can use the tangent plane to approximate a small piece of surface.
- ▶ The tangent vector in the  $x$ -directions is  $\mathbf{r}_x = (1, 0, \frac{\partial z}{\partial x})$ .

## Surface integrals

- ▶ Relate  $\delta S$  to the area of an element at the base  $\delta x \delta y$ .
- ▶ The special case where the surface  $S$  can be expressed as  $\mathbf{r} = (x, y, z(x, y))$  we can use the tangent plane to approximate a small piece of surface.
- ▶ The tangent vector in the  $x$ -directions is  $\mathbf{r}_x = (1, 0, \frac{\partial z}{\partial x})$ .
- ▶ The tangent vector in the  $y$ -direction is  $\mathbf{r}_y = (0, 1, \frac{\partial z}{\partial y})$ .

## Surface integrals

- ▶ Relate  $\delta S$  to the area of an element at the base  $\delta x \delta y$ .
- ▶ The special case where the surface  $S$  can be expressed as  $\mathbf{r} = (x, y, z(x, y))$  we can use the tangent plane to approximate a small piece of surface.
- ▶ The tangent vector in the  $x$ -directions is  $\mathbf{r}_x = (1, 0, \frac{\partial z}{\partial x})$ .
- ▶ The tangent vector in the  $y$ -direction is  $\mathbf{r}_y = (0, 1, \frac{\partial z}{\partial y})$ .
- ▶ Hence,

$$\delta S \approx |\mathbf{r}_x \delta x \times \mathbf{r}_y \delta y| = |\mathbf{r}_x \times \mathbf{r}_y| \delta x \delta y = \sqrt{1 + \left(\frac{\partial z}{\partial x}\right)^2 + \left(\frac{\partial z}{\partial y}\right)^2} \delta x \delta y$$

## Surface integrals

- ▶ Relate  $\delta S$  to the area of an element at the base  $\delta x \delta y$ .
- ▶ The special case where the surface  $S$  can be expressed as  $\mathbf{r} = (x, y, z(x, y))$  we can use the tangent plane to approximate a small piece of surface.
- ▶ The tangent vector in the  $x$ -directions is  $\mathbf{r}_x = (1, 0, \frac{\partial z}{\partial x})$ .
- ▶ The tangent vector in the  $y$ -direction is  $\mathbf{r}_y = (0, 1, \frac{\partial z}{\partial y})$ .
- ▶ Hence,

$$\delta S \approx |\mathbf{r}_x \delta x \times \mathbf{r}_y \delta y| = |\mathbf{r}_x \times \mathbf{r}_y| \delta x \delta y = \sqrt{1 + \left(\frac{\partial z}{\partial x}\right)^2 + \left(\frac{\partial z}{\partial y}\right)^2} \delta x \delta y$$

- ▶ The surface integrals becomes:

$$\iint_S f(x, y, z) dS = \iint_D f(x, y, z) \sqrt{1 + \left(\frac{\partial z}{\partial x}\right)^2 + \left(\frac{\partial z}{\partial y}\right)^2} dx dy.$$

where  $D$  is the projection of  $S$  onto the  $xy$ -plane.

# Surface integrals

## Example 6

Evaluate

$$\iint_S z^2 \, dS$$

where  $S$  is the hemisphere given by  $x^2 + y^2 + z^2 = 1$  with  $z \geq 0$ .

# Surface integrals


## Example 6

Evaluate

$$\int \int_S z^2 dS$$

where  $S$  is the hemisphere given by  $x^2 + y^2 + z^2 = 1$  with  $z \geq 0$ .

Answers





# Surface integrals

## Example 6


Evaluate

$$\int \int_S z^2 dS$$

where  $S$  is the hemisphere given by  $x^2 + y^2 + z^2 = 1$  with  $z \geq 0$ .

Answers

$$2\pi/3$$



## Surface integrals

A surface integral can also be used to calculate the area of a surface  $S$ .

$$\int \int_S 1 \, dS = \text{Area of surface } S$$

# Surface integrals

## Example 7


Find the area of the ellipse cut on the plane  $2x + 3y + 6z = 60$  by the circular cylinder  $x^2 + y^2 = 2x$ .

# Surface integrals

## Example 7

Find the area of the ellipse cut on the plane  $2x + 3y + 6z = 60$  by the circular cylinder  $x^2 + y^2 = 2x$ .

## Answers




# Surface integrals

## Example 7

Find the area of the ellipse cut on the plane  $2x + 3y + 6z = 60$  by the circular cylinder  $x^2 + y^2 = 2x$ .

## Answers

$$7\pi/6$$




## Surface integrals of vector fields

- ▶ To define *surface integrals of vector fields*, we need to consider *orientable surfaces* (2-sided).

## Surface integrals of vector fields

- ▶ To define *surface integrals of vector fields*, we need to consider *orientable surfaces* (2-sided).
- ▶ The *normal* to the surface gives the surface orientation. So there are two possible orientations for any orientable surface.



## Surface integrals of vector fields

- ▶ For a surface in the form  $f(x, y, z) = 0$  the *normal vector* is given by

$$\mathbf{n} = \left( \frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}, \frac{\partial f}{\partial z} \right)$$



## Surface integrals of vector fields

- ▶ For a surface in the form  $f(x, y, z) = 0$  the *normal vector* is given by

$$\mathbf{n} = \left( \frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}, \frac{\partial f}{\partial z} \right)$$

- ▶ For a surface in the form  $z = z(x, y)$  the *normal vector* is given by

$$\mathbf{n} = \left( \frac{\partial z}{\partial x}, \frac{\partial z}{\partial y}, -1 \right)$$

This one follows from the fact that  $\mathbf{r}_x \times \mathbf{r}_y$  is normal to the vectors  $\mathbf{r}_x$  and  $\mathbf{r}_y$  which lie in the tangent plane

# Surface integrals of vector fields

## Examples

- ▶ The normal to the plane  $f(x, y, z) = 2x + 7y + 3z - 50 = 0$  is:

$$\mathbf{n} = \left( \frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}, \frac{\partial f}{\partial z} \right) = (2, 7, 3)$$

# Surface integrals of vector fields

## Examples

- ▶ The normal to the plane  $f(x, y, z) = 2x + 7y + 3z - 50 = 0$  is:

$$\mathbf{n} = \left( \frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}, \frac{\partial f}{\partial z} \right) = (2, 7, 3)$$

- ▶ For the sphere  $x^2 + y^2 + z^2 - a^2 = 0$ , the normal is,  $(2x, 2y, 2z)$

## Surface integrals of vector fields

Imagine a fluid flowing through a surface  $S$ .  $\mathbf{F}$  is the force field and it is related to the velocity and density of the fluid flowing through the surface.

## Surface integrals of vector fields

Imagine a fluid flowing through a surface  $S$ .  $\mathbf{F}$  is the force field and it is related to the velocity and density of the fluid flowing through the surface.

A measure of the total flux (flow) across the surface is given by

$$\int \int_S \mathbf{F} \cdot \mathbf{n} \, dS,$$

where  $\mathbf{n}$  is the *unit normal*.

## Surface integrals of vector fields

Imagine a fluid flowing through a surface  $S$ .  $\mathbf{F}$  is the force field and it is related to the velocity and density of the fluid flowing through the surface.

A measure of the total flux (flow) across the surface is given by

$$\int \int_S \mathbf{F} \cdot \mathbf{n} \, dS,$$

where  $\mathbf{n}$  is the *unit normal*.

$\mathbf{F} \cdot \mathbf{n} \, dS$  tells us the mass of fluid flowing across a region  $dS$  in the direction of  $\mathbf{n}$ .

# Surface integrals of vector fields

- ▶ Gauss' Divergence Theorem will help us calculate

$$\iint_S \mathbf{F} \cdot \mathbf{n} \, dS$$

## Surface integrals of vector fields

- ▶ Gauss' Divergence Theorem will help us calculate

$$\int \int_S \mathbf{F} \cdot \mathbf{n} \, dS$$

- ▶ There are some similarities between Green's Theorem on  $\mathbb{R}^2$  and Gauss' Divergence Theorem in  $\mathbb{R}^3$ .



## Surface integrals of vector fields

- ▶ Gauss' Divergence Theorem will help us calculate

$$\int \int_S \mathbf{F} \cdot \mathbf{n} \, dS$$

- ▶ There are some similarities between Green's Theorem on  $\mathbb{R}^2$  and Gauss' Divergence Theorem in  $\mathbb{R}^3$ .

**Theorem:** Line integral round a boundary curve  $C$  of a *closed* region in  $\mathbb{R}^2$  = Double integral over the *enclosed* 2-dimensional region.

## Surface integrals of vector fields

- ▶ Gauss' Divergence Theorem will help us calculate

$$\int \int_S \mathbf{F} \cdot \mathbf{n} \, dS$$

- ▶ There are some similarities between Green's Theorem on  $\mathbb{R}^2$  and Gauss' Divergence Theorem in  $\mathbb{R}^3$ .

**Theorem:** Line integral round a boundary curve  $C$  of a *closed* region in  $\mathbb{R}^2$  = Double integral over the *enclosed* 2-dimensional region.

**Theorem:** Surface integral over a boundary surface  $S$  of a *closed* region in  $\mathbb{R}^3$  = Triple integral over the *enclosed* 3-dimensional region.

## Surface integrals of vector fields

- ▶ Gauss' Divergence Theorem will help us calculate

$$\int \int_S \mathbf{F} \cdot \mathbf{n} \, dS$$

- ▶ There are some similarities between Green's Theorem on  $\mathbb{R}^2$  and Gauss' Divergence Theorem in  $\mathbb{R}^3$ .

**Theorem:** Line integral round a boundary curve  $C$  of a *closed* region in  $\mathbb{R}^2$  = Double integral over the *enclosed* 2-dimensional region.

**Theorem:** Surface integral over a boundary surface  $S$  of a *closed* region in  $\mathbb{R}^3$  = Triple integral over the *enclosed* 3-dimensional region.

## Surface integrals of vector fields

- ▶ Gauss' Divergence Theorem will help us calculate

$$\int \int_S \mathbf{F} \cdot \mathbf{n} \, dS$$

- ▶ There are some similarities between Green's Theorem on  $\mathbb{R}^2$  and Gauss' Divergence Theorem in  $\mathbb{R}^3$ .

**Theorem:** Line integral round a boundary curve  $C$  of a *closed* region in  $\mathbb{R}^2$  = Double integral over the *enclosed* 2-dimensional region.

**Theorem:** Surface integral over a boundary surface  $S$  of a *closed* region in  $\mathbb{R}^3$  = Triple integral over the *enclosed* 3-dimensional region.

## Gauss' Divergence Theorem

Let  $V$  be a closed bounded volume on  $\mathbb{R}^3$  with boundary surface  $S$ , given with positive (*outward*) orientation. Let  $\mathbf{F}$  be a vector field whose component functions have continuous partial derivatives on an open region containing  $V$ . Then

$$\int \int_S \mathbf{F} \cdot \mathbf{n} \, dS = \int \int \int_V \operatorname{div} \mathbf{F} \, dx \, dy \, dz ,$$

where  $\mathbf{n}$  denotes the outward pointing *unit normal* at each point on the surface  $S$ .

# Divergence Theorem

## Example 8

Use Gauss' Divergence Theorem to evaluate

$$I = \int \int_S x^4 y + y^2 z^2 + xz^2 \, dS,$$

where  $S$  is the entire surface of the sphere  $x^2 + y^2 + z^2 = 1$ .

# Divergence Theorem

## Example 8

Use Gauss' Divergence Theorem to evaluate

$$I = \int \int_S x^4 y + y^2 z^2 + xz^2 \, dS,$$

where  $S$  is the entire surface of the sphere  $x^2 + y^2 + z^2 = 1$ .

## Answers

$$4\pi/15$$

# Divergence Theorem

## Example 9

Find  $I = \int \int_S \mathbf{F} \cdot \mathbf{n} \, dS$  where  $\mathbf{F} = (2x, 2y, 1)$  and where  $S$  is the entire surface consisting of  $S_2$  = the part of the paraboloid  $z = 1 - x^2 - y^2$  with  $z = 0$  together with  $S_1$  = disc  $\{(x, y) : x^2 + y^2 \leq 1\}$ . Here  $\mathbf{n}$  is the outward pointing unit normal.




# Divergence Theorem

## Example 9

Find  $I = \int \int_S \mathbf{F} \cdot \mathbf{n} \, dS$  where  $\mathbf{F} = (2x, 2y, 1)$  and where  $S$  is the entire surface consisting of  $S_2$  = the part of the paraboloid  $z = 1 - x^2 - y^2$  with  $z = 0$  together with  $S_1$  = disc  $\{(x, y) : x^2 + y^2 \leq 1\}$ . Here  $\mathbf{n}$  is the outward pointing unit normal.

## Answers



# Divergence Theorem

## Example 9

Find  $I = \int \int_S \mathbf{F} \cdot \mathbf{n} \, dS$  where  $\mathbf{F} = (2x, 2y, 1)$  and where  $S$  is the entire surface consisting of  $S_2$  = the part of the paraboloid  $z = 1 - x^2 - y^2$  with  $z = 0$  together with  $S_1$  = disc  $\{(x, y) : x^2 + y^2 \leq 1\}$ . Here  $\mathbf{n}$  is the outward pointing unit normal.

## Answers

$2\pi$

